

Mack's chain-ladder model in Lean 4: blueprint

Robert Sneiderman

August 25, 2026

Contents

1	Introduction	2
2	The chain-ladder estimators	3
2.1	Definitions	3
2.2	Deterministic identities	3
3	Mack 1999 equals Mack 1993	5
4	Bornhuetter-Ferguson on the chain-ladder pattern	6
5	The stochastic layer	7
5.1	Model	7
5.2	Mack's Theorem 2	7
5.3	Mack's Theorem 1	8
5.4	The variance assumptions and the estimation variance of $\hat{f}_k$	8
5.5	Unbiasedness of the variance estimator	9
5.6	Process variance and the exact MSEP (Theorem 3)	9
5.7	From independence across accident years to the $\mathcal{D}_k$ form	10
6	The variant catalogue: Mack versus conditional resampling	11
6.1	The total reserve: Mack's Corollary	12
7	Solvency II: the one-year view	13
7.1	The one-year claims development result	13
8	Röhr 2016: the error-propagation form	15
8.1	Definitions	15
8.2	The split, and the two Mack plug-ins	16
8.3	Catalogue row 3	16
8.4	The total reserve	17
8.5	The total reserve: the exact conditional MSEP	17
9	Non-vacuity: a nondegenerate model	19
10	Reference computation	20

Chapter 1

Introduction

This blueprint tracks the formalization of Mack’s distribution-free chain-ladder model [1] in Lean 4 on top of mathlib. Each statement below carries the name of its Lean declaration; green nodes in the dependency graph are proved, the others are open work listed in the roadmap. Statements are given in the paper’s own terms, and where a published formula is an estimator or an approximation rather than an identity, the text says so.

A run-off triangle with n accident years and n development years is a function $C : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{R}$; $C_{i,k}$ is the cumulative claims of accident year i at development year k , observed when $i+k \leq n-1$. The latest observed entry of accident year i is $C_{i,n-1-i}$.

Chapter 2

The chain-ladder estimators

2.1 Definitions

Definition 2.1 (Contributors). The accident years contributing to the development factor $k \rightarrow k+1$ are $i = 0, \dots, n-k-2$, those with $C_{i,k+1}$ observed.

Definition 2.2 (Column sums). $S_k = \sum_{i=0}^{n-k-2} C_{i,k}$ and $T_k = \sum_{i=0}^{n-k-2} C_{i,k+1}$.

Definition 2.3 (Development factor). $\hat{f}_k = T_k/S_k$. (In Lean, division by zero is zero.)

Definition 2.4 (Individual factor). $F_{i,k} = C_{i,k+1}/C_{i,k}$.

Definition 2.5 (Mack's variance estimator). $\hat{\sigma}_k^2 = \frac{1}{n-k-2} \sum_{i=0}^{n-k-2} C_{i,k} (F_{i,k} - \hat{f}_k)^2$. The value for $k = n-2$ (a single contributor) is an extrapolation convention in [1], not an estimator; no theorem here uses it.

Definition 2.6 (Projection). $\hat{C}_{i,k} = C_{i,n-1-i} \prod_{j=n-1-i}^{k-1} \hat{f}_j$ for $k \geq n-1-i$.

Definition 2.7 (Ultimate and reserve). $\hat{C}_{i,n-1}$ is the chain-ladder ultimate and $\hat{R}_i = \hat{C}_{i,n-1} - C_{i,n-1-i}$ the reserve.

Definition 2.8 (Mack's MSEP estimator). Mack (1993), Theorem 3:

$$\widehat{\text{mse}}(\hat{R}_i) = \hat{C}_{i,n-1}^2 \sum_{k=n-1-i}^{n-2} \frac{\hat{\sigma}_k^2}{\hat{f}_k^2} \left(\frac{1}{\hat{C}_{i,k}} + \frac{1}{S_k} \right).$$

2.2 Deterministic identities

Lemma 2.9 (Weighted-average form). If $C_{i,k} \neq 0$ for every contributor i , then $\hat{f}_k = \sum_i \frac{C_{i,k}}{S_k} F_{i,k}$.

Proof. Termwise: $\frac{C_{i,k}}{S_k} \cdot \frac{C_{i,k+1}}{C_{i,k}} = \frac{C_{i,k+1}}{S_k}$. □

Lemma 2.10. If $C_{i,k} \neq 0$ for every contributor, $T_k = \sum_i C_{i,k} F_{i,k}$.

Proof. Cancel the denominator termwise. □

Lemma 2.11 (Sum-of-squares decomposition). For any centre f , with $C_{i,k} \neq 0$ for every contributor, $\sum_i C_{i,k} (F_{i,k} - f)^2 = \sum_i C_{i,k} F_{i,k}^2 - 2fT_k + f^2S_k$.

Proof. Expand the square and use Lemma 2.10. □

Lemma 2.12 (Collapse at the chain-ladder centre). *If moreover $S_k \neq 0$, then $\sum_i C_{i,k}(F_{i,k} - \hat{f}_k)^2 = \sum_i C_{i,k}F_{i,k}^2 - S_k\hat{f}_k^2$. This is the identity behind the $n - k - 2$ degrees of freedom in $\hat{\sigma}_k^2$.*

Proof. Put $f = \hat{f}_k = T_k/S_k$ in Lemma 2.11. □

Lemma 2.13 (Projection step). *For $k \geq n - 1 - i$, $\hat{C}_{i,k+1} = \hat{C}_{i,k} \hat{f}_k$.*

Proof. Split off the top factor of the product. □

Lemma 2.14. $\hat{C}_{i,n-1-i} = C_{i,n-1-i}$.

Proof. Empty product. □

Lemma 2.15. *The oldest accident year is fully developed: $\hat{R}_0 = 0$.*

Proof. Empty product again. □

Chapter 3

Mack 1999 equals Mack 1993

Definition 3.1 (Mack's 1999 recursion). For $\alpha = 1$ and unit weights [2], starting from $\text{se}^2 = 0$ on the latest diagonal $d = n - 1 - i$ and stepping m times,

$$\text{se}_{m+1}^2 = \hat{C}_{i,d+m}^2 \left(\frac{\hat{\sigma}_{d+m}^2}{\hat{C}_{i,d+m}} + \frac{\hat{\sigma}_{d+m}^2}{S_{d+m}} \right) + \text{se}_m^2 \hat{f}_{d+m}^2.$$

Definition 3.2. The summand of the closed form: $\text{term}_{i,k} = \frac{\hat{\sigma}_k^2}{\hat{f}_k^2} \left(\frac{1}{\hat{C}_{i,k}} + \frac{1}{S_k} \right)$.

Lemma 3.3. $\widehat{\text{mse}}(\hat{R}_i) = \hat{C}_{i,n-1}^2 \sum_{k=n-1-i}^{n-2} \text{term}_{i,k}$.

Proof. By definition. □

Lemma 3.4 (Truncated closed form). If $\hat{f}_k \neq 0$ for $d \leq k < d+m$, then $\text{se}_m^2 = \hat{C}_{i,d+m}^2 \sum_{k=d}^{d+m-1} \text{term}_{i,k}$.

Proof. Induction on m using Lemma 2.13 and $\hat{f}_{d+m}^2 / \hat{f}_{d+m}^2 = 1$. □

Theorem 3.5 (Mack 1999 = Mack 1993). For $i \leq n - 1$ and $\hat{f}_k \neq 0$ along the row, i steps of the 1999 recursion return exactly Mack's 1993 estimator: $\text{se}_i^2 = \widehat{\text{mse}}(\hat{R}_i)$.

Proof. Lemma 3.4 with $m = i$, since $d + i = n - 1$. □

Chapter 4

Bornhuetter-Ferguson on the chain-ladder pattern

Definition 4.1 (Cumulative development factor, BF reserve and ultimate). $\text{cdf}_i = \prod_{k=d}^{n-2} \hat{f}_k$; for an a priori ultimate U , $R_i^{BF} = U(1 - 1/\text{cdf}_i)$ and $\hat{C}_{i,n-1}^{BF} = C_{i,d} + R_i^{BF}$.

Lemma 4.2 (Linearity). $R_i^{BF}(aU) = a R_i^{BF}(U)$ and $R_i^{BF}(U + V) = R_i^{BF}(U) + R_i^{BF}(V)$.

Proof. Ring identities. □

Theorem 4.3 (BF with the chain-ladder prior is chain ladder). *If $\text{cdf}_i \neq 0$ and $U = \hat{C}_{i,n-1}$, then $\hat{C}_{i,n-1}^{BF} = \hat{C}_{i,n-1}$ and $R_i^{BF} = \hat{R}_i$.*

Proof. $\hat{C}_{i,n-1} = C_{i,d} \text{cdf}_i$, so $U(1 - 1/\text{cdf}_i) = C_{i,d} \text{cdf}_i - C_{i,d}$. □

Lemma 4.4 (The BF reserve is a fraction of the prior). *If every development factor along the row is at least 1, then $\text{cdf}_i \geq 1$ and, for $U \geq 0$, $0 \leq R_i^{BF}(U) \leq U$.*

Proof. A product of numbers at least 1 is at least 1; then $0 \leq 1 - 1/\text{cdf}_i \leq 1$. □

Chapter 5

The stochastic layer

5.1 Model

Definition 5.1 (Random triangle). A random run-off triangle on a measurable space Ω is a family of random variables $C_{i,k} : \Omega \rightarrow \mathbb{R}$ together with a filtration $\mathcal{D}_k$ such that $C_{i,j}$ is $\mathcal{D}_k$ -measurable whenever $j \leq k$ ("everything observed up to development year k ").

Definition 5.2 (Assumption (M1)). $E[C_{i,k+1} \mid \mathcal{D}_k] = f_k C_{i,k}$ almost surely, for all i and k . This is Mack's first assumption in the form conditioned on $\mathcal{D}_k$; Mack states it conditioned on the row's own history and obtains this form from independence across accident years (M2). That passage is Theorem 5.30.

Definition 5.3 (Estimators as random variables). $S_k(\omega)$ and $\hat{f}_k(\omega)$ are the deterministic estimators applied to the triangle observed at ω .

Lemma 5.4. S_k is $\mathcal{D}_k$ -measurable.

Proof. A finite sum of $\mathcal{D}_k$ -measurable functions. □

5.2 Mack's Theorem 2

Theorem 5.5 (Conditional unbiasedness of the development factor). Assume (M1), that $S_k \neq 0$ almost surely, that each $C_{i,k+1}$ and $\hat{f}_k$ are integrable. Then $E[\hat{f}_k \mid \mathcal{D}_k] = f_k$ almost surely.

Proof. $\hat{f}_k = S_k^{-1} \sum_i C_{i,k+1}$. The factor S_k^{-1} is $\mathcal{D}_k$ -measurable and comes out of the conditional expectation; by (M1) the conditional expectation of the sum is $f_k \sum_i C_{i,k} = f_k S_k$; cancel S_k where it is nonzero. □

Lemma 5.6. $\hat{f}_j$ is $\mathcal{D}_k$ -measurable whenever $j + 1 \leq k$.

Proof. S_j is $\mathcal{D}_j \subseteq \mathcal{D}_k$ -measurable and each $C_{i,j+1}$ is $\mathcal{D}_k$ -measurable. □

Theorem 5.7 (Uncorrelated development factors, conditional form). Under (M1), on a finite measure space, for $j < k$ (with the integrability and $S \neq 0$ hypotheses of Theorem 5.5 for both columns and $\hat{f}_j \hat{f}_k$ integrable), $E[\hat{f}_j \hat{f}_k \mid \mathcal{D}_j] = f_j f_k$ almost surely.

Proof. Condition on $\mathcal{D}_k$ first: $\hat{f}_j$ is $\mathcal{D}_k$ -measurable and comes out, and $E[\hat{f}_k \mid \mathcal{D}_k] = f_k$ by Theorem 5.5. Then the tower property down to $\mathcal{D}_j$ and Theorem 5.5 again for column j . □

Theorem 5.8 (Mack 1993, Theorem 2). *On a probability space, under the same hypotheses, $E[\hat{f}_k] = f_k$ and $E[\hat{f}_j \hat{f}_k] = f_j f_k$ for $j < k$: the chain-ladder factors are unbiased and uncorrelated, which is what makes $\prod_k \hat{f}_k$ an unbiased estimator of $\prod_k f_k$.*

Proof. Integrate the conditional statements. $\square$

5.3 Mack's Theorem 1

Definition 5.9 (Chain-ladder projection as a random variable). $\hat{C}_{i,d+m} = C_{i,d} \prod_{k=d}^{d+m-1} \hat{f}_k$ with $d = n - 1 - i$.

Lemma 5.10. $\hat{C}_{i,d+m}$ is $\mathcal{D}_{d+m}$ -measurable.

Proof. Induction on m ; each new factor $\hat{f}_{d+m}$ is $\mathcal{D}_{d+m+1}$ -measurable. $\square$

Lemma 5.11 (Iterating (M1)). *For $d \leq k$, $E[C_{i,k+1} \mid \mathcal{D}_d] = f_k E[C_{i,k} \mid \mathcal{D}_d]$, hence $E[C_{i,d+m} \mid \mathcal{D}_d] = C_{i,d} \prod_{k=d}^{d+m-1} f_k$.*

Proof. Tower property through $\mathcal{D}_k$, then (M1); induction on m . $\square$

Theorem 5.12 (Mack 1993, Theorem 1). *Under (M1), with the partial products integrable and the column sums a.s. nonzero, $E[\hat{C}_{i,d+m} \mid \mathcal{D}_d] = C_{i,d} \prod_{k=d}^{d+m-1} f_k$, and therefore $E[\hat{C}_{i,n-1} \mid \mathcal{D}_d] = E[C_{i,n-1} \mid \mathcal{D}_d]$: the chain-ladder ultimate is a conditionally unbiased estimator of the true ultimate claims.*

Proof. Condition on $\mathcal{D}_{d+m}$: the partial product comes out and the last factor has conditional expectation f_{d+m} (Theorem 5.5); tower down to $\mathcal{D}_d$ and induct. Compare with Lemma 5.11. $\square$

5.4 The variance assumptions and the estimation variance of $\hat{f}_k$

Definition 5.13 (Residuals, (M3) and (M2')). With $\varepsilon_{i,k} = C_{i,k+1} - f_k C_{i,k}$: (M3) $E[\varepsilon_{i,k}^2 \mid \mathcal{D}_k] = \sigma_k^2 C_{i,k}$, and (M2') $E[\varepsilon_{i,k} \varepsilon_{j,k} \mid \mathcal{D}_k] = 0$ for $i \neq j$, the conditional uncorrelatedness across accident years that independence of accident years supplies.

Lemma 5.14. *On $\{S_k \neq 0\}$, $\hat{f}_k - f_k = S_k^{-1} \sum_i \varepsilon_{i,k}$ and $(\hat{f}_k - f_k)^2 = S_k^{-2} \sum_i \sum_j \varepsilon_{i,k} \varepsilon_{j,k}$.*

Proof. $T_k - f_k S_k = \sum_i \varepsilon_{i,k}$; expand the square of the sum. $\square$

Theorem 5.15 (Estimation variance of the development factor). *Under (M3) and (M2'), with the residual products integrable, on $\{S_k \neq 0\}$, $E[(\hat{f}_k - f_k)^2 \mid \mathcal{D}_k] = \sigma_k^2 / S_k$ (Mack 1993, proof of Theorem 3; Mack 1999, s.e. $(\hat{f}_k)^2 = \hat{\sigma}_k^2 / S_k$).*

Proof. S_k^{-2} is $\mathcal{D}_k$ -measurable and comes out; conditional expectation is linear over the double sum; cross terms vanish by (M2'); diagonal terms give $\sigma_k^2 \sum_i C_{i,k} = \sigma_k^2 S_k$. $\square$

5.5 Unbiasedness of the variance estimator

Lemma 5.16 (Sum of squares around $\hat{f}_k$ via residuals). *With all contributing $C_{i,k} \neq 0$ and $S_k \neq 0$, for any f , $\sum_i C_{i,k}(F_{i,k} - \hat{f}_k)^2 = \sum_i \varepsilon_i^2 / C_{i,k} - S_k(\hat{f}_k - f)^2$ where $\varepsilon_i = C_{i,k+1} - fC_{i,k}$.*

Proof. Two instances of Lemma 2.11 (centres $\hat{f}_k$ and f) and $T_k = \hat{f}_k S_k$. $\square$

Definition 5.17. $\hat{\sigma}_k^2$ and the weighted sum of squares $W_k = \sum_i C_{i,k}(F_{i,k} - \hat{f}_k)^2$ as random variables; $\hat{\sigma}_k^2 = W_k / (n - k - 2)$.

Theorem 5.18 (Unbiasedness of the variance estimator). *Under (M3) and (M2'), with all contributing $C_{i,k}$ and S_k a.s. nonzero and the stated integrability, $E[W_k \mid \mathcal{D}_k] = (n - k - 2) \sigma_k^2$ for $k + 2 \leq n$, and hence $E[\hat{\sigma}_k^2 \mid \mathcal{D}_k] = \sigma_k^2$ for $k + 3 \leq n$.*

Proof. By Lemma 5.16, $W_k = \sum_i \varepsilon_i^2 / C_{i,k} - S_k(\hat{f}_k - f_k)^2$. Each $C_{i,k}^{-1}$ is $\mathcal{D}_k$ -measurable, so (M3) gives σ_k^2 per term, $n - k - 1$ terms in all; S_k is $\mathcal{D}_k$ -measurable and Theorem 5.15 gives σ_k^2 for the subtracted term. $\square$

5.6 Process variance and the exact MSEP (Theorem 3)

Lemma 5.19 (Conditional second moment). *Under (M1) and (M3), $E[\varepsilon_{i,k} \mid \mathcal{D}_k] = 0$, $E[C_{i,k+1}^2 \mid \mathcal{D}_k] = f_k^2 C_{i,k}^2 + \sigma_k^2 C_{i,k}$, and for $d \leq k$, $E[C_{i,k+1}^2 \mid \mathcal{D}_d] = f_k^2 E[C_{i,k}^2 \mid \mathcal{D}_d] + \sigma_k^2 E[C_{i,k} \mid \mathcal{D}_d]$.*

Proof. Expand $C_{i,k+1} = f_k C_{i,k} + \varepsilon_{i,k}$; the cross term vanishes; tower property. $\square$

Definition 5.20 (Process variance). $V_0 = 0$, $V_{m+1} = f_{d+m}^2 V_m + \sigma_{d+m}^2 c \prod_{l < m} f_{d+l}$; in closed form $V_m = \sum_{j < m} \left(\prod_{l=j+1}^{m-1} f_{d+l} \right)^2 \sigma_{d+j}^2 c \prod_{l < j} f_{d+l}$.

Theorem 5.21 (Process variance along a row). *Under (M1) and (M3), $E[C_{i,d+m}^2 \mid \mathcal{D}_d] - E[C_{i,d+m} \mid \mathcal{D}_d]^2 = V_m$ with $c = C_{i,d}$.*

Proof. Induction on m with Lemma 5.19 and Lemma 5.11. $\square$

Lemma 5.22 (Conditional MSEP decomposition). *For a $\mathcal{D}$ -measurable predictor P of Y , $E[(P - Y)^2 \mid \mathcal{D}] = (E[Y^2 \mid \mathcal{D}] - E[Y \mid \mathcal{D}]^2) + (P - E[Y \mid \mathcal{D}])^2$.*

Proof. Expand the square; P comes out of the conditional expectation. $\square$

Theorem 5.23 (Mack's Theorem 3, exact form). *Let $\mathcal{D}$ be the observed data. Assume $\hat{C}_{i,d+m}$ is $\mathcal{D}$ -measurable and that $\mathcal{D}$ adds nothing to $\mathcal{D}_d$ about row i 's future (the conditional mean and second moment of $C_{i,d+m}$ given $\mathcal{D}$ agree with those given $\mathcal{D}_d$; this is what independence across accident years supplies). Then under (M1) and (M3)*

$$E[(\hat{C}_{i,d+m} - C_{i,d+m})^2 \mid \mathcal{D}] = V_m + (\hat{C}_{i,d+m} - C_{i,d} \prod_{l < m} f_{d+l})^2.$$

Proof. Lemma 5.22 with $P = \hat{C}_{i,d+m}$, then Theorem 5.21 and Lemma 5.11. $\square$

Definition 5.24 (Mack's estimators of the two terms). $\hat{V}$ is V_m with $\hat{f}, \hat{\sigma}^2$ plugged in. The estimation-error term $\hat{C}_{i,n-1}^2 \sum_k \hat{\sigma}_k^2 / (\hat{f}_k^2 S_k)$ is obtained from $(\hat{C} - M)^2$ by Mack's conditional-resampling step: replace $(\hat{f}_k - f_k)^2$ by σ_k^2 / S_k (Theorem 5.15), drop the cross terms (Theorem 5.7), and plug in $\hat{f}, \hat{\sigma}^2$. This is a definition, not a theorem: it is the approximation in Mack (1993), and the exact conditional MSEP above is the object it approximates (Buchwalder, Bühlmann, Merz and Wüthrich 2006 estimate the same object differently).

5.7 From independence across accident years to the $\mathcal{D}_k$ form

Mack states (M1) and (M3) conditioned on one accident year's own history, and assumes separately that the accident years are independent (M2). Every theorem above conditions on $\mathcal{D}_k$ instead. This section proves the passage.

Definition 5.25 (Row σ -algebras). $\sigma(C_{i,0}, \dots, C_{i,k})$, the history of accident year i up to development year k ; $\sigma(C_{i,j} : j)$, the whole of accident year i ; and the joins of these over the other accident years $i' \neq i$.

Definition 5.26 ((M2) and the row-conditioned assumptions). (M2) is **RowsIndep**: the σ -algebras of the individual accident years are independent. **RowsGenerated** says that $\mathcal{D}_k$ is what its name means, the join of the rows' histories up to development year k . (M1row) and (M3row) are Mack's assumptions conditioned on $\sigma(C_{i,0}, \dots, C_{i,k})$ rather than on $\mathcal{D}_k$.

Lemma 5.27 (Rectangles). *If m'_1 and m_2 are independent, F is m'_1 -measurable and integrable, $a \in m'_1$ and $b \in m_2$, then $\int_{a \cap b} F = \mu(b) \int_a F$.*

Proof. Apply `mathlib's condExp_indep_eq` to the indicator $1_a F$, which is m'_1 -measurable, and integrate the resulting constant over b . $\square$

Theorem 5.28 (Conditioning on an independent enlargement changes nothing). *Let $m_1 \leq m'_1$, let f be m'_1 -measurable and integrable, and let m_2 be independent of m'_1 . Then $E[f \mid m_1 \vee m_2] = E[f \mid m_1]$ almost surely.*

Proof. The sets $a \cap b$ with $a \in m_1$, $b \in m_2$ form a π -system generating $m_1 \vee m_2$. On such a set both $\int f$ and $\int E[f \mid m_1]$ factor by Lemma 5.27 into $\mu(b)$ times the integral over a , and those agree by the defining property of $E[f \mid m_1]$. The class of sets on which the two set integrals agree is closed under complements and countable disjoint unions, so it is all of $m_1 \vee m_2$; conclude by the uniqueness of the conditional expectation. Mathlib has the case $m_1 = \perp$ only. $\square$

Lemma 5.29 (One row against the others). *Under (M2), accident year i is independent of all the other accident years taken together, and so of their observed histories. Under **RowsGenerated**, $\mathcal{D}_k$ is the join of row i 's history up to k with the other rows' histories up to k .*

Proof. `indep_iSup_of_disjoint` for $\{i\}$ against its complement, then monotonicity of independence; the splitting of $\mathcal{D}_k$ is `iSup_split_single`. $\square$

Theorem 5.30 ((M1) and (M3) in the $\mathcal{D}_k$ form). *Assume (M2), that $\mathcal{D}_k$ is generated by the rows' histories, and the integrability of the entries. Then Mack's row-conditioned (M1row) implies (M1), and (M3row) implies (M3).*

Proof. Split $\mathcal{D}_k$ by Lemma 5.29, apply Theorem 5.28 with m_1 the row's history up to k , m'_1 the whole row and m_2 the other rows, and finish with the row-conditioned assumption. $\square$

Theorem 5.31 ((M2') is a consequence, not an assumption). *Under (M2), **RowsGenerated** and (M1row), the residuals of two different accident years are conditionally uncorrelated given $\mathcal{D}_k$: assumption (M2'), which the variance theorems take as a hypothesis, follows from independence across accident years.*

Proof. Condition on $\mathcal{D}_k$ together with the whole of accident year j : row j 's residual comes out of the conditional expectation, and what is left is row i 's residual conditioned on a σ -algebra that adds nothing about row i , so it equals $E[\varepsilon_{i,k} \mid \sigma(C_{i,0}, \dots, C_{i,k})] = 0$ by Theorem 5.28 and (M1row). The tower property brings the identity back to $\mathcal{D}_k$. $\square$

Chapter 6

The variant catalogue: Mack versus conditional resampling

Write $a_k = \hat{\sigma}_k^2 / (\hat{f}_k^2 S_k)$ for the relative estimation variance of development factor k along the row. Mack's estimation-error term is $\hat{C}_{i,n-1}^2 \sum_k a_k$; the conditional-resampling term of Buchwalder, Bühlmann, Merz and Wüthrich (2006), which Murphy (1994) also reaches, is $\hat{C}_{i,n-1}^2 (\prod_k (1 + a_k) - 1)$.

Definition 6.1. a_k and the conditional-resampling estimator $\hat{C}_{i,n-1}^2 (\prod_k (1 + a_k) - 1)$.

Lemma 6.2 (Product minus sum). *For $a_k \geq 0$: $1 + \sum a_k \leq \prod (1 + a_k) \leq \exp(\sum a_k)$; for two terms $(1 + a)(1 + b) - 1 - (a + b) = ab$; and $\prod (1 + a_k) - 1 - \sum a_k = 0$ when at most one a_k is nonzero.*

Proof. Induction over the index set for the Weierstrass inequality; $1 + x \leq e^x$ termwise for the exponential bound. $\square$

Theorem 6.3 (Catalogue row: Mack is the first-order part of BMW). *With $a_k \geq 0$ along the row: (i) Mack $\leq$ BMW; (ii) the difference is exactly $\hat{C}_{i,n-1}^2 (\prod_k (1 + a_k) - 1 - \sum_k a_k)$; (iii) the two coincide for the second-latest accident year, which has a single development factor; (iv) the difference is at most $\hat{C}_{i,n-1}^2 (e^{\sum a_k} - 1 - \sum a_k)$, second order in the relative estimation variances.*

Proof. Lemma 6.2 scaled by $\hat{C}_{i,n-1}^2$. $\square$

Theorem 6.4 (Strict counterexample). *There is a run-off triangle (four accident years, four development years, with unequal individual factors in the first two columns) and an accident year on which Mack's estimation-error term is strictly smaller than the conditional-resampling term: the two are different estimators, not two presentations of one. On the exhibited triangle $a_0 = 1/297$, $a_1 = 1/2809$, and the remainder is $\hat{C}^2 a_0 a_1 > 0$. Every number is checked by `norm_num` from the definitions.*

Proof. Evaluate the definitions on the triangle; apply Theorem 6.3(ii). $\square$

6.1 The total reserve: Mack's Corollary

Definition 6.5 (Total-reserve estimators). Mack's Corollary: $\widehat{\text{mse}}(\hat{R}) = \sum_i \widehat{\text{mse}}_i + \sum_i \hat{C}_i (\sum_{j>i} \hat{C}_j) 2 \sum_{k \in \text{row } i} a_k$, the cross terms running over the older year's remaining factors. The conditional-resampling aggregate replaces $2 \sum_k a_k$ by $2(\prod_k (1 + a_k) - 1)$ (Buchwalder et al. 2006, Section 4.3).

Theorem 6.6 (Aggregated catalogue row). *The aggregated estimation-error terms differ by $\sum_i (\hat{C}_i^2 + 2\hat{C}_i \sum_{j>i} \hat{C}_j) (\prod_{k \in \text{row } i} (1 + a_k) - 1 - \sum_{k \in \text{row } i} a_k)$, which is nonnegative when the ultimates and the a_k are: Mack's total is at most the resampling total, by the same row-wise second-order remainder as in the single-year case.*

Proof. Termwise algebra and Lemma 6.2. □

Chapter 7

Solvency II: the one-year view

7.1 The one-year claims development result

Solvency II prices the uncertainty of the next twelve months, not of the run-off to ultimate. Merz and Wüthrich (2008) call the twelve-month move of the predicted ultimate the claims development result and separate the true one from the observable one. The filtration here is the one carried by **RandomTriangle**: $\mathcal{D}_k$ is everything observed up to development year k , for every accident year, the filtration in which (M1) is assumed in this development. Merz and Wüthrich index information by calendar year instead; the martingale property below is the same tower property in either filtration, and the closed form is stated in the filtration (M1) is assumed in.

Definition 7.1 (True one-year claims development result). For accident year i and development year k ,

$$\text{CDR}_i(k) = E[C_{i,n-1} \mid \mathcal{D}_k] - E[C_{i,n-1} \mid \mathcal{D}_{k+1}],$$

the change in the predicted ultimate claim when the information grows by one year.

Theorem 7.2 (Merz-Wüthrich 2008: the true CDR has conditional mean zero). *On a finite measure space, $E[\text{CDR}_i(k) \mid \mathcal{D}_k] = 0$ almost surely, for every accident year i and every k : the reserve set at time k is unbiased for the reserve that will be set at time $k+1$ together with the payments made in between.*

Proof. Linearity of the conditional expectation, then the tower property twice: the inner $\mathcal{D}_k$ -conditional expectation is idempotent, and $E[E[C_{i,n-1} \mid \mathcal{D}_{k+1}] \mid \mathcal{D}_k] = E[C_{i,n-1} \mid \mathcal{D}_k]$ because $\mathcal{D}_k \subseteq \mathcal{D}_{k+1}$. The two terms cancel. The only hypotheses are that $\mathcal{D}$ is a filtration inside the ambient σ -algebra and that the measure is finite; no integrability assumption on $C_{i,n-1}$ is needed for the identity as stated, since both conditional expectations are 0 off the integrable case. $\square$

Lemma 7.3 (Iterating (M1) from an arbitrary development year). *Under (M1), $E[C_{i,d+m} \mid \mathcal{D}_d] = C_{i,d} \prod_{j=d}^{d+m-1} f_j$ for every starting year d , hence $E[C_{i,n-1} \mid \mathcal{D}_k] = C_{i,k} \prod_{j=k}^{n-2} f_j$ for $k \leq n-1$.*

Proof. The induction of Lemma 5.11 with the starting year free rather than pinned to the latest observed year $n-1-i$ of the row; the base case is that $C_{i,d}$ is $\mathcal{D}_d$ -measurable. $\square$

Theorem 7.4 (Merz-Wüthrich 2008: the true CDR as a one-step residual). *Under (M1), for $k < n - 1$ and integrable claims,*

$$\text{CDR}_i(k) = \left(\prod_{j=k+1}^{n-2} f_j \right) (f_k C_{i,k} - C_{i,k+1})$$

almost surely: everything the next year of experience does to the predicted ultimate is carried by the single deviation of the new observation from its (M1) prediction.

Proof. Both predicted ultimates are given by Lemma 7.3; splitting $\prod_{j=k}^{n-2} f_j = f_k \prod_{j=k+1}^{n-2} f_j$ at the bottom index leaves the common factor $\prod_{j>k} f_j$ times $f_k C_{i,k} - C_{i,k+1}$. $\square$

Definition 7.5 (Observable one-year claims development result). The difference of the two chain-ladder predictions an actuary computes: the ultimate estimated from the triangle with n diagonals minus the ultimate estimated a year later, with every development factor re-estimated, from the triangle with $n + 1$ diagonals, $\widehat{\text{CDR}}_i = \hat{C}_{i,n-1}^{(n)} - \hat{C}_{i,n-1}^{(n+1)}$, together with its value along a random triangle outcome by outcome.

Lemma 7.6 (The displayed form). $\widehat{\text{CDR}}_i = \hat{R}_i^{(n)} - ((C_{i,n-i} - C_{i,n-1-i}) + \hat{R}_i^{(n+1)})$: *opening reserve minus the payments of the year and the closing reserve, the form Merz and Wüthrich display. Algebra on the definitions.*

Proof. Expand both reserves; the two claim entries cancel. $\square$

Nothing is proved here about the distribution of the observable CDR. The conditional mean squared error of prediction of $\widehat{\text{CDR}}_i$, which is what a Solvency II one-year reserve risk figure reports, rests in Merz and Wüthrich on an approximation of the same kind as Mack's: the estimated development factors are treated as resampled ones and terms of second order in the residuals are dropped. That step is not formalized, so the true-CDR statements above are exact and the observable-CDR statements of the paper are not yet available here.

Chapter 8

Röhr 2016: the error-propagation form

Röhr, *Chain ladder and error propagation*, ASTIN Bulletin 46 (2016) 293–330, derives the chain-ladder prediction error from the error-propagation formula. The chain-ladder ultimate $\hat{C}_{i,n-1} = C_{i,d} \prod_{k=d}^{n-2} \hat{f}_k$, $d = n - 1 - i$, is a product, so a first-order expansion in the development factors turns the relative squared error of the product into the *sum* of the relative squared errors of the factors. The paper’s classical-case display is

$$\frac{\widehat{\text{mse}}(\hat{R}_i)}{\hat{C}_{i,n-1}^2} = \sum_k \hat{u}_k^2,$$

where $\hat{u}_k^2$ is the relative uncertainty attached to development step k , and the derivation splits it into a process part and a parameter part,

$$\hat{u}_k^2 = \frac{\hat{\sigma}_k^2}{\hat{f}_k^2 \hat{C}_{i,k}} + \frac{\hat{\sigma}_k^2}{\hat{f}_k^2 S_k},$$

the first from the conditional variance $\sigma_k^2 C_{i,k}$ of the step itself, the second from the estimation variance σ_k^2/S_k of $\hat{f}_k$.

Source note. The full text of Röhr (2016) is paywalled and was not available for this formalization. What is formalized is the classical-case (ultimate run-off, single accident year) display quoted above, taken from the paper’s own abstract — “in the classical case treated by Mack (1993) the mean squared prediction error divided by the squared estimated ultimate loss can be written as $\sum_j \hat{u}_j^2$, where $\hat{u}_j$ measures the (relative) uncertainty around the j -th development factor and the proportion of the estimated ultimate loss that it affects”, with “a split into process error and parameter error” — together with the two parts in the form above. The paper’s claims-development-result formulas between two arbitrary future horizons are not formalized here, and neither is its own aggregation over accident years.

8.1 Definitions

Definition 8.1 (Röhr’s relative uncertainties). $\hat{u}_{k,\text{proc}}^2 = \hat{\sigma}_k^2/(\hat{f}_k^2 \hat{C}_{i,k})$, $\hat{u}_{k,\text{par}}^2 = \hat{\sigma}_k^2/(\hat{f}_k^2 S_k)$, and $\hat{u}_k^2 = \hat{u}_{k,\text{proc}}^2 + \hat{u}_{k,\text{par}}^2$.

Definition 8.2 (Röhr's linearized prediction error). The process term $\hat{C}_{i,n-1}^2 \sum_{k=d}^{n-2} \hat{u}_{k,\text{proc}}^2$, the parameter term $\hat{C}_{i,n-1}^2 \sum_{k=d}^{n-2} \hat{u}_{k,\text{par}}^2$, and their sum $\text{rohrMsep}_i = \hat{C}_{i,n-1}^2 \sum_{k=d}^{n-2} \hat{u}_k^2$.

8.2 The split, and the two Mack plug-ins

Lemma 8.3 (Process plus parameter). rohrMsep_i is the process term plus the parameter term.

Proof. Termwise, by construction. $\square$

Lemma 8.4 (Parameter term is Mack's estimation term). $\hat{u}_{k,\text{par}}^2 = a_k$, so Röhr's parameter term is Mack's estimation-error term $\hat{C}_{i,n-1}^2 \sum_k a_k$ exactly.

Proof. $\hat{\sigma}_k^2 / (\hat{f}_k^2 S_k) = (\hat{\sigma}_k^2 / \hat{f}_k^2) / S_k$. $\square$

Lemma 8.5 (Splitting the projection). For $n-1-i \leq k \leq m$, $\hat{C}_{i,m} = \hat{C}_{i,k} \prod_{l=k}^{m-1} \hat{f}_l$.

Proof. Split the product of the definition at k . $\square$

Theorem 8.6 (Process term is Mack's process variance). Let $i \leq n-1$ and suppose $\hat{f}_k \neq 0$ and $\hat{C}_{i,k} \neq 0$ for every $k = d, \dots, n-2$. Then Röhr's process term equals procVar , Mack's process variance recursion $V_{m+1} = \hat{f}_{d+m}^2 V_m + \hat{\sigma}_{d+m}^2 \hat{C}_{i,d+m}$, $V_0 = 0$, run for i steps with $\hat{f}, \hat{\sigma}^2$ plugged in.

Proof. Write $\hat{C}_{i,n-1} = \hat{C}_{i,k} \hat{f}_k \prod_{l=k+1}^{n-2} \hat{f}_l$ by Lemma 8.5. Then $\hat{C}_{i,n-1}^2 \hat{\sigma}_k^2 / (\hat{f}_k^2 \hat{C}_{i,k}) = (\prod_{l=k+1}^{n-2} \hat{f}_l)^2 \hat{\sigma}_k^2 \hat{C}_{i,k}$, which is the k -th summand of the closed form of procVar ; reindex $k = d+j$, $j = 0, \dots, i-1$. $\square$

8.3 Catalogue row 3

Theorem 8.7 (Röhr = Mack 1993). $\text{rohrMsep}_i = \widehat{\text{mse}}(\hat{R}_i)$, with no side condition: the linearized error-propagation form and Mack's 1993 closed form are equal term by term. This catalogue row therefore has zero correction term, in contrast with Theorem 6.3. Dividing by $\hat{C}_{i,n-1}^2 \neq 0$ recovers Röhr's display $\widehat{\text{mse}}(\hat{R}_i) / \hat{C}_{i,n-1}^2 = \sum_k \hat{u}_k^2$.

Proof. $\hat{\sigma}_k^2 / (\hat{f}_k^2 \hat{C}_{i,k}) + \hat{\sigma}_k^2 / (\hat{f}_k^2 S_k) = (\hat{\sigma}_k^2 / \hat{f}_k^2) (1 / \hat{C}_{i,k} + 1 / S_k)$ termwise. $\square$

Corollary 8.8 (Mack's closed form is process plus estimation). Under the hypotheses of Theorem 8.6, $\widehat{\text{mse}}(\hat{R}_i) = \text{mackProcess}_i + \text{mackEstimation}_i$: reading Mack's formula through Röhr's split identifies its two halves with the two plug-in estimators of the stochastic layer.

Proof. Chain Theorem 8.7, Lemma 8.3, Theorem 8.6 and Lemma 8.4. $\square$

Theorem 8.9 (What the linearization costs). Röhr's parameter term is the first-order part of the conditional-resampling term: their difference is exactly $\hat{C}_{i,n-1}^2 (\prod_k (1+a_k) - 1 - \sum_k a_k)$, and with $a_k \geq 0$ along the row Röhr's term is the smaller one. Linearizing in the estimated factors is precisely the step that replaces the product by the sum.

Proof. Lemma 8.4 and Theorem 6.3. $\square$

8.4 The total reserve

Definition 8.10 (Aggregate). Röhr's single-year terms carried into Mack's Corollary, with Mack's cross terms: $\sum_i \text{rohrMsep}_i + \sum_i \text{mackCross}_i$. The cross terms are Mack's; the paper's own aggregation over accident years could not be checked against the source, so this is recorded as a definition rather than attributed.

Theorem 8.11. *That aggregate is Mack's total-reserve estimator $\widehat{\text{mse}}(\hat{R})$.*

Proof. Theorem 8.7 row by row. □

8.5 The total reserve: the exact conditional MSEP

Mack's Corollary estimates the mean squared error of prediction of the total reserve $\hat{R} = \sum_i \hat{R}_i$. The estimator is Definition 6.5; the object it estimates is the conditional MSEP of the sum, and this section proves what that object is.

Write $\hat{C}_i = \hat{C}_{i,n-1}$ for the chain-ladder ultimate of accident year i , $C_i = C_{i,n-1}$ for the true ultimate, $\mathcal{D}$ for the σ -algebra of the observed data (the one Theorem 5.23 conditions on) and $M_i = E[C_i \mid \mathcal{D}]$.

Definition 8.12 (Cross-term condition). For a finite set s of accident years and targets Y_i ,

$$E[(Y_i - E[Y_i \mid \mathcal{D}])(Y_j - E[Y_j \mid \mathcal{D}]) \mid \mathcal{D}] = 0 \quad \text{for } i \neq j \text{ in } s.$$

This is what independence across accident years supplies, in the same way that the $\mathcal{D}_k$ -conditioned form of (M1) (Definition 5.2) is what it supplies for a single row. Like (M1), (M3) and the “no extra information” hypotheses of Theorem 5.23, it is carried as a hypothesis; the passage from independence of the rows to it is not formalized.

Lemma 8.13 (Conditional variance as a centred second moment). *For $Y \in L^2$, $E[(E[Y \mid \mathcal{D}] - Y)^2 \mid \mathcal{D}] = E[Y^2 \mid \mathcal{D}] - E[Y \mid \mathcal{D}]^2$.*

Proof. Lemma 5.22 with the predictor taken to be $E[Y \mid \mathcal{D}]$ itself, so that the estimation-error term is zero. Square integrability supplies every product integrability side condition by Hölder. □

Lemma 8.14 (Conditional variances add). *If $E[Z_i Z_j \mid \mathcal{D}] = 0$ for $i \neq j$ in s and each $Z_i \in L^2$, then $E[(\sum_i Z_i)^2 \mid \mathcal{D}] = \sum_i E[Z_i^2 \mid \mathcal{D}]$.*

Proof. Expand the square into the double sum $\sum_{i,j} Z_i Z_j$, take conditional expectations term by term, and drop the off-diagonal terms. □

Theorem 8.15 (Conditional MSEP of a sum). *For $\mathcal{D}$ -measurable predictors P_i of square-integrable targets Y_i whose centred versions satisfy Definition 8.12,*

$$E\left[\left(\sum_i P_i - \sum_i Y_i\right)^2 \mid \mathcal{D}\right] = \sum_i (E[Y_i^2 \mid \mathcal{D}] - E[Y_i \mid \mathcal{D}]^2) + \left(\sum_i (P_i - E[Y_i \mid \mathcal{D}])\right)^2.$$

The conditional variances add; the estimation errors add before they are squared.

Proof. Lemma 5.22 applied to the totals, then Lemma 8.13 to rewrite the conditional variance of the total as a centred second moment, then Lemma 8.14 to split it, then Lemma 8.13 again on each accident year. □

Lemma 8.16 (Theorem 1 and the process variance at the ultimate). *Under (M1), (M3) and the hypotheses of Theorem 5.23 that $\mathcal{D}$ carries no information about row i 's future beyond $\mathcal{D}_{n-1-i}$,*

$$M_i = C_{i,n-1-i} \prod_{l=n-1-i}^{n-2} f_l, \quad E[C_i^2 \mid \mathcal{D}] - M_i^2 = \text{procVar}(C_{i,n-1-i}, f, \sigma^2, n-1-i).$$

Proof. Specialize Lemma 5.11 and Theorem 5.21 to $m = i$, where $(n-1-i) + i = n-1$, and transport them from $\mathcal{D}_{n-1-i}$ to $\mathcal{D}$. $\square$

Theorem 8.17 (Mack's Corollary, exact form). *Under (M1), (M3), square integrability of the claims and of the chain-ladder predictions, the "no extra information" hypotheses of Theorem 5.23 for each accident year, and Definition 8.12 for the true ultimates,*

$$E\left[\left(\sum_i \hat{C}_i - \sum_i C_i\right)^2 \mid \mathcal{D}\right] = \sum_i \text{procVar}_i + \left(\sum_i (\hat{C}_i - M_i)\right)^2.$$

The process variances add; the estimation errors add before being squared, and expanding that square is the exact origin of the cross terms of Definition 6.5.

Proof. Theorem 8.15 with $P_i = \hat{C}_i$ and $Y_i = C_i$, then Lemma 8.16 term by term. $\square$

Corollary 8.18 (The total MSEP dominates the process variance). $\sum_i \text{procVar}_i \leq E[(\sum_i \hat{C}_i - \sum_i C_i)^2 \mid \mathcal{D}]$ almost everywhere, whatever the chain-ladder predictions happen to be.

Proof. The estimation-error term of Theorem 8.17 is a square. $\square$

Theorem 8.19 (The plug-in has the same two-part shape). *On a row whose development factors $\hat{f}_k$ and projections $\hat{C}_{i,k}$ do not vanish, the $1/\hat{C}_{i,k}$ half of Mack's formula is exactly procVar with $\hat{f}, \hat{\sigma}^2$ substituted and the $1/S_k$ half is exactly the plug-in estimation error: $\widehat{\text{mse}}_i = \text{mackProcess}_i + \text{mackEstimation}_i$, hence $\widehat{\text{mse}}(\hat{R}) = \sum_i \text{mackProcess}_i + \text{mackTotalEstimation}$. This is the plug-in counterpart of Theorem 8.17: process variances added, and everything else, single-year terms and cross terms alike, estimating the squared summed estimation error.*

Proof. Reindex the row sum over $\text{Ico}(n-1-i, n-1)$ by $j < i$, split the product $\prod_{l=n-1-i}^{n-2} \hat{f}_l$ at the factor $\hat{f}_{n-1-i+j}$, and cancel; then sum over accident years. $\square$

Theorem 8.20 (The cross-term condition is not vacuous). *On the nondegenerate model of Theorem 9.1 the three true ultimates are genuinely random and their centred versions are conditionally uncorrelated given the trivial σ -algebra: Definition 8.12 holds for $s = \{0, 1, 2\}$ without holding for want of randomness.*

Proof. The unconditional mean of each ultimate is cf^2 ; the centred ultimates are proportional to the shocks ξ_i , which are orthogonal. $\square$

Chapter 9

Non-vacuity: a nondegenerate model

Theorem 9.1 (A nondegenerate Mack model exists). *There is a probability space, a random triangle with three accident years, and constants f_k, σ_k^2 with $\sigma_0^2 > 0$, satisfying (M1), (M3) and (M2'), in which the first development step is genuinely random. The model: eight equally likely outcomes, an independent ± 1 shock per accident year, $C_{i,0} = c$ and $C_{i,k} = cf^k(1 + s\xi_i)$ for $k \geq 1$, with $\mathcal{D}_0$ trivial and $\mathcal{D}_k$ everything for $k \geq 1$.*

Proof. Conditional expectations given the trivial σ -algebra are eight-term averages; the shocks sum to zero, have unit square, and are orthogonal. $\square$

Theorem 9.2 (Headline theorems instantiated). *On that model, $E[\hat{f}_0 \mid \mathcal{D}_0] = 2$ although $\hat{f}_0$ is random; $E[\hat{C}_{2,2} \mid \mathcal{D}_0] = E[C_{2,2} \mid \mathcal{D}_0]$; $E[(\hat{f}_0 - 2)^2 \mid \mathcal{D}_0] = \sigma_0^2/S_0$; and $E[\hat{\sigma}_0^2 \mid \mathcal{D}_0] = \sigma_0^2 = 4$.*

Proof. Direct instantiation of the general theorems; every hypothesis is discharged. $\square$

Chapter 10

Reference computation

Mack's 1993 example (the Taylor and Ashe 1983 triangle) is reproduced to every printed digit (nine reserves, the total 18,681 thousand, all standard-error percentages) by a dependency-free script implementing the definitions above in floating point. The Lean definitions are over $\mathbb{R}$ and noncomputable; the script is the reference computation, not the Lean code.

Bibliography

- [1] T. Mack, Distribution-free calculation of the standard error of chain ladder reserve estimates, ASTIN Bulletin 23 (1993) 213-225.
- [2] T. Mack, The standard error of chain ladder reserve estimates: recursive calculation and inclusion of a tail factor, ASTIN Bulletin 29 (1999) 361-366.