

Tutorial 3: Dimensionality Reduction

Rui Zhao

rzhao@ee.cuhk.edu.hk

Feb. 6, 2014

Outline

- 1 Generalized Rayleigh Quotient
- 2 Lagrange Optimization
- 3 Exercises

Outline

- 1 Generalized Rayleigh Quotient
- 2 Lagrange Optimization
- 3 Exercises

1. Generalized Rayleigh Quotient

Fisher Criterion

$$J(\mathbf{w}) = \frac{\mathbf{w}^t \mathbf{S}_B \mathbf{w}}{\mathbf{w}^t \mathbf{S}_W \mathbf{w}} \quad (1)$$

$J(\mathbf{w})$ is the generalized Rayleigh quotient. A vector \mathbf{w} that maximizes $J(\cdot)$ must satisfy

$$\mathbf{S}_B \mathbf{w} = \lambda \mathbf{S}_W \mathbf{w} \quad (2)$$

for some constant λ .

1. Generalized Rayleigh Quotient

Maximizing $J(\mathbf{w})$ is equivalent to

$$\max_{\mathbf{w}} \quad \mathbf{w}^t \mathbf{S}_B \mathbf{w} \quad (3)$$

$$s.t. \quad \mathbf{w}^t \mathbf{S}_W \mathbf{w} = K \quad (4)$$

which can be solved using Lagrange multipliers.

1. Generalized Rayleigh Quotient