

Center of Mass of a Conical Shell

To find the center of mass of a conical shell with radius a and height h , we consider the shell to be thin and use the concept of continuous mass distribution. The center of mass (z_{CM}) of the conical shell is calculated by considering an infinitesimally thin circular slice of the cone at a height z from the base, with thickness dz and radius $r(z)$. The relationship between the radius of the slice and its height is linear, given by $r(z) = \frac{a}{h}z$.

The center of mass z_{CM} is given by:

$$z_{\text{CM}} = \frac{\int_0^h z \, dm}{\int_0^h dm}$$

The mass of an infinitesimal slice (dm) is proportional to its area. Considering the surface area of the infinitesimal slice for a thin shell, the mass distribution is uniform over its surface, hence $dm = \sigma 2\pi r(z) dz$, with σ being the surface mass density (mass per unit area) of the shell.

Substituting $r(z) = \frac{a}{h}z$ into the expression for dm gives:

$$dm = \sigma 2\pi \left(\frac{a}{h}z \right) dz$$

Substituting dm into the integral for z_{CM} , we get:

$$z_{\text{CM}} = \frac{\int_0^h z \sigma 2\pi \left(\frac{a}{h}z \right) dz}{\int_0^h \sigma 2\pi \left(\frac{a}{h}z \right) dz}$$

Simplifying the integrals:

$$z_{\text{CM}} = \frac{\int_0^h z^2 dz}{\int_0^h z dz} = \frac{\left[\frac{1}{3}z^3 \right]_0^h}{\left[\frac{1}{2}z^2 \right]_0^h} = \frac{\frac{1}{3}h^3}{\frac{1}{2}h^2} = \frac{2}{3}h$$

Therefore, the center of mass of the conical shell is located at a distance of $\frac{h}{3}$ from the base along the axis of symmetry, correcting the initial mistake in the explanation.