

Robert Davidson
BSc Mathematics and Computer Science

Linear Algebra (MA283) notes

Date: April 10, 2025

Abstract

Linear algebra provides a framework to understand and analyze systems defined by linear equations and transformations, serving as the foundation of diverse areas in mathematics, science, engineering, and beyond. These notes introduce and build upon core concepts, beginning with **systems of linear equations** and extending into abstract vector spaces, transformations, and geometric interpretations.

We first explore the fundamental notions of **linear independence**, **subspaces**, and **span**, which clarify the structure underlying linear systems. Equipped with these tools, we define **basis** and **dimension**, revealing how vector spaces can be succinctly and precisely described. This naturally leads us to **coordinates**, enabling clear communication about vectors across different bases, and the crucial concepts of **rank** and **nullity**, capturing essential properties of linear transformations.

Next, we delve into **inner products**, **orthogonality**, and **projections**, extending geometric intuition into abstract vector spaces. These ideas give rise to practical methods like the Gram-Schmidt process, orthogonal projections, and least squares approximations, which are indispensable in both theoretical and real-world contexts.

Finally, we examine **eigenvalues**, **eigenvectors**, and **diagonalizability**, exploring how linear transformations can be simplified and understood by choosing optimal bases. These powerful tools provide insight into the intrinsic properties of matrices and transformations, enabling deeper analysis and efficient computation.

Together, these notes offer a coherent journey through linear algebra, balancing theoretical depth with practical applications, equipping readers with both the conceptual clarity and computational skills needed to leverage linear algebra across various fields.

Contents

1	Review of Matrix Algebra	3
2	Systems of linear equations	5
2.1	Linear equations and Solution Sets	5
2.1.1	Interpreting Linear Systems as Matrix Equations	6
2.1.2	Cases of Solutions	6
2.2	Elementary Row Operations	6
2.2.1	REF and Strategy	7
2.2.2	Row Reduced Echelon Form	7
2.3	Leading variables and free variables	7
2.4	Consistent and Inconsistent Systems	8
2.5	Possible Outcomes when solving a system of equations	8
2.6	Elementary Row Operations as Matrix Transformations	8
2.6.1	Multiplying a Row by a Non-Zero Scalar	8
2.6.2	Switching Two Rows	9
2.6.3	Adding a Multiple of One Row to Another	9
2.7	EROs and Inverses	10
2.8	Determinant	10
2.9	Key Takeaways	11
3	Subspaces, Span, and Linear Independence	12
3.1	Subspaces	12
3.2	The Image and Kernel of a Linear Transformation	12
3.2.1	Image of T	13
3.2.2	Kernel of T	13
3.2.3	Column Dependence	13
3.3	The span : how to make subspaces	14
3.4	Spanning sets	14
3.4.1	Questions about spanning sets	14
3.5	Linear Dependence and Linear Independence	15
3.6	Test for linear independence	15
3.7	Key Takeaways	16
4	Basis, Dimension, and Rank	16
4.1	Finite Dimensional Spaces	16
4.2	Basis: The Structural Backbone	17
4.3	Steinitz Replacement Lemma	17
4.4	Consequences of the replacement theorem	18
4.5	Every basis has the same number of elements	18
4.6	Row rank and column rank	18
4.7	Row rank = column rank	19
4.8	Key Takeaways	20
5	Coordinates, Rank-Nullity, and Change of Basis	20
5.1	Coordinates	20
5.2	Coordinates with respect to different bases	21
5.3	The Rank Nullity Theorem	21
5.4	Linear transformations and change of basis	22
5.5	More on Change of Basis	22
5.6	Key Takeaways	23
6	Similarity, Eigenvalues, and Diagonalizability	23
6.1	Similarity (The relation of A and A')	24
6.2	Similar Matrices	25
6.3	Two interpretations of diagonalizability	25
6.4	Eigenvalues and diagonalizability	25
6.5	Two examples of non-diagonalizability	26
6.6	A Shear in \mathbb{R}^2	26
6.7	Eigenvector for distinct eigenvalues are independent	26
6.8	At most n distinct eigenvalues	27
6.9	Notes about determinants and characteristic polynomials	27
6.10	Multiplicity of Eigenvalues	27
6.11	Multiplicity of Eigenvalues	27
6.12	Geometric multiplicity \leq Algebraic multiplicity	28
6.13	Key Takeaways	28
7	Inner Products, Orthogonality, and Projections	29
7.1	Inner Products	29
7.2	Real Inner Products	29
7.2.1	Examples of Inner Products	30
7.3	Length, Distance and Scaling in an Inner Product Space	30
7.4	Orthogonality in an inner product space	30
7.5	Orthogonal Projection	31
7.6	Orthogonal projection of one vector on another	31
7.7	Orthogonal Bases (the Gram-Schmidt process)	31
7.8	Orthogonal projection on a subspace	32
7.9	Calculating the projection on a subspace	32
7.10	$\text{proj}_W(v)$ is the nearest point of W to v	32
7.11	Recall	32
7.12	Orthogonal Basis	33
7.13	Projection on a subspace	33
7.14	How to calculate $\text{proj}_W(v)$	33
7.15	Application: least squares for overdetermined systems	34

1 Review of Matrix Algebra

Matrix Addition

If a matrix has m rows and n columns, we say it is $m \times n$. **Two matrices can only be added if they have the same size..** In this case, we just add the entries in each position.

The $m \times n$ **zero** matrix is a matrix with all entries equal to 0. It is the **Identity element** for matrix addition (adding it to any matrix does not change the matrix)

Matrix Multiplication by a Scalar

This simply means multiplying each entry of the matrix by the scalar. For example:

$$\alpha \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} \alpha & 2\alpha \\ 3\alpha & 4\alpha \end{bmatrix}$$

Vector Space

Definition 1.1

The most typical example of a **vector space** is \mathbb{R}^n , the all n sized vector of real numbers

$$\{(x_1, \dots, x_n) \mid x_i \in \mathbb{R}\}$$

What makes this a vector space is that we can add two n -vector together:

$$(x, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n)$$

and we can multiply each vector by a scalar, α :

$$\alpha(x_1, \dots, x_n) = (\alpha x_1, \dots, \alpha x_n)$$

Linear Combinations

Suppose v_1, v_2, \dots, v_k are elements that can be added together and multiplied by scalars.

A Linear Combination of v_1, v_2, \dots, v_k is an expression of the form:

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_k v_k$$

where $a_i \in \mathbb{R}$ are scalars, called **coefficients**.

Matrix-Vector Multiplication

Let A be a $m \times n$ matrix, and \mathbf{v} be a column vector with n entries ($n \times 1$ matrix).

Then the matrix vector product Av is the column vector, with m entries, obtained by taking the linear combination of the columns of A with the entries of \mathbf{v} as coefficients.

Example:

$$\begin{bmatrix} -1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix} \begin{bmatrix} 7 \\ 6 \\ 9 \end{bmatrix} = 7 \begin{bmatrix} -1 \\ 0 \end{bmatrix} + 6 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + 9 \begin{bmatrix} 4 \\ 3 \end{bmatrix} = \begin{bmatrix} 41 \\ 33 \end{bmatrix}.$$

Remark: Av , if defined, has the same number of rows as A and the same number of columns as \mathbf{v} .

Matrix-Matrix Multiplication

Let A and B be matrices of size $m \times p$ and $p \times n$, respectively. Write v_1, \dots, v_n for the columns of B . Then the product AB is the $m \times n$ matrices whose columns are Av_1, \dots, Av_n .

The entry at row i and column j of the matrix A is given by A_{ij} . The entry in the i, j position of the product AB is the i th entry of the vector Av_j , where the vector v_j is the j th column of B . In other words, the entry in the i, j position of the product AB is given by:

$$(AB)_{ij} = A_{i1}B_{1j} + A_{i2}B_{2j} + \dots + A_{ip}B_{pj} = \sum_{k=1}^p A_{ik}B_{kj}$$

Alternatively, if A is $m \times p$ with rows u_1, \dots, u_m and B is $p \times n$ with columns v_1, \dots, v_n , then the product AB is:

$$AB = \begin{bmatrix} u_1 \cdot v_1 & u_1 \cdot v_2 & \dots & u_1 \cdot v_n \\ u_2 \cdot v_1 & u_2 \cdot v_2 & \dots & u_2 \cdot v_n \\ \vdots & \vdots & \ddots & \vdots \\ u_m \cdot v_1 & u_m \cdot v_2 & \dots & u_m \cdot v_n \end{bmatrix}$$

Example:

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \quad B = \begin{bmatrix} 7 & 8 \\ 9 & 10 \\ 11 & 12 \end{bmatrix} \quad AB = \begin{bmatrix} 1 \cdot 7 + 2 \cdot 9 + 3 \cdot 11 & 1 \cdot 8 + 2 \cdot 10 + 3 \cdot 12 \\ 4 \cdot 7 + 5 \cdot 9 + 6 \cdot 11 & 4 \cdot 8 + 5 \cdot 10 + 6 \cdot 12 \end{bmatrix} = \begin{bmatrix} 58 & 64 \\ 139 & 154 \end{bmatrix}$$

For matrices A and B , the products AB and BA are generally not equal, even if they are both defined and even if both have the same size.

Linear Transformations

Let m and n be positive integers.

A **linear transformation** T from \mathbb{R}^n to \mathbb{R}^m is a function $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ that satisfies:

- $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$
- $T(\lambda \mathbf{u}) = \lambda T(\mathbf{u})$ for all $\mathbf{u} \in \mathbb{R}^n$ and $\lambda \in \mathbb{R}$

Matrix of a Linear Transformation

Suppose $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ is the linear transformation:

$$T \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -6 \\ 7 \end{bmatrix}$$

Then for the vector in \mathbb{R}^3 with entries a, b, c :

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = aT \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + bT \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + cT \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 & -6 \\ -3 & 4 & 7 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

Where the 2×3 matrix M_T is called the **standard matrix** of A . A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ can be completely represented by an $m \times n$ matrix M_T .

Understanding the Matrix Representation

- The columns of matrix M_T are the images of the standard basis vectors e_1, e_2, \dots, e_n under T .
- For any vector $v \in \mathbb{R}^n$, we calculate $T(v)$ by multiplying: $M_T \cdot v$.
- Therefore, matrix-vector multiplication is simply evaluating a linear transformation.

Correspondence: Any $m \times n$ matrix A defines a linear transformation $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by: $T_A(v) = Av$. Linear transformations include rotations, reflections and scaling

Efficiency of Representation: A remarkable property of linear transformations is their information efficiency:

- To completely define $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, we need only mn values.
- These values are the coordinates of the n transformed basis vectors in \mathbb{R}^m .
- This differs fundamentally from general continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$, which cannot be fully determined by their values at finitely many points.

Matrix multiplication is composition

Suppose that $T : \mathbb{R}^n \rightarrow \mathbb{R}^p$ and $S : \mathbb{R}^p \rightarrow \mathbb{R}^m$ are linear transformations. Then the composition $S \circ T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is also a linear transformation from \mathbb{R}^n to \mathbb{R}^m defined for $\mathbf{v} \in \mathbb{R}^n$ by:

$$S \circ T(\mathbf{v}) = S(T(\mathbf{v}))$$

To see how that the $m \times n$ matrix $M_{S \circ T}$ depends on the matrix $M_S(m \times p)$ and $M_T(p \times n)$ we look at the definition of $M_{S \circ T}$:

- The first column has coordinates $S \circ T(e_1) = S(T(e_1))$
- $T(e_1)$ is first column of M_T
- Then $S(T(e_1))$ is the matrix-vector product $M_S \cdot M_T(e_1)$
- Same for all other columns $\implies M_{S \circ T} = M_S \cdot M_T$

Thus, we conclude **matrix multiplication is composition of linear transformations**.

Key Takeaways

Key Takeaways 1.1: Review of Matrix Algebra

- **Vector Space:** A set of objects (vectors) that can be added and multiplied by scalars.
- **Matrix Addition:** Only defined for matrices of the same size. Entrywise: $(A + B)_{ij} = A_{ij} + B_{ij}$. The zero matrix is the additive identity.
- **Scalar Multiplication:** Multiply each entry by the scalar α : $\alpha \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} \alpha & 2\alpha \\ 3\alpha & 4\alpha \end{bmatrix}$
- **Linear Combinations:** $\alpha_1 v_1 + \dots + \alpha_k v_k$ with $\alpha_i \in \mathbb{R}$ defines a linear combination.
- **Matrix-Vector Multiplication:** For $A \in \mathbb{R}^{m \times n}$ and $v \in \mathbb{R}^n$, the product $Av \in \mathbb{R}^m$ is:

$$Av = v_1 A_1 + v_2 A_2 + \dots + v_n A_n$$

- **Matrix-Matrix Multiplication:** If $A \in \mathbb{R}^{m \times p}$, $B \in \mathbb{R}^{p \times n}$, then:

$$(AB)_{ij} = \sum_{k=1}^p A_{ik} B_{kj}$$

- **Not commutative** The matrix product $AB \neq BA$ in general.
- **Linear Transformations:** $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear if:

$$T(u + v) = T(u) + T(v), \quad T(\lambda u) = \lambda T(u)$$

- **Standard Matrix of A:** A matrix M_T such that $T(v) = M_T v$. The columns of M_T are the images of the standard basis vectors.
- **Composition = Multiplication:** For $S : \mathbb{R}^p \rightarrow \mathbb{R}^m$, $T : \mathbb{R}^n \rightarrow \mathbb{R}^p$,

$$(S \circ T)(v) = S(T(v)) \quad \Rightarrow \quad M_{S \circ T} = M_S M_T$$

2 Systems of linear equations

This section introduces the fundamental notion of a **linear equation** in several variables, examines how to combine multiple linear equations into a **system**, and discusses systematic methods for finding all solutions. We begin by defining a linear equation and its **solution set**, and then show how to represent an entire system of equations in a concise **matrix form**. From there, we take a closer look at how to solve such systems using **elementary row operations** (EROs), the building blocks of a powerful technique known as Gaussian elimination.

A central theme here is how EROs transform a matrix into forms (REF and RREF) that make solutions transparent. This leads to an understanding of **leading variables** versus **free variables**, which determines whether a system has a unique solution, infinitely many solutions, or no solution at all (i.e., whether the system is **consistent** or **inconsistent**). We then see that the same row operations can be captured by multiplying on the left by specially designed **elementary matrices**, each of which is invertible. This viewpoint not only deepens our understanding of how systems are solved but also shows that any invertible matrix can be decomposed into a product of elementary matrices.

As a culminating example, we use these methods to **find the inverse of a matrix** by performing row operations on an augmented matrix $[A \mid I]$ until A becomes the identity. All these ideas form the core machinery for handling systems of linear equations throughout linear algebra.

2.1 Linear equations and Solution Sets

A linear equation in the variables x and y is an equation of the form

$$2x + y = 3$$

If we replace x and y with some numbers, the statement **becomes true or false**.

A pair, $(x_0, y_0) \in \mathbb{R}$, is a solution to a linear equation if setting $x = x_0$ and $y = y_0$ **makes the equation true**.

Definition 2.1: Solution set

The **solution set** is the set of all solutions to a linear equation.

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n = b \quad \text{where } a_i, b \in \mathbb{R}$$

is an **affine hyperplane** in \mathbb{R}^n ; geometrically resembles a copy of \mathbb{R}^{n-1} inside \mathbb{R}^n .

2.1.1 Interpreting Linear Systems as Matrix Equations

For some equations:

$$\begin{aligned} a_{11}x_1 + \cdots + a_{1n}x_n &= b_1 \\ &\vdots \\ a_{m1}x_1 + \cdots + a_{mn}x_n &= b_m \end{aligned}$$

We can write:

$$\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}, \quad A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

and our unknowns as:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

Then, we can rewrite our system of linear equations in form form:

$$A\mathbf{x} = \mathbf{b}$$

For Example

$$\begin{aligned} x + 2y - z &= 5 \\ 3x + y - 2z &= 9 \\ -x + 4y + 2z &= 0 \end{aligned} \Rightarrow \begin{bmatrix} 1 & 2 & -1 \\ 3 & 1 & -2 \\ -1 & 4 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 9 \\ 0 \end{bmatrix}$$

2.1.2 Cases of Solutions

- **$m > n$ no solutions** (more equations than unknowns)
For example, let $m = 3, n = 2$, that is we have 3 equations and 2 unknowns, this corresponds to the fact that three lines in a plane will usually have no common point of intersection.
- **$m < n$ infinitely many solutions** (more unknowns than equations)
For example, $m = 2, n = 3$, that is 2 equations, with 3 unknowns, this corresponds to the fact that two planes in space will usually intersect in an entire line.
- **$m = n$:** Much of linear algebra deals with this remaining case.




Figure 1: Geometric interpretation of linear systems

2.2 Elementary Row Operations

Suppose that the square matrix has an inverse A^{-1} , which means $A^{-1}A = I$. With I being the identity matrix. Then our solution will be:

$$\mathbf{x} = A^{-1}\mathbf{b}$$

Since:

$$A\mathbf{x} = A(A^{-1}\mathbf{b}) = I\mathbf{b} = \mathbf{b}$$

To solve a system of linear equations we associate an **augmented matrix** to the system of equations. For example:

$$\begin{array}{rrcr} x & + & 2y & - & z & = & 5 \\ 3x & + & y & - & 2z & = & 9 \\ -x & + & 4y & + & 2z & = & 0 \end{array} \Rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 5 \\ 3 & 1 & -2 & 9 \\ -1 & 4 & 2 & 0 \end{array} \right]$$

To solve, we can perform the following **Elementary Row Operations (EROs)**:

1. Multiply a row by a non-zero constant.
2. Add a multiple of one row to another row.
3. Swap two rows.

The goal of these operations is to transform the augmented matrix into **row echelon form (REF)** or **reduced row echelon form (RREF)**.

2.2.1 REF and Strategy

We say a matrix is in **row echelon form (REF)** if:

- The first non zero entry in each row is a 1 (called the **leading 1**).
- If a column has a leading 1, then all entries below it are 0.
- The leading 1 in each row is to the right of the leading 1 in the previous row.
- All rows of 0s are at the bottom of the matrix.

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 1 & -1 \end{array} \right]$$

Example of REF

We have produced a new system of equations. This is easily solved by back substitution.

Concept 2.1: Strategy for Obtaining REF

- Get a 1 as the top left entry
- Use this 1 to clear the entries below it
- Move to the next column and repeat
- Continue until all leading 1s are in place
- Use back substitution to solve the system

2.2.2 Row Reduced Echelon Form

A matrix is in **reduced row echelon form (RREF)** if:

- It is in REF
- The leading 1 in each row is the only non-zero entry in its column.

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \end{array} \right]$$

Example of RREF

2.3 Leading variables and free variables

We'll start by an example:

$$\begin{array}{rrcr} x_1 & - & x_2 & - & x_3 & + & 2x_4 & = & 0 \\ 2x_1 & + & x_2 & - & x_3 & + & 2x_4 & = & 8 \\ x_1 & - & 3x_2 & + & 2x_3 & + & 7x_4 & = & 2 \end{array} \Rightarrow \left[\begin{array}{cccc|c} 1 & -1 & -1 & 2 & 0 \\ 2 & 1 & -1 & 2 & 8 \\ 1 & -3 & 2 & 7 & 2 \end{array} \right]$$

Solving this system of equations, we get:

$$\text{RREF: } \left[\begin{array}{cccc|c} 1 & 0 & 0 & 2 & 4 \\ 0 & 1 & 0 & -1 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{array} \right] \Rightarrow \begin{array}{l} x_1 + 2x_4 = 4 \\ x_2 - x_4 = 2 \\ x_3 + x_4 = 2 \end{array} \Rightarrow \begin{array}{l} x_1 = 4 - 2x_4 \\ x_2 = 2 + x_4 \\ x_3 = 2 - x_4 \end{array}$$

This RREF tells us how the **leading variables** (x_1, x_2, x_3) depend on the **free variable** (x_4). The free variable can take any value in \mathbb{R} . We write the solution set as:

$$x_1 = 4 - 2t, \quad x_2 = 2 + t, \quad x_3 = 2 - t, \quad x_4 = t \quad \text{where } t \in \mathbb{R}$$

$$(x_1, x_2, x_3, x_4) = (4 - 2t, 2 + t, 2 - t, t); \quad t \in \mathbb{R}$$

Definition 2.2: Leading and Free Variables

- **Leading variable** : A variable whose columns in the RREF contain a leading 1
- **Free variable** : A variable whose columns in the RREF do not contain a leading 1

2.4 Consistent and Inconsistent Systems

Consider the following system of equations:

$$\begin{array}{rrcr} 3x & + & 2y & - & 5z & = & 4 \\ x & + & y & - & 2z & = & 1 \\ 5x & + & 3y & - & 8z & = & 6 \end{array} \Rightarrow \left[\begin{array}{ccc|c} 3 & 2 & -5 & 4 \\ 1 & 1 & -2 & 1 \\ 5 & 3 & -8 & 6 \end{array} \right] \Rightarrow \left[\begin{array}{ccc|c} 1 & 1 & -2 & 1 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 1 \end{array} \right] \quad (\text{REF})$$

We can see the last row of the REF is:

$$0x + 0y + 0z = 1$$

This equation clearly has no solution, and hence the system has no solutions. We say the system is **inconsistent**. Alternatively, we say the system is **consistent** if it has at least one solution.

2.5 Possible Outcomes when solving a system of equations

- The system may be **inconsistent** (no solutions) - i.e:

$$[0 \ 0 \ \dots \ 0 \ | \ a] \quad a \neq 0$$

- The system may be **consistent** which occurs if:

- **Unique Solutions** each column (aside from the rightmost) contains a single leading 1. - i.e:

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 4 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

- **Infinitely many solutions** at least one variable does not appear as a leading 1 in any row, making it a free variable - i.e:

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

2.6 Elementary Row Operations as Matrix Transformations

Elementary row operations may be interpreted as **matrix multiplication**. To see this, first we introduce the **Identity matrix**:

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

The I_m Identity matrix is an $m \times m$ matrix with 1s on the diagonal and 0s elsewhere. We also introduce the $E_{i,j}$ matrix which has 1 in the (i, j) position and 0s elsewhere. For example:

$$E_{1,2} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Then:

$$I_3 + 4E_{1,2} = \begin{bmatrix} 1 & 4 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Performing a row operation on A is the same as multiplying A by an appropriate matrix E on the left. These matrices are called **elementary matrices**. They are **always invertible**, and **their inverses are also elementary matrices**. The statement:

"every matrix can be reduced to RREF through EROs"

is equivalent to saying that

"for every matrix A with m rows, there exists a $m \times m$ matrix B which is a product of elementary matrices such that BA is in RREF."

2.6.1 Multiplying a Row by a Non-Zero Scalar

When multiplying row i of matrix A by a scalar $\alpha \neq 0$, we can use the matrix:

$$I_m + (\alpha - 1)E_{i,i}$$

This works because it modifies only the (i, i) entry of the identity matrix to be α while keeping all other entries unchanged. When multiplied with A , it scales row i by α and leaves all other rows intact.

Example: If $\alpha = 5$ and $i = 2$, then:

$$I_3 + 4E_{2,2} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \quad (I_3 + 4E_{2,2})A = \begin{bmatrix} 1 & 2 & 3 \\ 20 & 25 & 30 \\ 7 & 8 & 9 \end{bmatrix}$$

2.6.2 Switching Two Rows

To swap rows i and k , we use:

$$S = I_m + E_{i,k} + E_{k,i} - E_{i,i} - E_{k,k}$$

This works by:

- Removing the 1's at positions (i, i) and (k, k) from the identity matrix
- Adding 1's at positions (i, k) and (k, i)

Example: Swapping rows 1 and 3:

$$S = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \quad SA = \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

2.6.3 Adding a Multiple of One Row to Another

To replace row k with row $k + \alpha \times$ row i , use:

$$I_m + \alpha E_{k,i}$$

This adds α times row i to row k while leaving all other rows unchanged because:

- For any row $j \neq k$, the corresponding row in this matrix is just the standard basis row
- Row k becomes the sum of the standard basis row k plus α times the standard basis row i

Example: Adding 3 times row 1 to row 2:

$$I_3 + 3E_{2,1} = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \quad (I_3 + 3E_{2,1})A = \begin{bmatrix} 1 & 2 & 3 \\ 7 & 11 & 15 \\ 7 & 8 & 9 \end{bmatrix}$$

Example 2.1

Write the inverse of an elementary matrix and show it is an elementary matrix.

Multiplying a row by a nonzero scalar:

- **Operation:** Multiply row i by $\alpha \neq 0$.
- **Elementary Matrix:** $E = I_m + (\alpha - 1)E_{i,i}$
- **Inverse:** To reverse the operation, multiply row i by $1/\alpha$. Hence, the inverse is

$$E^{-1} = I_m + \left(\frac{1}{\alpha} - 1\right) E_{i,i} = I_m + \frac{1 - \alpha}{\alpha} E_{i,i}.$$

Swapping two rows:

- **Operation:** Swap rows i and k .
- **Elementary Matrix:** $S = I_m - E_{i,i} - E_{k,k} + E_{i,k} + E_{k,i}$
- **Inverse:** Since swapping the same two rows twice returns them to their original positions,

$$S^{-1} = S.$$

Adding a multiple of one row to another:

- **Operation:** Add α times row i to row k .
- **Elementary Matrix:** $E = I_m + \alpha E_{k,i}$
- **Inverse:** To undo the operation, subtract α times row i from row k . Therefore,

$$E^{-1} = I_m - \alpha E_{k,i}.$$

Example 2.2

Prove that every invertible matrix in $M_n(\mathbb{R})$ is a product of elementary matrices.

Let A be an invertible matrix in $M_n(\mathbb{R})$. Since A is invertible, we can use Gaussian elimination to transform A into the identity matrix I_n .

Let E_1, E_2, \dots, E_k be the elementary matrices corresponding to the row operations used in the elimination process. Then, we have:

Multiplying a row by a scalar: $I_n + (\alpha - 1)E_{i,i}$

Swapping two rows: $I_n + E_{i,k} + E_{k,i} - E_{i,i} - E_{k,k}$

Adding a multiple of one row to another: $I_n + \alpha E_{k,i}$

Applying these in sequence to A gives:

$$E_k \cdots E_2 E_1 A = I_n$$

Since $E_k \cdots E_2 E_1 = I_n$, we can multiply both sides by $(E_k \cdots E_2 E_1)^{-1}$ on the left to obtain:

$$A = (E_k \cdots E_2 E_1)^{-1} I_n = (E_k \cdots E_2 E_1)^{-1}$$

Using the property:

$$(E_k \cdots E_2 E_1)^{-1} = E_1^{-1} E_2^{-1} \cdots E_k^{-1},$$

we can express A as a product of elementary matrices:

$$A = E_1^{-1} E_2^{-1} \cdots E_k^{-1}$$

Since each E_i is an elementary matrix, its inverse is also an elementary matrix. Therefore, A can be expressed as a product of elementary matrices.

2.7 EROs and Inverses

Elementary Row Operations can be used to find the inverse of a square matrix. Consider a square matrix $A \in M_n(\mathbb{F})$ (that is, an $n \times n$ matrix over a field \mathbb{F}). If A is invertible, let

$$A^{-1} = \left[\begin{array}{c|c|c|c} | & | & & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & & | \end{array} \right]$$

be its inverse, where each \mathbf{v}_i is the i th column of A^{-1} . By definition of the matrix inverse, we have

$$A A^{-1} = A \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix} = \begin{bmatrix} A\mathbf{v}_1 & A\mathbf{v}_2 & \cdots & A\mathbf{v}_n \end{bmatrix} = I_n,$$

the $n \times n$ identity matrix. This implies that

$$A\mathbf{v}_i = \mathbf{e}_i, \quad \text{for each } i = 1, 2, \dots, n,$$

where \mathbf{e}_i is the i th column of I_n (which has a 1 in the i th row and 0 everywhere else). In other words, each column \mathbf{v}_i of A^{-1} is the unique solution to the linear system

$$A\mathbf{v}_i = \mathbf{e}_i.$$

To find A^{-1} effectively, we form the augmented matrix $[A \mid I_n]$ and apply EROs to transform A into I_n . When this is achieved, the augmented portion becomes A^{-1} . Thus, we have

$$\text{RREF}([A \mid I_n]) = [I_n \mid A^{-1}].$$

2.8 Determinant

Before the definition, we start with notation. For an $n \times n$ matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}$$

, we define A_{ij} as the $(n-1) \times (n-1)$ matrix obtained by deleting the i th row and j th column of A . For example:

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \quad \text{then } A_{12} = \begin{bmatrix} 4 & 6 \\ 7 & 9 \end{bmatrix}$$

Definition 2.3

$$\det(A) = \sum_{k=1}^n (-1)^{k+1} a_{1k} \det(A_{1k})$$

where A_{1k} is the $(n-1) \times (n-1)$ matrix obtained by deleting the first row and k th column of A .

Example 2.3

Find A^{-1} if $A = \begin{bmatrix} 3 & 4 & -1 \\ 1 & 0 & 3 \\ 2 & 5 & -4 \end{bmatrix}$.

We form a 3×6 matrix $A' = [A \mid I_3]$:

$$A' = \left[\begin{array}{ccc|ccc} 3 & 4 & -1 & 1 & 0 & 0 \\ 1 & 0 & 3 & 0 & 1 & 0 \\ 2 & 5 & -4 & 0 & 0 & 1 \end{array} \right]$$

We apply the following EROs to A' :

- $R_1 \leftrightarrow R_2$
- $R_2 \rightarrow R_2 - 3R_1$
- $R_3 \rightarrow R_3 - 2R_1$
- $R_3 \rightarrow R_3 + R - 2$
- $R_3 \leftrightarrow R_2$
- $R_3 \rightarrow R_3 - 4R_2$
- $R_3 \times (-\frac{1}{10})$
- $R_1 \rightarrow R_1 - 3R_3$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{3}{2} & -\frac{11}{10} & -\frac{6}{5} \\ 0 & 1 & 0 & -1 & 1 & 1 \\ 0 & 0 & 1 & -\frac{1}{2} & \frac{7}{10} & \frac{2}{5} \end{array} \right]$$

That is:

$$A^{-1} = \begin{bmatrix} \frac{3}{2} & -\frac{11}{10} & -\frac{6}{5} \\ -1 & 1 & 1 \\ -\frac{1}{2} & \frac{7}{10} & \frac{2}{5} \end{bmatrix}$$

It is easily checked that $AA^{-1} = I_3$.

2.9 Key Takeaways**Key Takeaways 2.1**

- A **solution set** is the set of all equations that satisfy $a_1X_1 + a_2X_2 + \dots + a_nX_n = b$, where $a_i, b \in \mathbb{R}$.
- The **solution set** is a copy of \mathbb{R}^{n-1} inside \mathbb{R}^n
- **REF** is a matrix form where the first non-zero entry in each row is a 1, and all entries below it are 0.
- **RREF** is a matrix form where the leading 1 in each row is the only non-zero entry in its column.
- **Leading variables** are variables whose columns in the RREF contain a leading 1, while **free variables** do not.
- **Inconsistent systems** have no solutions, while **consistent systems** have at least one solution.
- **Identity Matrix** (I_m) is an $m \times m$ matrix with 1s on the diagonal and 0s elsewhere.
- $E_{i,j}$ matrix has 1 in the (i,j) position and 0s elsewhere.
- **ERO's as matrix multiplication:**
 - Multiply a row by a non-zero scalar: $I_m + (\alpha - 1)E_{i,i}$
 - Switch two rows: $S = I_m + E_{i,k} + E_{k,i} - E_{i,i} - E_{k,k}$
 - Add a multiple of one row to another: $I_m + \alpha E_{k,i}$
- **ERO to find inverse:** To find the inverse of a square matrix A , form the augmented matrix $[A \mid I_n]$ and apply EROs to transform A into I_n . The augmented portion becomes A^{-1} . that is

$$RREF([A \mid I_n]) = [I_n \mid A^{-1}]$$

3 Subspaces, Span, and Linear Independence

Introduction

In this section, we explore a series of interconnected ideas that shape our understanding of how sets of vectors can form specialized structures called **subspaces**, how certain subsets of vectors can generate entire spaces (**spanning**), and how to determine whether one vector is dependent on others (**linear dependence** versus **linear independence**). These notions not only formalize what it means for a collection of vectors to fill a space (or a portion of it) but also lay the groundwork for designing and working with efficient representations of those spaces.

We begin with the concepts of **image** and **kernel** in a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$. The image (also called the column space) comprises all possible outputs of the transformation, while the kernel (or nullspace) is the set of all inputs that map to the zero vector. Both are prime examples of subspaces of \mathbb{R}^m and \mathbb{R}^n , respectively.

From there, we introduce the formal definition of a **subspace** of \mathbb{R}^n . A subspace is any set of vectors in \mathbb{R}^n that is closed under addition and scalar multiplication, meaning sums and scalar multiples of vectors in the set remain in the set. Identifying these subspaces is crucial, for instance, in analyzing solution sets of linear systems and transformations.

Next, we discuss the idea of the **span** of a set of vectors, demonstrating how taking linear combinations of a few vectors can give rise to entire subspaces. This leads to the concept of a **spanning set**, a collection of vectors whose linear combinations recreate the entire subspace in question. Understanding which (and how many) vectors actually suffice to generate a subspace is a step toward more advanced ideas like dimension.

Finally, we turn to the topics of **linear dependence** and **linear independence**. These properties reveal whether one vector in a set can be written as a linear combination of the others. A set is linearly dependent if at least one vector is redundant, while a linearly independent set has no such redundancy. Checking for linear independence typically involves setting up a linear combination that yields the zero vector; if the only solution is all zero coefficients, the set is independent. This distinction shapes the way we reduce or optimize spanning sets and reveals the essence of a vector space's most efficient basis.

Overall, these ideas—image, kernel, subspaces, span, and linear independence—are at the heart of linear algebra. They clarify how different vector sets form coherent structures and pave the way for higher-level topics such as rank, dimension, and bases, all of which are built upon the foundation laid out here.

Recall:

Definition 3.1

A set, V , is a vector space over the real numbers \mathbb{R} if there are maps:

- $\mathbb{R} \times V \rightarrow V$: Denoted by $\alpha \mathbf{v}$, where $\alpha \in \mathbb{R}$ and $\mathbf{v} \in V$.
- $V \times V \rightarrow V$: Denoted by $\mathbf{u} + \mathbf{v}$, $\forall \mathbf{u}, \mathbf{v} \in V$

Definition 3.2

A linear transformation, $T : V \rightarrow W$ is a function from vector space V to a vector space W , such that, for any real numbers a_1, a_2 and v_1, v_2 , we have:

$$T(a_1 v_1 + a_2 v_2) = a_1 T(v_1) + a_2 T(v_2)$$

That is the additivity and homogeneity properties hold.

3.1 Subspaces

Definition 3.3

A non empty subset \mathbf{V} of \mathbb{R}^n is a **subspace** if:

- **Closed under scalar multiplication:** $\alpha \mathbf{u} \in \mathbf{V}$, $\mathbf{u} \in \mathbf{V}$, $\alpha \in \mathbb{R}$
- **Closed under addition:** $\mathbf{u} + \mathbf{v} \in \mathbf{V}$, $\mathbf{u}, \mathbf{v} \in \mathbf{V}$

Given some linear transformation $T : V \rightarrow W$, there are naturally occurring subspaces of both V and W , namely the **kernel** and **image**.

3.2 The Image and Kernel of a Linear Transformation

Let: $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, defined by matrix $M_T = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 5 \\ 1 & 1 & 1 \end{bmatrix}$

3.2.1 Image of T

Definition 3.4

$$\text{Im}(T) = \{w \in W, \text{there exists } v \in V \text{ such that } T(v) = w\}$$

The **image** of T , denoted $\text{im}(T)$ is the set of all vectors, w that can be written as $T(\mathbf{v})$, for some $\mathbf{v} \in \mathbb{R}^3$. Algebraically, this is the column space of the matrix M_T . **Explicitly:**

$$\text{im}(T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 5 \\ 1 \end{bmatrix} \right\}$$

3.2.2 Kernel of T

Definition 3.5

$$\ker(T) = \{v \in V : T(v) = 0\}$$

The **kernel** of T , denoted $\ker(T)$ is the set of all vectors $\mathbf{v} \in \mathbb{R}^3$ such that $T(\mathbf{v}) = \mathbf{0}$. Algebraically, this is the **null space** of the matrix M_T . To find the kernel solve:

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 5 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \xrightarrow{\text{Row reducing}} \begin{bmatrix} 1 & 2 & 0 & | & 0 \\ 0 & -5 & 5 & | & 0 \\ 1 & 1 & 1 & | & 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 2 & | & 0 \\ 0 & 1 & -1 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

$$\text{General Solution: } \begin{bmatrix} a \\ b \\ c \end{bmatrix} = t \cdot \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix} \quad \text{where } t \in \mathbb{R}$$

So the solution is is:

$$\ker(T) = \left\{ t \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix} : t \in \mathbb{R} \right\}$$

This is a one-dimensional subspace of \mathbb{R}^3 : all scalar multiples of a single nonzero vector. Geometrically, that's a line through the origin. Every vector in the kernel lies along that line.

3.2.3 Column Dependence

The fact the kernel is one-dimensional implies that the columns of M_T are linearly dependent. specially, the third column is a linear combination of the first two:

$$-2 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + 1 \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} + 1 \begin{bmatrix} 0 \\ 5 \\ 1 \end{bmatrix} \xrightarrow{\text{Rearranging}} 2 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} - 1 \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 5 \\ 1 \end{bmatrix}$$

Thus, the image of T is the plane:

$$\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} \right\}$$

Examples of subspaces

- $\{(x, y, z) \in \mathbb{R}^3 : x + y + z = 1\}$ is not a subspace of \mathbb{R}^3 . the vectors $(1, 0, 0)$ and $(0, 1, 0)$ are in the set, but their sum $(1, 1, 0)$ is not in the set.
- $\{(x, y, z) \in \mathbb{R}^3 : (x, y, z) \cdot (1, 2, 3) = 0\}$ is a subspace of \mathbb{R}^3 .
- $\{(x, y, z) \in \mathbb{R}^3 : (x, y, z) \cdot (1, 2, 3) \neq 0\}$ is not a subspace of \mathbb{R}^3 .
- The kernel of any linear transformation is a subspace of \mathbb{R}^n .
- The image of any linear transformation is a subspace of \mathbb{R}^n .

3.3 The span : how to make subspaces

Definition 3.6

Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be a finite subset of \mathbb{R}^n .

The **S**, denoted $\langle S \rangle$ is the set of all linear combinations of the vectors in S :

$$\langle S \rangle = \{a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_k\mathbf{v}_k : a_i \in \mathbb{R}\}$$

The span, $\langle S \rangle$ is a subspace of \mathbb{R}^n .

Proof that $\langle S \rangle$ is a subspace of \mathbb{R}^n

- **Closed under scalar multiplication:**

Let $u \in \langle S \rangle$ and $\alpha \in \mathbb{R}$ (that is, \mathbf{u} can be written as a combination of the vector in S)

We need to show that cu is a linear combination of v_1, \dots, v_k . We have $u = a_1v_1 + a_2v_2 + \dots + a_kv_k$ for some $a_i \in \mathbb{R}$. Then:

$$cu = c(a_1v_1 + a_2v_2 + \dots + a_kv_k) = (ca_1)v_1 + (ca_2)v_2 + \dots + (ca_k)v_k$$

so $cu \in \langle S \rangle$.

- **Closed under addition:**

Let $\mathbf{u}, \mathbf{v} \in \langle S \rangle$. Then $\mathbf{u} = a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_k\mathbf{v}_k$ and $\mathbf{v} = b_1\mathbf{v}_1 + b_2\mathbf{v}_2 + \dots + b_k\mathbf{v}_k$ for some $a_i, b_i \in \mathbb{R}$. We see that:

$$u + v = (a_1 + b_1)\mathbf{v}_1 + (a_2 + b_2)\mathbf{v}_2 + \dots + (a_k + b_k)\mathbf{v}_k$$

So S is closed under addition.

3.4 Spanning sets

Definition 3.7

A set of vectors, (v_1, \dots, v_n) span the vector space, V , if given any vector $v \in V$, there are scalar $a_1, \dots, a_n \in \mathbb{R}$ with:

$$v = a_1v_1 + a_2v_2 + \dots + a_nv_n$$

Example

The set $\{e_1, e_2, e_3\}$ is a spanning set of \mathbb{R}^3 . We know that:

$$e_1 = [1, 0, 0], \quad e_2 = [0, 1, 0], \quad e_3 = [0, 0, 1]$$

We can represent every element of \mathbb{R}^3 as a linear combination of e_1, e_2, e_3 :

$$\begin{bmatrix} 2 \\ -3 \\ 4 \end{bmatrix} = 2e_1 - 3e_2 + 4e_3$$

Remark A set S of three column vectors in \mathbb{R}^3 is a spanning set of \mathbb{R}^3 if and only if the three vectors are linearly independent. This occurs only if the 3×3 matrix whose columns are the three vectors has S as an inverse.

3.4.1 Questions about spanning sets

- Does \mathbb{R}^3 have a spanning set fewer than three vectors?
 - **No.** A spanning set for \mathbb{R}^3 must contain at least three linearly independent vectors, since the dimension of \mathbb{R}^3 is 3. Fewer than three vectors cannot span all of \mathbb{R}^3 .
- Does every spanning set of \mathbb{R}^3 have three vectors?
 - **No.** A spanning set can have more than three vectors, but not necessarily exactly three. Redundant vectors (linearly dependent ones) can be included, so a spanning set might have more than three vectors.
- Does every spanning set of \mathbb{R}^3 contain one with exactly three elements?
 - **Yes.** Every spanning set of \mathbb{R}^3 contains a basis, and since the dimension is 3, there exists a subset of exactly three linearly independent vectors that still span \mathbb{R}^3 .
- If V is a subspace of \mathbb{R}^3 does V have a spanning set with at most three elements?
 - **Yes.** Any subspace of \mathbb{R}^3 has a basis, and since \mathbb{R}^3 has dimension 3, the basis of any of its subspaces can have at most 3 elements. Hence, every subspace can be spanned by at most three vectors.

- If V is a proper subspace of \mathbb{R}^3 , does V have a spanning set with fewer than three elements?
 - **Yes.** A proper subspace of \mathbb{R}^3 has dimension less than 3, so it can be spanned by fewer than three vectors.

3.5 Linear Dependence and Linear Independence

Definition 3.8

A set of at least two vectors in \mathbb{R}^n is **linearly dependent** if one of its elements is a linear combination of the others.

A set of vectors in \mathbb{R}^n is **linearly independent** if it is not linearly dependent.

For a subset $\{v_1, \dots, v_k\}$ of \mathbb{R}^n , suppose that v_k is a linear combination of $\{v_1, \dots, v_{k-1}\}$. Then every linear combination of $\{v_1, \dots, v_k\}$ is **already a linear combination** of v_1, \dots, v_{k-1} :

$$\langle v_1, \dots, v_k \rangle = \langle v_1, \dots, v_{k-1} \rangle$$

If we are interested in the span of $\{v_1, \dots, v_k\}$, we can throw away v_k and this wouldn't change the span.

Linear independence means that throwing away any element of the set **shrinks the span**

Example 3.1

$$\begin{array}{cccccc} x_1 & + & 3x_2 & + & 5x_3 & - & 9x_4 = 5 \\ 3x_1 & - & x_2 & - & 5x_3 & + & 13x_4 = 5 \\ 2x_1 & - & 3x_2 & - & 8x_3 & + & 18x_4 = 1 \end{array} \Rightarrow \left[\begin{array}{cccc|c} 1 & 3 & 5 & -9 & 5 \\ 3 & -1 & -5 & 13 & 5 \\ 2 & -3 & -8 & -18 & 51 \end{array} \right] \Rightarrow \left[\begin{array}{cccc|c} 1 & 0 & -1 & 3 & 2 \\ 0 & 1 & 2 & -4 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

The three equations of the system form a linearly dependent set. One row was eliminated by adding a linear combination of the other two rows. Thus, all the information in the system was contained in the first two equations.

The non-zero rows of the RREF are linearly independent, they span the row space of the matrix. The row space is the subspace of \mathbb{R}^5 spanned by the rows of the matrix.

3.6 Test for linear independence

Definition 3.9

Vectors (v_1, \dots, v_n) in a vector space V are linearly independent, if whenever:

$$a_1 v_1 + \dots + a_n v_n = 0$$

, it must be the case that all scalars a_1, \dots, a_n must be zero. If $\forall i \rightarrow a_i = 0$, then the set is linearly independent. If not, the set is linearly dependent.

Intuitively, a collection of vectors are linearly independent if they all point in different directions.

Example 3.2

Decide whether the set $\{[1, 0, 1], [1, 0, -1], [1, 1, 1]\}$ is linearly independent or dependent.

To solve, we use ERO and find:

$$\left[\begin{array}{ccc} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & -1 & 1 \end{array} \right] \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \Rightarrow a = b = c = 0$$

The set is linearly independent

3.7 Key Takeaways

Key Takeaways 3.1

- **Image of a linear transformation** T is the set of all vectors in the co-domain (W) that can be obtained by applying T to vectors in the domain (V), we also call this the **column space** of T .
- **Kernel of a linear transformation** T is the set of all vectors in the domain (V) that are mapped to the zero vector in the co-domain (W), we also call this the **null space** of T .
- **Subspace:** A non-empty subset of \mathbb{R}^n that is closed under addition and scalar multiplication.

$$u + v \in \mathbb{V}, \quad \alpha u \in \mathbb{V} \quad u, v \in \mathbb{V}, \alpha \in \mathbb{R}$$

- **Span:** The set of all possible linear combinations of a finite subset of \mathbb{R}^n .
- **Span is a subspace of \mathbb{R}^n :** by the criteria of a subspace (closed under addition and scalar multiplication)
- **Spanning sets:** S is a spanning set of V (or S spans V) if every element of V is a linear combination of the elements of S - where V is a subspace of \mathbb{R}^n and S is a subset of V .
- **Linear dependence:** A set of at least two vectors in \mathbb{R}^n is linearly dependent if one of its elements is a linear combination of the others.
- **Linear independence:** A set of vectors in \mathbb{R}^n is linearly independent if it is not linearly dependent, throwing away any element of the set shrinks the span.
 - S is linearly independent only if no element of S is a linear combination of the other elements of S - equivalently if no proper subset of S spans the same space as S .

4 Basis, Dimension, and Rank

Introduction

This section deepens our investigation of vector spaces by addressing how to describe them succinctly and measure their “size.” First, we focus on the notion of a **finite dimensional** space: one that can be generated by a finite set of vectors. When that set of generators (or **spanning set**) is as small as possible while still spanning the entire space, we call it a **basis**. Crucially, a basis is both a minimal spanning set and a set of vectors that are linearly independent—no vector is redundant.

The number of vectors in any basis of a space defines its **dimension**. That this number is well-defined follows from important results like the Steinitz Replacement Lemma, which tells us that if you have a linearly independent set of the same size as a known basis, it must also be a basis. Consequently, every basis of a finite dimensional vector space has the same cardinality.

Moving from abstract vector spaces to matrices, we explore the concept of **rank**, which extends the idea of dimension to an $m \times n$ matrix. The rank is the dimension of either the row space or the column space of the matrix—two apparently different subspaces of \mathbb{R}^n and \mathbb{R}^m , respectively, but which always share the same dimension. Hence, the matrix’s **row rank** equals its **column rank**. This single integer is a powerful indicator of how many degrees of freedom the matrix transformation has, or equivalently, how many independent rows and columns the matrix contains.

Understanding basis, dimension, and rank is fundamental because they illuminate why some linear systems have unique solutions, why others have infinitely many solutions, and why still others have none. They also provide a systematic way to reduce large sets of vectors to essential generators, simplify linear transformations, and reason about the capacity or complexity of a given space. Taken together, these concepts form a bedrock for more advanced topics in linear algebra, such as diagonalization, eigenvalues, and beyond.

4.1 Finite Dimensional Spaces

Definition 4.1

A vector space V is said to be finite dimensional if there exists a finite set of vectors $\{v_1, \dots, v_k\} \subseteq V$ such that every vector in V can be expressed as a linear combination of these vectors. That is, the set spans V .

Examples

- \mathbb{R}^n is finite dimensional with $\{e_1, \dots, e_n\}$ as a spanning set. The dimension of \mathbb{R}^n is n .
- The space of $m \times n$ real matrices, denoted $M_{m \times n}(\mathbb{R})$ is finite dimensional. A spanning set is the collection of matrices with a single 1 in each position and 0s elsewhere (denoted $E_{i,j}$).

- An example of an infinite dimensional space is the set, $\mathbb{R}[x]$, of all polynomials with real coefficients. This set is infinite dimensional because it contains an infinite number of linearly independent vectors, such as $\{1, x, x^2, \dots\}$.

4.2 Basis: The Structural Backbone

Definition 4.2

A basis of a vector space is a set of vectors that satisfies two conditions:

- Linearly independent: No vector in the set can be expressed as a linear combination of the others.
- Spanning: The set of vectors can be combined to form any vector in the space.

Equivalently, (v_1, \dots, v_n) , form a basis for the vector space, V , if for any given vector $v \in V$, there are unique scalars $a_1, \dots, a_n \in \mathbb{R}$ such that:

$$v = a_1 v_1 + a_2 v_2 + \dots + a_n v_n$$

Definition 4.3

The dimension of a vector space, V , denoted $\dim(V)$ is the number of elements in a basis.

"I can represent everything, using only these, and no fewer."

A basis is both the minimal spanning set (remove any vector, and it no longer spans) and the maximal linearly independent set (add any vector, and it becomes linearly dependent)

Examples:

- $\{e_1, \dots, e_n\}$ is a basis for \mathbb{R}^n .
- $\{(1, 3), (1, 4)\}$ is a basis for \mathbb{R}^2 .
- If S is a finite spanning set of a vector space V , then S contains a basis of V . If S is not linearly independent, then some $v \in S$ is a linear combination of the other elements of S . Throwing away v leaves a smaller set that still spans V . This process can be repeated until a basis is obtained.

4.3 Steinitz Replacement Lemma

This is the core result guaranteeing that any two bases of a finite-dimensional space have the same number of elements. You can always replace vectors from a spanning set with vectors from an independent set, one at a time, while preserving the ability to span the space

Lemma 4.1

Let V be a vector space that has a basis with n elements. Then every linearly independent set with n elements in V is a basis for V .

Proof (for $n = 3$)

Suppose $B = \{b_1, b_2, b_3\}$ is a basis of V and let $\{y_1, y_2, y_3\}$ be a linearly independent subset of V .

Step 1.

$y_1 = a_1 b_1 + a_2 b_2 + a_3 b_3$ for scalars a_1, a_2, a_3 , not all zero. We can assume (after maybe relabelling the b_i), that $a_1 \neq 0$. Then

$$b_1 = a_1^{-1} y_1 - a_1^{-1} a_2 b_2 - a_1^{-1} a_3 b_3.$$

So $b_1 \in \langle y_1, b_2, b_3 \rangle$ and $\{y_1, b_2, b_3\}$ spans V . (Note that we have to use the fact that we can divide by non-zero scalars to write b_1 as a linear combination of y_1, b_2, b_3 .)

Step 2.

Now $y_2 \in \langle y_1, b_2, b_3 \rangle$ and y_2 is not a scalar multiple of y_1 (because $\{y_1, y_2, y_3\}$ is linearly independent).

So b_2 (or b_3) has non-zero coefficient in any description of y_2 as a linear combination of y_1, b_2, b_3 .

Replace again: $\{y_1, y_2, b_2\}$ spans V .

Step 3.

Same reasoning: we can replace b_2 with y_3 to conclude $\{y_1, y_2, y_3\}$ spans V .

Conclusion: $\{y_1, y_2, y_3\}$ is a basis of V .

4.4 Consequences of the replacement theorem

Theorem 4.1

Let V be a vector space that has a basis with n elements.

Then every linearly independent set with n elements in V is a basis for V .

- A linearly independent set in V cannot have more than n elements, if V has a spanning set with n elements.
- A spanning set in V must have at least n elements, if V has a linearly independent set with n elements.

Concept 4.1

The number of elements of a linearly independent set cannot exceed the number in a spanning set. Every spanning set has at least as many elements as the biggest independent set.

4.5 Every basis has the same number of elements

Let V be a finite dimensional vector space and let B and B' be the bases of V . Then:

- B is linearly independent and B' is a spanning set, so B has **at most** as many elements as B' .
- B is a spanning set and B' is linearly independent, so B has **at least** as many elements as B' .

It follows that B and B' have the same number of elements.

Definition 4.4

The dimension of V is the number of elements in a basis of V .

Note: Every vector space that has a finite spanning set has a finite basis (since we can discard elements from a finite spanning set until a basis remains).

Examples:

- The set $\{1, x, x^2, x^3\}$ is a basis for the vector space P_3 of all polynomials of degree at most 3 with real coefficients.

It is linearly independent because the only way to write the zero polynomial as

$$a_3x^3 + a_2x^2 + a_1x + a_0 = 0$$

is by taking $a_0 = a_1 = a_2 = a_3 = 0$.

Another basis of P_3 , preferable for some applications, consists of the first four Legendre polynomials:

$$\{1, x, \frac{1}{2}(3x^2 - 1), \frac{1}{2}(5x^3 - 3x)\}.$$

- The **row space** of an $m \times n$ matrix is the subspace of \mathbb{R}^n spanned by its rows. When we reduce a matrix to row-reduced echelon form (RREF), we are computing a basis of its row space.
- In \mathbb{R}^2 , the reflection in the line $y = 2x$ sends:

$$(1, 0) \mapsto \left(-\frac{3}{5}, \frac{4}{5}\right), \quad (0, 1) \mapsto \left(\frac{4}{5}, \frac{3}{5}\right).$$

Its standard matrix is:

$$\begin{bmatrix} -\frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{bmatrix}.$$

The same reflection sends:

$$(1, 2) \mapsto (1, 2), \quad (2, -1) \mapsto (-2, 1).$$

It is easier to describe this transformation in terms of the basis:

$$\left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}.$$

4.6 Row rank and column rank

Let $A \in \mathbb{A}^{m \times n}$ be a matrix. We define:

- Row space: The span of the rows \rightarrow subspace of \mathbb{R}^n .
- Column space: The span of the columns \rightarrow subspace of \mathbb{R}^m .
- Row rank r : The dimension of the row space
- Column rank c : The dimension of the column space

Rank = dimension of the image = number of pivot columns = number of nonzero rows in RREF.

4.7 Row rank = column rank

Theorem 4.2

The row rank and column rank are the same for every matrix

Therefore, we refer to this common value simply as the **rank** of the matrix.

Let A be an $m \times n$ matrix. The row rank r is the number of non-zero rows in the RREF of A , and the column rank c is the number of linearly independent columns of A . To show that $c \leq r$, consider a basis for the row space of A and arrange its vectors as the rows of an $r \times n$ matrix P . Since every row of A is a linear combination of the rows of P , there exists an $m \times r$ matrix Q such that

$$A = QP.$$

It follows that each column of A is a linear combination of the r columns of Q , implying that $\dim(\text{col}(A)) \leq r$. Hence, $c \leq r$.

Conversely, to show that $r \leq c$, take a basis for the column space of A and arrange its vectors as the columns of an $m \times c$ matrix P' . Since every column of A is a linear combination of the columns of P' , there exists a $c \times n$ matrix Q' such that

$$A = P'Q'.$$

Therefore, each row of A is a linear combination of the c rows of Q' , and $\dim(\text{row}(A)) \leq c$. Hence, $r \leq c$. Combining both inequalities, we conclude that $r = c$, i.e., the row rank and column rank are equal.

Example 4.1

Step 1: Determine the Rank. Reduce A to its reduced row echelon form (RREF):

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 1 & 1 & 1 \end{bmatrix} \implies \text{RREF}(A) = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}.$$

There are 2 non-zero rows, so the row rank is $r = 2$. Examining the columns, we observe that the third column is a linear combination of the first two, so the column rank $c = 2$ as well.

Step 2: Show that $c \leq r$ via $A = QP$. Choose a basis for the row space from the non-zero rows of RREF

$$A = QP = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 1 & 1 & 1 \end{bmatrix}.$$

Thus, each column of A is a linear combination of the $r = 2$ columns of Q , implying that $\dim(\text{col}(A)) \leq r$.

Step 3: Show that $r \leq c$ via $A = P'Q'$. Take a basis for the column space of A

$$A = P'Q' = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 1 & 1 & 1 \end{bmatrix}.$$

So, each row of A is a linear combination of the $c = 2$ rows of Q' , and $\dim(\text{row}(A)) \leq c$.

Conclusion. Since $c \leq r$ and $r \leq c$, it follows that $r = c = 2$. This example confirms that the row rank equals the column rank.

4.8 Key Takeaways

Key Takeaways 4.1

- **Finite dimensional space** is one that can be spanned by a finite set of vectors (akk uts vectors can be expresses as linear combinations of vectors)
- **Basis**
 - S is a basis f V if S is **linearly independent** S **spans** V
 - A basis is a minimal spanning set - meaning that every element is needed and does not contain a smaller spanning set.
 - A basis is a maximal linearly independent set - meaning that no element of S is a linear combination of the other elements of S .
- **Steinitz Replace Lemma:** "Every linearly independent set with n elements in V is a basis for V ." From the lemma, we conclude
 - A linearly independent in V cannot have more than n elements, if V has a spanning set with n elements.
 - A spanning set in V must have at least n elements, if V has a linearly independent set with n elements.
- **Dimension** of V is the number of elements in a basis of V . The dimension of \mathbb{R}^n is n .
- **Row rank** of A is the maximum number of linearly independent rows A .
- **Column rank** of A is the maximum number of linearly independent columns A .
- **Row rank = column rank** for every matrix.
- **Rank** of a matrix is the dimension of its row space or column space (since they are equal).

5 Coordinates, Rank-Nullity, and Change of Basis

Introduction

In this section, we extend our understanding of vector spaces by examining how to describe vectors with respect to an arbitrary basis (rather than always defaulting to the standard basis). This leads to the powerful notion of **coordinates**, which capture exactly how each vector in a space is formed as a linear combination of basis elements. Understanding coordinates is crucial for efficiently switching between different viewpoints on the same vector space—especially when one basis simplifies certain linear transformations more than another.

We begin by highlighting the uniqueness of coordinates in a given basis: any vector in a finite-dimensional space can be written in one and only one way as a linear combination of the basis vectors. From there, we see how to translate coordinates between different bases by way of a **change of basis** matrix. This concept underlies many practical calculations in linear algebra, including representing vectors or transformations in forms that reveal specific properties, such as diagonalization.

We also introduce the **Rank-Nullity Theorem**, one of the cornerstones of linear algebra, which states that for any linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, the dimension of the domain (n) is the sum of the dimension of the kernel and the rank of T . In other words, every vector in \mathbb{R}^n is either mapped to zero or contributes to the image; no other outcomes are possible.

Finally, we close by looking more deeply into how a linear map appears when expressed in different bases. A single transformation can have dramatically simpler matrices in certain bases—sometimes even diagonal. Understanding why and how to choose a basis that simplifies a map is a fundamental skill. Taken together, these ideas on coordinates, rank-nullity, and change of basis help us confidently move between abstract concepts (subspaces, images, kernels) and concrete matrix manipulations, ensuring we never lose sight of the geometry and structure behind the algebra.

5.1 Coordinates

Lemma 5.1

If $\{b_1, \dots, b_n\}$ is a basis of a vector space V , then every element of V has a unique expression of a linear combination of b_1, \dots, b_n :

Proof: Suppose, for some $v \in V$, that:

$$\begin{aligned}v &= a_1 b_1 + a_2 b_2 + \dots + a_n b_n, & a_i &\in \mathbb{R} \\v &= a'_1 b_1 + a'_2 b_2 + \dots + a'_n b_n, & a'_i &\in \mathbb{R}\end{aligned}$$

Then:

$$0_v = (a_1 - a'_1)b_1 + (a_2 - a'_2)b_2 + \cdots + (a_n - a'_n)b_n$$

Since B is linearly independent, we have:

$$a_i - a'_i = 0 \implies a_i = a'_i, \quad \forall i$$

Example 5.1

In \mathbb{R}^2 , the standard coordinates of $\begin{bmatrix} 4 \\ 3 \end{bmatrix}$ are $(4, 3)$.

With respect to the basis, $\left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$ the coordinates of $\begin{bmatrix} 4 \\ 3 \end{bmatrix}$ are $(2, -1)$. Which is saying:

$$\begin{bmatrix} 4 \\ 3 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + (-1) \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

5.2 Coordinates with respect to different bases

Let B be the ordered basis of \mathbb{R}^3 with elements:

$$b_1 = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix}, \quad b_2 = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}, \quad b_3 = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix} \implies B = \begin{bmatrix} 1 & 2 & 4 \\ 0 & -1 & 0 \\ 4 & 0 & 2 \end{bmatrix}$$

Given an element of \mathbb{R}^3 , say v , how do we find the B -coordinates of v ?

$$v = \begin{bmatrix} 2 \\ -3 \\ 4 \end{bmatrix}$$

We know

$$v = 2e_1 - 3e_2 + 4e_3 = [v]_B = 2[e_1]_B - 3[e_2]_B + 4[e_3]_B.$$

To find $[e_1]_B$:

$$e_1 = xb_1 + yb_2 + zb_3 = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix} x + \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} y + \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix} z = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \implies \begin{bmatrix} 1 & 2 & 4 \\ 0 & -1 & 0 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

This is saying that $[e_1]_B$ is the first column of the inverse of the matrix B . Write a matrix P which has b_1, b_2, b_3 as columns:

$$P = \begin{bmatrix} 1 & 2 & 4 \\ 0 & -1 & 0 \\ 4 & 0 & 2 \end{bmatrix} \implies P^{-1} = \begin{bmatrix} -\frac{1}{7} & -\frac{2}{7} & \frac{2}{7} \\ 0 & -1 & 0 \\ \frac{2}{7} & \frac{4}{7} & -\frac{1}{14} \end{bmatrix}$$

To find the B -coordinates of any $v \in \mathbb{R}^3$, we can multiply v on the left by P^{-1} :

$$\begin{bmatrix} 2 \\ -3 \\ 4 \end{bmatrix}_B = P^{-1}v = \begin{bmatrix} -\frac{1}{7} & -\frac{2}{7} & \frac{2}{7} \\ 0 & -1 & 0 \\ \frac{2}{7} & \frac{4}{7} & -\frac{1}{14} \end{bmatrix} \begin{bmatrix} 2 \\ -3 \\ 4 \end{bmatrix} = \begin{bmatrix} \frac{12}{7} \\ 3 \\ -\frac{10}{7} \end{bmatrix}$$

This is saying that $v = \frac{12}{7}b_1 + 3b_2 - \frac{10}{7}b_3$. P^{-1} is called the **change of basis matrix** from the standard basis to the basis B .

5.3 The Rank Nullity Theorem

The Rank-Nullity Theorem **relates the dimensions of the kernel, image and domain** of a linear transformation. The dimension of the image of a linear transformation is called the **rank** and the dimension of the the kernel is called the **nullity**. The rank of T is equal to the rank of matrix X , since the image of T is the column space of this matrix.

Theorem 5.1

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Where V and W are finite-dimensional vector spaces, over some field \mathbb{F} . Then:

$$\dim(\ker(T)) + \text{rank}(T) = n$$

Informally: The rank-nullity theorem says the full dimension of the domain must be accounted for in the combination of the kernel and image.

Proof:

1. Write k for $\dim(\ker(T))$ and let $\{b_1, \dots, b_k\}$ be a basis of $\ker(T)$.
2. Extend this to a basis: $\{b_1, \dots, b_k, b_{k+1}, \dots, b_n\}$ of \mathbb{R}^n .
3. Since T sends each b_i to 0, the image under T of every element of \mathbb{R}^n is a linear combination of $T(c_{k+1}), \dots, T(c_n)$.
4. Also, $\{T(c_{k+1}), \dots, T(c_n)\}$ is a linearly independent subset of \mathbb{R}^m . To see this suppose for some scalar a_{k+1}, \dots, a_n that $a_{k+1}T(c_{k+1}) + \dots + a_nT(c_n) = 0$. Then:

$$a_{k+1}c_{k+1} + \dots + a_n c_n \in \ker T \implies a_{k+1}c_{k+1} + \dots + a_n c_n \in \langle b_1, \dots, b_k \rangle.$$

Since $\{b_1, \dots, b_k, c_{k+1}, \dots, c_n\}$ is linearly independent in \mathbb{R}^n , this means that $a_{k+1}c_{k+1} + a_{k+2}c_{k+2} + \dots + a_n c_n = 0$ for each $a_j = 0$.

5. It follows that $\{T(c_{k+1}), \dots, T(c_n)\}$ is a basis for the image of T , so this image has dimensions $n - k$ as required.

5.4 Linear transformations and change of basis

Definition 5.1

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation and let $B = \{b_1, \dots, b_n\}$ be a basis of \mathbb{R}^n .

The **matrix of T with respect to the basis B** is the $n \times n$ matrix that has the B -coordinates of $T(b_1), \dots, T(b_n)$ as its n columns. This matrix M satisfies

$$[T(v)]_B = M[v]_B, \quad \forall v \in \mathbb{R}^n.$$

Example 5.2

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation defined by $v \rightarrow Av$ and B be the ordered basis of \mathbb{R}^3 with elements:

$$A = \begin{bmatrix} -2 & 2 & 1 \\ 4 & 5 & -1 \\ -4 & -8 & 3 \end{bmatrix} \quad \left\| \right\| \quad b_1 = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix}, \quad b_2 = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}, \quad b_3 = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix} \implies B = \begin{bmatrix} 1 & 2 & 4 \\ 0 & -1 & 0 \\ 4 & 0 & 2 \end{bmatrix}$$

A diagonal representation of the matrix A' of T with respect to the basis B

$$T(b_1) = \begin{bmatrix} -2 & 2 & 1 \\ 4 & 5 & -1 \\ -4 & -8 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 8 \end{bmatrix} = 2b_1 \implies [T(b_1)]_B = \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$$

$$T(b_2) = \begin{bmatrix} -2 & 2 & 1 \\ 4 & 5 & -1 \\ -4 & -8 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} -6 \\ 3 \\ 0 \end{bmatrix} = -3b_2 \implies [T(b_2)]_B = \begin{bmatrix} 0 \\ -3 \\ 0 \end{bmatrix}$$

$$T(b_3) = \begin{bmatrix} -2 & 2 & 1 \\ 4 & 5 & -1 \\ -4 & -8 & 3 \end{bmatrix} \begin{bmatrix} 0 \\ -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ -7 \\ 14 \end{bmatrix} = 7b_3 \implies [T(b_3)]_B = \begin{bmatrix} 0 \\ 0 \\ 7 \end{bmatrix}$$

The matrix of A' of T with respect to B is **diagonal**. For describing this transformation, T , the basis B is preferable to the standard basis.

$$A' = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 7 \end{bmatrix}$$

This means for any $v \in \mathbb{R}^3$:

$$\underbrace{[T(v)]_B}_{\text{B-coordinates of } T(v)} = \underbrace{A'[v]_B}_{\text{matrix-vector product}}$$

5.5 More on Change of Basis

Let P be the matrix with the basis vectors from B as columns. As we've seen, P^{-1} is the change of basis matrix from the standard basis to the basis B .

For any element, $v \in \mathbb{R}^n$, its B -coordinates are given by:

$$[v]_B = P^{-1}v$$

Equivalently, if we start with the B -coordinates, then the standard coordinates of v are given by:

$$v = P[v]_B$$

So P itself, is the change of basis matrix from the basis B to the standard basis.

5.6 Key Takeaways

Key Takeaways 5.1

- **Coordinates** of a vector v with respect to a basis B are the coefficients of the linear combination of the basis vectors that yield v .
- **Lemma 5.1:** *If $\{b_1, \dots, b_n\}$ is a basis of a vector space V , then every element of V has a unique expression of a linear combination of b_1, \dots, b_n .*
- **Change of basis matrix** a matrix that allows you to express the coordinates of a vector with respect to a new basis, given its coordinates in the original basis
- **Rank-Nullity Theorem:** *Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then $\dim(\ker(T)) + \text{rank}(T) = n$.*

6 Similarity, Eigenvalues, and Diagonalizability

Recall, given a basis for an n dimensional vector space, V , we can represent a linear transformation:

$$T : V \rightarrow V$$

as an $n \times n$ matrix A . Unfortunately, if we choose a different basis for V , the matrix A representing T will change.

We aim to clearly explain when two matrices represent the same linear transformation, just in different bases.

k

We want to see two matrices are similar when they represent the same linear transformation. Choose two bases for the vector space V , and W : $\{v_1, \dots, v_n\}$ and $\{w_1, \dots, w_n\}$.

Given the v basis, we can write any vector $z \in V$ as a linear combination of the basis vectors, and similar for the w basis:

$$z = a_1 v_1 + \dots + a_n v_n = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \quad z = b_1 w_1 + \dots + b_n w_n = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

where a_i, b_i are unique scalars.

Let A be the matrix representing the linear transformation, T , for the v basis, and B be the matrix representing the same linear transformation for the w basis:

$$T(z) = A \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \quad \text{and} \quad T(z) = B \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

Now, we introduce a matrix P , that maps the v basis to the w basis, called the **change-of-basis** matrix:

$$P \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix} \quad \Rightarrow \quad \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = P^{-1} \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

k

We want to see two matrices are similar when they represent the same linear transformation. Choose two bases for the vector space V , and W : $\{v_1, \dots, v_n\}$ and $\{w_1, \dots, w_n\}$ and let A be the matrix representing the linear transformation, T , for the v basis, and B be the matrix representing the same linear transformation for the w basis.

Given the v basis, we can write each vector $z \in V$ as an $n \times 1$ column vector:

$$z = a_1 v_1 + a_2 v_2 + \dots + a_n v_n = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

where a_i are unique scalars. Similar for the w basis:

$$z = b_1 w_1 + b_2 w_2 + \cdots + b_n w_n = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

where b_i are unique scalars.

We know, the linear transformation T acts as:

$$T(z) = T(a_1 v_1 + a_2 v_2 + \cdots + a_n v_n) = A \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

Then, we can introduce a matrix P that maps the v basis to the w basis, called the **change-of-basis** matrix:

$$P \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

So applying the transformation, then converting to the w basis gives:

$$P \left(A \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \right)$$

On the other hand, if B represent the transformation of T in the w basis, we can write:

$$B \left(P \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \right)$$

So:

$$PA = BP \implies A = P^{-1}BP$$

Our goal is to relate the matrices A and B — that is, to find how the matrix representation the shared linear transformation T changes when we change from the v -basis to the w -basis.

6.1 Similarity (The relation of A and A')

Starting with A , the matrix $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with respect to the standard basis, to we find A' , the matrix of T with respect to the basis B we:

- **Take a vector $v \in \mathbb{R}^3$ written in B -coordinates as the column $[v]_B$.**
- **Convert to standard coordinates:** take the product $P[v]_B$.
- **Apply T :** Left multiply by A , to get $AP[v]_B$. This column is the standard coordinates of $T(v)$.
- **Convert to B -coordinates:** left multiply by P^{-1} (change of basis matrix from standard to B) to get $P^{-1}AP[v]_B$. This column has the B -coordinates of $T(v)$.
- **Conclusion:** For any element $v \in \mathbb{R}^3$, the B -coordinates of $T(v)$ are given by:

$$[T(v)]_B = P^{-1}AP[v]_B$$

The B matrix of T is: $P^{-1}AP$, where P has the elements of B as columns.

6.2 Similar Matrices

Definition 6.1

Two matrices A and B are **similar** if there exists an invertible matrix P such that:

$$B = P^{-1}AP$$

Notes:

- Two distinct matrices are similar only if they **represent the same linear transformation** with respect to different bases.
- We can't tell if two square matrices are similar just by looking. Instead, we look at the **trace** of the matrix, the sum of the diagonal elements, if two matrices have the same trace, they are similar.
- Similar matrices have some features in common, such as the same determinant, the same eigenvalues, etc (more on this later).
- Our example showed that the matrix:

$$A = \begin{bmatrix} -2 & 2 & 1 \\ 4 & 5 & -1 \\ -4 & -8 & 3 \end{bmatrix}$$

is similar to the diagonal matrix:

$$\text{diag}(2, -3, 7) = A' = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 7 \end{bmatrix}$$

In this situation, we say that A is **diagonalizable**.

6.3 Two interpretations of diagonalizability

Interpretation 1:

From the diagonal form of A' we have:

$$T(b_1) = 2b_1, \quad T(b_2) = -3b_2, \quad T(b_3) = 7b_3$$

This means that each of the basis elements b_1, b_2, b_3 is mapped by T to a scalar multiple of itself. - each of them is an eigenvector of T

Interpretation 2:

We can rearrange the version of $P^{-1}AP = A'$ to get:

$$AP = P'A$$

Bearing in mind that that:

$$P = \begin{bmatrix} | & | & | \\ b_1 & b_2 & b_3 \\ | & | & | \end{bmatrix} \quad \text{and} \quad A' = \text{diag}(2, -3, 7)$$

this is saying that:

$$A \begin{bmatrix} | & | & | \\ b_1 & b_2 & b_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | \\ b_1 & b_2 & b_3 \\ | & | & | \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 7 \end{bmatrix} = \begin{bmatrix} | & | & | \\ Ab_1 & Ab_2 & Ab_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | \\ 2b_1 & -3b_2 & 7b_3 \\ | & | & | \end{bmatrix}$$

This means that $Ab_1 = 2b_1$, $Ab_2 = -3b_2$, $Ab_3 = 7b_3$, so that $B = \{b_1, b_2, b_3\}$ is a basis of \mathbb{R}^3 consisting of eigenvectors of A .

6.4 Eigenvalues and diagonalizability

Definition 6.2

An **eigenvector** of a square matrix A is a non zero column vector v , for which there exists a scalar λ such that:

$$Av = \lambda v$$

The scalar λ is called the **eigenvalue** of A corresponding to the eigenvector v .

The eigenvalues of A are the roots of its **characteristic polynomial**:

$$\det(xI_n - A) = 0$$

The **eigenspace** corresponding to a particular eigenvalue, λ , is the set of all vector, v , satisfying: $Av = \lambda v$. It is a subspace of the relevant \mathbb{R}^n of dimension at least 1.

The matrix $A \in M_n(\mathbb{R})$ is **diagonalizable** only if \mathbb{R}^n has a basis consisting of eigenvectors of A . In this case $P^{-1}AP$ is diagonal, where P is a matrix whose n columns are linearly independent eigenvectors of A . The diagonal entries of $P^{-1}AP$ are the corresponding eigenvalues.

6.5 Two examples of non-diagonalizability

For $A \in M_n(\mathbb{R})$, it does not always happen that \mathbb{R}^n has a basis consisting of eigenvectors of A .

Example 1:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

The matrix A is diagonalizable in $M_2(\mathbb{C})$, but not in $M_2(\mathbb{R})$. The matrix represents an anti-clockwise rotation of 90° in \mathbb{R}^2 . It does not fix any line in \mathbb{R}^2 . Its characteristic polynomial is:

$$\det(xI_2 - A) = \det \begin{bmatrix} x & 1 \\ -1 & x \end{bmatrix} = x^2 + 1 = 0$$

Example 2:

$$B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$


The matrix B is not diagonalizable in $M_2(\mathbb{C})$. The matrix represents **horizontal shear**. Its characteristic polynomial is:

$$\det(xI_2 - B) = \det \begin{bmatrix} x-1 & -1 \\ 0 & x-1 \end{bmatrix} = (x-1)^2 = 0$$

but its 1-dimensional eigenspace only consists of the X -axis. It does not have two linearly independent eigenvectors.

6.6 A Shear in \mathbb{R}^2

$$B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$



The linear transformation T described by B send $(x, y) \in \mathbb{R}^2$ to $(x+y, y)$. This is a **horizontal shear**: it shifts every point horizontally by its y -coordinate.

For every point $v \in \mathbb{R}^2$, $T(v)$ is on the same horizontal line as v . It follows that $T(v)$ is a scalar multiply of v , only if v lies on the X -axis; in this case, $T(v) = v$.

The characteristic polynomial of B (and T) is:

$$\det(xI_2 - B) = \det \begin{bmatrix} x-1 & -1 \\ 0 & x-1 \end{bmatrix} = (x-1)^2 = (\lambda-1)^2 = 0$$

The only eigenvalue is $\lambda = 1$, and it has **algebraic multiplicity** 2, meaning it appears twice as a root of the characteristic polynomial.

Its **geometric multiplicity** is only 1, meaning its corresponding eigenspace is only 1-dimensional - just the line $y = 0$.

6.7 Eigenvector for distinct eigenvalues are independent

Theorem 6.1

Let $A \in M_n \mathbb{R}$ and v_1, \dots, v_k be eigenvector of A in \mathbb{R}^n corresponding to distinct eigenvalues $\lambda_1, \dots, \lambda_k$ of A . Then $\{v_1, \dots, v_k\}$ is a linearly independent subset of \mathbb{R}^n .

Proof (for $k = 3$): Note that no two v_1, v_2, v_3 are multiples of each other, since they correspond to distinct eigenvalues.

Suppose: $a_1 v_1 + a_2 v_2 + a_3 v_3 = 0$ for some $a_i \in \mathbb{R}$. We want to show that $a_1 = a_2 = a_3 = 0$. Multiplying on the left by A gives:

$$a_1 A v_1 + a_2 A v_2 + a_3 A v_3 = 0 \implies a_1 \lambda_1 v_1 + a_2 \lambda_2 v_2 + a_3 \lambda_3 v_3 = 0$$

and multiplying the same equation by λ_1 gives:

$$a_1\lambda_1v_1 + a_2\lambda_1v_2 + a_3\lambda_1v_3 = 0$$

Subtract to get:

$$a_2(\lambda_1 - \lambda_2)v_2 + a_3(\lambda_1 - \lambda_3)v_3 = 0$$

Since v_2 and v_3 are linearly independent, and $\lambda_1 - \lambda_2 \neq 0$ and $\lambda_1 - \lambda_3 \neq 0$, it follows that $a_2 = a_3 = 0$ and hence $a_1 = 0$.

6.8 At most n distinct eigenvalues

The following consequence of the previous theorem shows that a matrix can have at most n distinct eigenvalues. We already knew this, since the eigenvalues are roots of a polynomial of degree n .

Theorem 6.2: Corollary

Let $A \in M_N(\mathbb{R})$. Then A has at most n distinct eigenvalues in \mathbb{R} .

Proof:

If A has k distinct eigenvalues, with corresponding eigenvectors $v_1, \dots, v_k \in \mathbb{R}^n$, then k cannot exceed the dimension of \mathbb{R}^n , since $\{v_1, \dots, v_k\}$ is a linearly independent set in \mathbb{R}^n .

Theorem 6.3: Corollary

If $A \in M_n(\mathbb{R})$ has n distinct eigenvalues, then A is diagonalizable.

Proof:

A set consisting of one eigenvector for each of the n eigenvalues is linearly independent, so hence is a basis.

6.9 Notes about determinants and characteristic polynomials

1. The **characteristic polynomial** of the square matrix $A \in M_n(\mathbb{R})$ is the determinant of $\lambda I_n - A$.
2. If t is a root of this polynomial, the t -eigenspace of A is the nullspace of the matrix $tI_n - A$.
3. The determinant of a diagonal or upper triangular matrix is the product of the entries on its main diagonal.
4. A square matrix is **block diagonal** if its non-zero entries are all contained in square blocks along its diagonal. The determinant of a block diagonal matrix is the product of the determinants of its diagonal blocks.
5. Similar matrices have the same characteristic polynomial and the same eigenvalues and eigenspace dimensions, since they represent the same linear transformation.

6.10 Multiplicity of Eigenvalues

Suppose that A and B are similar square matrices, so that $B = P^{-1}AP$ for some invertible matrix P . Then

$$\begin{aligned} \det(xI - B) &= \det(xI - P^{-1}AP) \\ &= \det(P^{-1}(xI)P - P^{-1}AP) \\ &= \det(P^{-1}(xI - A)P) \\ &= \det P^{-1} \det(xI - A) \det P \\ &= \det(xI - A). \end{aligned}$$

If two matrices have the same characteristic polynomial, they are not necessarily similar. For example $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ have the same characteristic polynomial but are not similar.

6.11 Multiplicity of Eigenvalues

Let λ be an eigenvalue of a matrix $A \in M_n(\mathbb{R})$. The **algebraic multiplicity** of λ is the number of times that λ occurs as a root of the characteristic polynomial. The **geometric multiplicity** is the dimension of the t -eigenspace of A .

Example 6.1

$$A = \begin{bmatrix} 3 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 0 & 4 \end{bmatrix}$$

The matrix has two distinct eigenvalues, 3 and 4. Both have algebraic multiplicity 2; the characteristic polynomial is $(\lambda - 3)^2(\lambda - 4)^2$. The 3-eigenspace has dimension 2, its elements are $\begin{bmatrix} a \\ b \\ 0 \\ 0 \end{bmatrix}$, for $a, b \in \mathbb{R}$.

The 4-eigenspace only has dimension 1, its elements are $\begin{bmatrix} 0 \\ 0 \\ c \\ 0 \end{bmatrix}$, for $c \in \mathbb{R}$.

This A is not diagonalizable since it does not have four independent eigenvectors.

6.12 Geometric multiplicity \leq Algebraic multiplicity

Theorem 6.4

The geometric multiplicity of an eigenvalue is at most equal to its algebraic multiplicity.

Proof:

Suppose that t has geometric multiplicity k as an eigenvalue of the square matrix $A \in M_n(\mathbb{R})$, and let $\{v_1, \dots, v_k\}$ be a basis for the t -eigenspace of A .

Extend this to a basis \mathcal{B} of \mathbb{R}^n , and let P be the matrix whose columns are the elements of \mathcal{B} . Then the first k columns of $P^{-1}AP$ have t in the diagonal position and zeros elsewhere.

It follows that $\lambda - t$ occurs at least k times as a factor of $\det(\lambda I_n - P^{-1}AP)$, so the algebraic multiplicity of t is at least k .

Theorem 6.5: Corollary

A matrix is diagonalizable if and only if the geometric multiplicity of each of its eigenvalues is equal to the algebraic multiplicity

6.13 Key Takeaways

Key Takeaways 6.1

- The B coordinates of a vector a linear transformation T is given by:

$$[T(v)]_B = P^{-1}AP[v]_B$$

- Two matrices are **similar** there exists an invertible matrix P such that $B = P^{-1}AP$.
- Eigenvector** a vector which when operated on by a given operator gives a scalar multiple of itself.
- Eigenvalue** the scalar multiple of an eigenvector.
- Characteristic polynomial** $\det(\lambda I_n - A)$ is a polynomial in λ whose roots are the eigenvalues of A .
- Eigenspace** the set of all vectors v such that $Av = \lambda v$ for a given eigenvalue λ .
- Diagonalizability** a matrix is diagonalizable if it has a basis of eigenvectors.
- Horizontal Shear** a shear mapping that shifts every point horizontally by its y -coordinate.
- Algebraic multiplicity** the number of times an eigenvalue appears as a root of the characteristic polynomial.
- Geometric multiplicity** the dimension of the eigenspace corresponding to an eigenvalue.
- The set of eigenvectors corresponding to distinct eigenvalues is linearly independent.
- Given a $n \times n$ matrix A , A has at most n distinct eigenvalues.
- If A has n distinct eigenvalues, then A is diagonalizable.
- The geometric multiplicity of an eigenvalue is \leq to its algebraic multiplicity.
- A matrix is diagonalizable if and only if the geometric multiplicity of each of its eigenvalues is equal to the algebraic multiplicity.

7 Inner Products, Orthogonality, and Projections

Introduction

In this section, we introduce the notion of an **inner product**, a tool that lets us measure angles, lengths, and distances within a vector space—much like the dot product does in \mathbb{R}^n . Once we have an inner product, we can talk about **orthogonality**, or perpendicular vectors, which lays the groundwork for a host of geometric insights and computational techniques.

First, we generalize the usual dot product to an arbitrary vector space by defining a function $\langle x, y \rangle$ that must satisfy certain symmetry, linearity, and positivity conditions. This innately geometric perspective extends to the concepts of **length** (or norm), **distance**, and **angle** in any such space. Orthogonality—where two vectors satisfy $\langle x, y \rangle = 0$ —then becomes the keystone for analyzing how vectors project and decompose relative to each other.

We examine how to build an **orthogonal basis**, a set of mutually perpendicular vectors that simplifies calculations significantly. The **Gram–Schmidt process** provides a systematic way to convert any basis into one that is orthogonal. Orthogonal bases enable us to define the **orthogonal projection** of a vector onto a subspace, i.e., the nearest point in the subspace to a given vector. This projection is at the heart of geometric approximations—particularly the solution of certain systems that may not have exact solutions.

Finally, we apply these ideas to the method of **least squares** for **overdetermined systems**, where there are more equations than unknowns. By projecting the system's right-hand side vector onto the column space of the coefficient matrix, we find an approximate solution that minimizes the total error. This illustrates just how powerful inner-product-based tools can be in both pure geometry and practical problem-solving contexts in data analysis, engineering, and beyond.

7.1 Inner Products

In \mathbb{R}^n , the dot product of the vectors $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and $y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ is defined as:

$$x \cdot y = x_1 y_1 + x_2 y_2 = x^T = y^T = y \cdot x$$

We can interpret the length ($\|x\|$) of x as the length of the line segment from origin to (x_1, x_2) , which by the Pythagorean theorem is:

$$\sqrt{x_1^2 + x_2^2} = \sqrt{x \cdot x}$$

Once we have a concept of length, we can define distance ($d(x, y)$) between two vectors x, y as the length of the line segment from x to y , which is:

$$d(x, y) = \|x - y\|$$

Similarly, from the Cosine Rule:

$$x \cdot y = \|x\| \cdot \|y\| \cdot \cos(\theta)$$

where θ is the angle between x and y .

We say that x and y are **orthogonal** (or $x \perp y$) if $x \cdot y = 0$.

So the scalar product encodes geometric information in \mathbb{R}^2 , and also provides a way to define length, distance and orthogonality. The scalar product is an example of an **inner product**.

7.2 Real Inner Products

A **Inner Product** on a vector space V is a function from $V \times V$ to \mathbb{R} that assigns an element of \mathbb{R} to every ordered pair of elements of V , and has the following properties.

1. **Symmetry:** $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$
2. **Linearity in both slots (bilinearity):** For all $x, y, z \in V$ and all $a, b \in \mathbb{R}$, we have $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ and $\langle x, ay + bz \rangle = a\langle x, y \rangle + b\langle x, z \rangle$.
3. **Non-negativity:** $\langle x, x \rangle \geq 0$ for all $x \in V$, and $\langle x, x \rangle = 0$ only if $x = 0_V$.

The ordinary scalar product on \mathbb{R}^n is the best known example of an inner product.

7.2.1 Examples of Inner Products

1. The ordinary scalar product on \mathbb{R}^n .
2. Let C be the vector space of all continuous real-valued functions on the interval $[0, 1]$. The analogue of the ordinary scalar product on C is the inner product given by

$$\langle f, g \rangle = \int_0^1 f(x)g(x) dx, \quad \text{for } f, g \in C.$$

3. On the space $M_{m \times n}(\mathbb{R})$, the **Frobenius inner product** or **trace inner product** is defined by

$$\langle A, B \rangle = \text{trace}(A^T B).$$

Note that $\text{trace}(A^T B)$ is the sum over all positions (i, j) of the products $A_{ij}B_{ij}$. So this is closely related to the ordinary scalar product, if the matrices A and B were regarded as vectors with mn entries over \mathbb{R} .

It is possible for a single vector space to have many different inner products defined on it, and if there is any risk of ambiguity we need to specify which one we are considering.

7.3 Length, Distance and Scaling in an Inner Product Space

Definition 7.1

We define the **length** or **norm** of any vector v by:

$$\|v\| = \sqrt{\langle v, v \rangle}$$

Definition 7.2

We define the distance between two vectors, u and v , by:

$$d(u, v) = \|u - v\|$$

Concept 7.1: Scaling

Every vector, v , and scalar, c , satisfy:

$$\|cv\| = |c| \cdot \|v\|$$

Since:

$$\|cv\| = \sqrt{\langle cv, cv \rangle} = \sqrt{c^2 \langle v, v \rangle} = |c| \cdot \|v\|$$

So we can adjust the norm of any element of V while preserving its direction, by multiplying by a scalar.

Definition 7.3

If v is a non-zero vector in an inner product space V , then

$$\hat{v} := \frac{1}{\|v\|} v$$

is a unit vector in the direction of v , called the **normalization** of v .

7.4 Orthogonality in an inner product space

Let V be a vector space with an inner product $\langle \cdot, \cdot \rangle$ (such as the ordinary scalar product).

Definition 7.4

We say that the vectors u and v are **orthogonal** (with respect to $\langle \cdot, \cdot \rangle$) if $\langle u, v \rangle = 0$.

All these definitions are consistent with “typical” geometrically motivated concepts of distance and orthogonality.

Examples

1. $(2, 5)$ and $(5, -2)$ are orthogonal with respect to the ordinary scalar product in \mathbb{R}^2 .
2. $\sin \pi x$ and $\cos \pi x$ are orthogonal with respect to the scalar product on the space of continuous functions on $[0, 1]$ defined in Lecture 18; this is saying that

$$\int_0^1 \sin(\pi x) \cos(\pi x) dx = 0 \quad \left(= \frac{1}{2\pi} \sin^2(\pi x) \Big|_0^1 \right).$$

7.5 Orthogonal Projection

Lemma 7.1

Let u and v be non-zero vectors in an inner product space V . Then it is possible to write (in a unique way) $v = au + v'$, where a is a scalar and v' is orthogonal to u .

- If v is orthogonal to u , take $a = 0$ and $v' = v$.
- If v is a scalar multiple of u , take $au = v$ and $v' = 0$.
- Otherwise, to solve for a and v' in the equation $v = au + v'$ (with $u \perp v'$), take the inner product with u on both sides. Then

$$\langle u, v \rangle = a\langle u, u \rangle + 0 \implies a = \frac{\langle u, v \rangle}{\|u\|^2}, \quad v' = v - \frac{\langle u, v \rangle}{\|u\|^2}u.$$

We can verify directly that the two components in this expression are orthogonal to each other.

Example

$$\text{Write } u = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad v = \begin{bmatrix} 6 \\ -2 \end{bmatrix} \quad \text{Then } u = \begin{bmatrix} \frac{3}{2} \\ \frac{3}{4} \end{bmatrix} + \begin{bmatrix} \frac{3}{2} \\ -\frac{11}{4} \end{bmatrix}$$

7.6 Orthogonal projection of one vector on another

Definition 7.5

For non-zero vectors u and v in an inner product space V , the vector

$$\frac{\langle u, v \rangle}{\|u\|^2}u$$

is called the projection of v on the 1-dimensional space spanned by u .

It is denoted by $\text{proj}_u(v)$ and it has the property that $v - \text{proj}_u(v)$ is orthogonal to u

Lemma 7.2

$\text{proj}_u(v)$ is the unique element of $\langle u \rangle$ whose distance from v is minimal.

Proof: Let au be a scalar multiple of u . Then

$$d(au, v)^2 = \langle au - v, au - v \rangle = a^2\langle u, u \rangle - 2a\langle u, v \rangle + \langle v, v \rangle$$

Regarded as a quadratic function of a , this has a minimum when its derivative is 0, i.e. when

$$2a\langle u, u \rangle - 2\langle u, v \rangle = 0, \quad \text{when } a = \frac{\langle u, v \rangle}{\|u\|^2}.$$

7.7 Orthogonal Bases (the Gram-Schmidt process)

Concept 7.2

Every finite-dimensional inner product space has an **orthogonal basis**

We can start with any basis $\{b_1, \dots, b_n\}$, and adjust the elements one by one (by subtracting off orthogonal projections of later vectors on earlier ones). The process ends with an orthogonal basis $\{v_1, \dots, v_n\}$.

1. Set $v_1 = b_1$, and

$$v_2 = b_2 - \text{proj}_{v_1}(b_2) = b_2 - \frac{\langle v_1, b_2 \rangle}{\langle v_1, v_1 \rangle}v_1.$$

Then the pairs b_1, b_2 and v_1, v_2 span the same space, and $v_1 \perp v_2$.

2. Write $v_3 = b_3 - \text{proj}_{v_1}(b_3) - \text{proj}_{v_2}(b_3)$. Then $\{v_1, v_2, v_3\}$ and $\{b_1, b_2, b_3\}$ span the same space, and $v_3 \perp v_1, v_3 \perp v_2$. To see this look at $\langle v_3, v_1 \rangle$ and $\langle v_3, v_2 \rangle$, noting that

$$v_3 = b_3 - \frac{\langle b_3, v_1 \rangle}{\langle v_1, v_1 \rangle}v_1 - \frac{\langle b_3, v_2 \rangle}{\langle v_2, v_2 \rangle}v_2.$$

3. Continue: at the k th step, form v_k by subtracting from b_k its projections on v_1, \dots, v_{k-1} .

7.8 Orthogonal projection on a subspace

The result of this process is a basis $\{v_1, \dots, v_n\}$ whose elements satisfy

$$\langle v_i, v_j \rangle = 0 \text{ for } i \neq j$$

We can adjust this basis to a **orthonormal basis** (consisting of orthogonal unit vectors) by replacing each v_i with its normalization \hat{v}_i . From the Gram–Schmidt process, we have

Theorem 7.1

If V is a finite-dimensional inner product space, then V has an orthogonal (or orthonormal) basis.

Now let W be a subspace of V , and let $v \in V$. The **orthogonal projection** of v on W , denoted $\text{proj}_W(v)$, is defined to be the unique element u of W for which

$$v = u + v',$$

and $v' \perp w$ for all $w \in W$.

7.9 Calculating the projection on a subspace

Example 7.1: I

\mathbb{R}^3 , find the unique point of the plane $W : x + 2y - z = 0$ that is nearest to the point $v : (1, 2, 2)$.

First find an orthogonal basis for W : for example $\{b_1, b_2\}$, where

$$b_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad b_2 = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}$$

Then

$$\begin{aligned} \text{proj}_W(v) &= \text{proj}_{b_1}(v) + \text{proj}_{b_2}(v) \\ &= \frac{\langle b_1, v \rangle}{\langle b_1, b_1 \rangle} b_1 + \frac{\langle b_2, v \rangle}{\langle b_2, b_2 \rangle} b_2 \\ &= \frac{3}{2} b_1 - \frac{3}{3} b_2 \\ &= \left(\frac{3}{2}, 0, \frac{3}{2} \right) - (1, -1, -1) = \left(\frac{1}{2}, 1, \frac{5}{2} \right) \end{aligned}$$

7.10 $\text{proj}_W(v)$ is the nearest point of W to v

Let $u = \text{proj}_W(v)$ and let w be any element of W . Note that $v - u$ is orthogonal to both w and u , hence to $w - u$. Then

$$\begin{aligned} d(v, w)^2 &= \langle v - w, v - w \rangle \\ &= \langle (v - u) + (u - w), (v - u) + (u - w) \rangle \\ &= \langle v - u, v - u \rangle + 2\langle v - u, u - w \rangle + \langle u - w, u - w \rangle \\ &= \langle v - u, v - u \rangle + \langle u - w, u - w \rangle \\ &\geq d(v, u)^2, \end{aligned}$$

with equality only if $w = u = \text{proj}_W(v)$.

7.11 Recall

Let u and v be elements in an inner product space V . The **projection of v on u** is

$$\text{proj}_u v = \frac{\langle u, v \rangle}{\langle u, u \rangle} u$$

Then $\text{proj}_u v$ is the unique nearest element to v of the 1-dimensional subspace spanned by u (in the distance determined by the inner product), and

$$\langle v', u \rangle = 0, \quad v' \perp u, \quad \text{where } v' = v - \text{proj}_u v.$$

So v is the sum of its projection on u and a component orthogonal to u .

In general, if W is any subspace of V and $v \in V$, then there is a unique nearest element in W to v , called $\text{proj}_W v$ or the **projection of v on W** , and $v - \text{proj}_W v$ is orthogonal to **every element of W** .

7.12 Orthogonal Basis

An **orthogonal basis** of an inner product space V is a basis whose elements are all orthogonal to each other.

Theorem 7.2

If V is a finite-dimensional inner product space, then V has an orthogonal (or orthonormal) basis.

Let $\{v_1, \dots, v_n\}$ be any basis. We convert it to an orthogonal basis $\{b_1, \dots, b_n\}$ as follows.

1. Set $b_1 = v_1$.
2. Set $b_2 = v_2 - \text{proj}_{b_1} v_2$. Then $b_2 \perp b_1$ and $\text{span}(b_1, b_2) = \text{span}(v_1, v_2)$.
3. Set $b_3 = v_3 - \text{proj}_{b_1} v_3 - \text{proj}_{b_2} v_3$. Then $b_3 \perp b_1$ and $b_3 \perp b_2$ and $\text{span}(b_1, b_2, b_3) = \text{span}(b_1, b_2, v_3) = \text{span}(v_1, v_2, v_3)$.
4. Continue in this way: Set $b_k = v_k -$ (the sum of its projections on b_1, \dots, b_{k-1}).

The result is an orthogonal basis of V .

7.13 Projection on a subspace

Let W be a subspace of an inner product space V and let v be any element of V .

Then there is a unique element w of W for which $v = w + v'$, and $v' \perp w$ for all $w \in W$.

Then w is called the **orthogonal projection of v on W** , denoted $\text{proj}_W v$.

$$\text{proj}_W v = \text{proj}_{b_1} v + \dots + \text{proj}_{b_k} v \quad \text{for an orthogonal basis } \{b_1, \dots, b_k\} \text{ of } W.$$

Example

In \mathbb{R}^3 , let W be the subspace $x + y + 3z = 0$, and let $v = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$. Then

$$v = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \underbrace{\begin{bmatrix} 6/11 \\ 6/11 \\ -4/11 \end{bmatrix}}_{\text{proj}_W(v)} + \underbrace{\begin{bmatrix} 5/11 \\ 5/11 \\ 15/11 \end{bmatrix}}_{\perp W}$$

7.14 How to calculate $\text{proj}_W(v)$

Example 7.2

In \mathbb{R}^3 , let W be the subspace $x + y + 3z = 0$, and let $v = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$. Find an orthogonal basis of W

For example $\{b_1, b_2\}$ with

$$b_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \quad b_2 = \begin{bmatrix} 3 \\ 3 \\ -2 \end{bmatrix}.$$

Then

$$\begin{aligned} \text{proj}_W v &= \text{proj}_{b_1} v + \text{proj}_{b_2} v = \frac{\langle v, b_1 \rangle}{\langle b_1, b_1 \rangle} b_1 + \frac{\langle v, b_2 \rangle}{\langle b_2, b_2 \rangle} b_2 \\ &= \frac{0}{2} b_1 + \frac{4}{22} b_2 = \frac{4}{22} b_2 = \begin{bmatrix} 6/11 \\ 6/11 \\ -4/11 \end{bmatrix}. \end{aligned}$$

$$v = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \underbrace{\begin{bmatrix} 6/11 \\ 6/11 \\ -4/11 \end{bmatrix}}_{\text{proj}_W(v)} + \underbrace{\begin{bmatrix} 5/11 \\ 5/11 \\ 15/11 \end{bmatrix}}_{\perp W}$$

Example 7.3: I

\mathbb{R}^3 , find the unique point of the plane $W : x + 2y - z = 0$ that is nearest to the point $v : (1, 2, 2)$.

First find an orthogonal basis for W : for example $\{b_1, b_2\}$, where

$$b_1 = (1, 0, 1), \quad b_2 = (1, -1, -1).$$

Then

$$\begin{aligned} \text{proj}_W(v) &= \text{proj}_{b_1}(v) + \text{proj}_{b_2}(v) \\ &= \frac{\langle b_1, v \rangle}{\langle b_1, b_1 \rangle} b_1 + \frac{\langle b_2, v \rangle}{\langle b_2, b_2 \rangle} b_2 \\ &= \frac{3}{2} b_1 - \frac{3}{3} b_2 \\ &= \left(\frac{3}{2}, 0, \frac{3}{2} \right) - (1, -1, -1) = \left(\frac{1}{2}, 1, \frac{5}{2} \right). \end{aligned}$$

7.15 Application: least squares for overdetermined systems

Example Consider the following **overdetermined** linear system.

$$\begin{aligned} 2x + y &= 3 \\ x - y &= 0 \\ x - 3y &= -4 \end{aligned} \quad \Longleftrightarrow \quad \begin{bmatrix} 2 & 1 \\ 1 & -1 \\ 1 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ -4 \end{bmatrix}$$

This system has three equations and only two variables. It is inconsistent and **overdetermined** — each pair of equations has a simultaneous solution, but all three do not.

Overdetermined systems arise quite often in applications, from observed data. Even if they do not have exact solutions, approximate solutions are of interest.

For a vector $b \in \mathbb{R}^3$, the system

$$\underbrace{\begin{bmatrix} 2 & 1 \\ 1 & -1 \\ 1 & -3 \end{bmatrix}}_A \begin{bmatrix} x \\ y \end{bmatrix} = b$$

has a solution if and only if b belongs to the 2-dimensional linear span W of the columns of the coefficient matrix

$$A: v_1 = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \text{ and } v_2 = \begin{bmatrix} 1 \\ -1 \\ -3 \end{bmatrix}.$$

If not, then the nearest element of W to b is $b' = \text{proj}_W(b)$, and our approximate solutions for x and y are the entries of the vector c in \mathbb{R}^2 for which $Ac = b'$. We know that $b' - b$ is orthogonal to v_1 and v_2 , which are the rows of A^T . Hence

$$A^T(b' - b) = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \implies A^T b' = A^T A c = A^T b \implies c = (A^T A)^{-1} A^T b$$

In our example,

$$\begin{bmatrix} 2 & 1 \\ 1 & -1 \\ 1 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ -4 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & 1 \\ 1 & -1 \\ 1 & -3 \end{bmatrix}, \quad A^T = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & -3 \end{bmatrix}, \quad A^T A = \begin{bmatrix} 6 & -2 \\ -2 & 11 \end{bmatrix}, \quad A^T b = \begin{bmatrix} 2 \\ 15 \end{bmatrix}$$

The least squares solution is given by

$$\begin{bmatrix} x \\ y \end{bmatrix} = c = (A^T A)^{-1} A^T b = \frac{1}{62} \begin{bmatrix} 11 & 2 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} 2 \\ 15 \end{bmatrix} = \begin{bmatrix} \frac{26}{31} \\ \frac{47}{31} \end{bmatrix}.$$