

IMPARTIAL ACHIEVEMENT & AVOIDANCE GAMES FOR GENERATING FINITE GROUPS

ACGT Seminar at NAU

Dana C. Ernst

Northern Arizona University

September 26 & October 3, 2017

Joint work with Bret Benesh and Nándor Sieben

Intuitive Definition

Combinatorial Game Theory (CGT) is the study of two-person games satisfying:

- Two players alternate making moves.
- No hidden information.
- No random moves.

Combinatorial games

- Chess
- Go
- Connect Four
- Nim
- Tic-Tac-Toe
- X-Only Tic-Tac-Toe

Combinatorial games

- Chess
- Go
- Connect Four
- Nim
- Tic-Tac-Toe
- X-Only Tic-Tac-Toe

Non-combinatorial games

- Battleship (hidden information)
- Rock-Paper-Scissors (non-alternating and random)
- Poker (hidden information and random)

Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

Partizan

- Chess
- Go
- Connect Four
- Tic-Tac-Toe

Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

Partizan

- Chess
- Go
- Connect Four
- Tic-Tac-Toe

Impartial

- Nim
- X-Only Tic-Tac-Toe

Comments

- We are interested in impartial games.
- We will require that game sequence is finite and there are no ties.
- Player that moves first is called α and second player is called β .
- **Normal Play:** The last player to move wins.
- **Misère Play:** The last player to move loses.

Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Question

Is there an optimal strategy for either player?

Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Question

Is there an optimal strategy for either player?

Answer

Boring: α always wins; just take the whole pile.

Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Question

Is there an optimal strategy for either player?

Answer

Boring: α always wins; just take the whole pile.

Multi-pile Nim

Start with k piles consisting of n_1, \dots, n_k stones, respectively. Each player chooses at least one stone from a single pile. The player that takes the last stone wins. Denoted $*n_1 + \dots + *n_k$.

Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.


Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.




Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.




Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.




Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



Example


Let's play $*1 + *2 + *2$. Here's a possible sequence.



In this case, α wins.

Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



In this case, α wins.

Question

In general, is there an optimal strategy for either player?

Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



In this case, α wins.

Question

In general, is there an optimal strategy for either player?

Answer

Short answer is yes: write sizes of piles in binary, do binary addition without carry (XOR), and if possible, hand your opponent a sum of 0. If players make optimal moves, this is only possible for one of the players.

Definition

An **impartial game** is a finite set X of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

Definition

An **impartial game** is a finite set X of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

- Two players take turns choosing a single available option in $\text{Opt}(Q)$ of current position Q .
- Player who encounters empty option set cannot move and loses.
- **P-position**: previous player (player that just moved) wins
- **N-position**: next player (player that is about to move) wins

Definition

An **impartial game** is a finite set X of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

- Two players take turns choosing a single available option in $\text{Opt}(Q)$ of current position Q .
- Player who encounters empty option set cannot move and loses.
- **P-position**: previous player (player that just moved) wins
- **N-position**: next player (player that is about to move) wins

From perspective of the player that is about to move, a **P-position** is a **losing position** while an **N-position** is a **winning position**.

Examples

- $*n$ is an N-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position
- $*1 + *2$ is an N-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position
- $*1 + *2$ is an N-position
- Empty X-Only Tic-Tac-Toe board is an N-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position
- $*1 + *2$ is an N-position
- Empty X-Only Tic-Tac-Toe board is an N-position

Definition

If G and H are games, then $G + H$ is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

Definition

If G and H are games, then $G + H$ is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

Theorem

$G + G$ is a P-position.

Definition

If G and H are games, then $G + H$ is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

Theorem

$G + G$ is a P-position.

Proof



Copy cat.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

$$\cdot \ast 1 + \ast 1 = \ast 0$$

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

· $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

- $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.
- $*1 + *2 = *3$

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

- $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.
- $*1 + *2 = *3$ since $*1 + *2 + *3$ is a P-position.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

- $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.
- $*1 + *2 = *3$ since $*1 + *2 + *3$ is a P-position.

Theorem

$G_1 = G_2$ if and only if $G_1 + H$ and $G_2 + H$ have the same outcome for all H .

Definition

If A is a finite subset of nonnegative integers, then $\text{mex}(A)$ is the smallest nonnegative integer not in A .

Definition

If A is a finite subset of nonnegative integers, then $\text{mex}(A)$ is the smallest nonnegative integer not in A .

Examples

- $\text{mex}(\{0, 1, 2, 4, 5\}) = 3$
- $\text{mex}(\{1, 3\}) = 0$
- $\text{mex}(\{0, 1\}) = 2$
- $\text{mex}(\emptyset) = 0$

Definition

If G is a game, then

$$\text{nim}(G) := \text{mex}(\{\text{nim}(Q) \mid Q \in \text{Opt}(G)\}).$$

This is a recursive definition. We start computing with terminal positions (empty option set).

Definition

If G is a game, then

$$\text{nim}(G) := \text{mex}(\{\text{nim}(Q) \mid Q \in \text{Opt}(G)\}).$$

This is a recursive definition. We start computing with terminal positions (empty option set).

Examples

- $\text{nim}(*0) = \text{mex}(\emptyset) = 0$
- $\text{nim}(*1) = \text{mex}(\{\text{nim}(*0)\}) = \text{mex}(\{0\}) = 1$
- $\text{nim}(*2) = \text{mex}(\{\text{nim}(*0), \text{nim}(*1)\}) = \text{mex}(\{0, 1\}) = 2$
- $\text{nim}(*n) = n$
- $\text{nim}(*1 + *1) = \text{mex}(\{\text{nim}(*1)\}) = \text{mex}(\{1\}) = 0$
- $\text{nim}(*1 + *2) = \text{mex}(\{\text{nim}(*2), \text{nim}(*1), \text{nim}(*1 + *1)\})$
 $= \text{mex}(\{2, 1, 0\}) = 3$

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

We can think of nim-numbers as “isomorphism” classes of games.

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

We can think of nim-numbers as “isomorphism” classes of games.

Theorem

2nd player β wins G if and only if $G = *0$.

Let G be a finite (possibly trivial) group.

Let G be a finite (possibly trivial) group.

Generate Game

For the achievement game $\text{GEN}(G)$:

Let G be a finite (possibly trivial) group.

Generate Game

For the achievement game $\text{GEN}(G)$:

- 1st player chooses any $g_1 \in G$.
- At k th turn, designated player selects $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$ to create position $\{g_1, \dots, g_k\}$.
- Player wins on the n th turn if $\langle g_1, \dots, g_n \rangle = G$.

Let G be a finite (possibly trivial) group.

Generate Game

For the achievement game $\text{GEN}(G)$:

- 1st player chooses any $g_1 \in G$.
- At k th turn, designated player selects $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$ to create position $\{g_1, \dots, g_k\}$.
- Player wins on the n th turn if $\langle g_1, \dots, g_n \rangle = G$.

Positions of $\text{GEN}(G)$ are subsets of terminal positions, which are certain generating sets of G .

MATCH-UP



Name: LeBron James

Bret Benesh

Height: 6'8"

6'5"

Weight: 260 lbs

180 lbs

Age: 32 years

>32 years

Salary: \$30.96 million/year

\$0 million/year

Accolades: 3x NBA Champion
4x NBA MVP
2x Olympic gold medalist
11x NBA All-Star

Never had a cavity
Sagittarius

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron

P

$\langle P \rangle$

Bret

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	S_3	

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	$(1, 3, 2)$
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	S_3	

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	$(1, 3, 2)$
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	S_3	

Let G be a finite nontrivial group.

Do Not Generate Game

For the **avoidance game** $\text{DNG}(G)$:

- 1st player chooses $g_1 \in G$ such that $\langle g_1 \rangle \neq G$.
- At the k th turn, designated player selects $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$ such that $\langle g_1, \dots, g_k \rangle \neq G$ to create position $\{g_1, \dots, g_k\}$.
- Player that cannot select an element without building a generating set is loser.

Positions of $\text{DNG}(G)$ are exactly the non-generating subsets of G and terminal positions are the maximal subgroups of G .

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron

P

$\langle P \rangle$

Bret

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
e	$\{(1, 2, 3), (1, 3, 2), e\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
e	$\{(1, 2, 3), (1, 3, 2), e\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
e	$\{(1, 2, 3), (1, 3, 2), e\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron

P

$\langle P \rangle$

Bret

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
	$\{r^2\}$	\mathbb{Z}_2	r^2

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
	$\{r^2\}$	\mathbb{Z}_2	r^2
r^3	$\{r^2, r^3\}$	\mathbb{Z}_4	

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
r^3	$\{r^2\}$	\mathbb{Z}_2	r^2
	$\{r^2, r^3\}$	\mathbb{Z}_4	
	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
r^3	$\{r^2\}$	\mathbb{Z}_2	r^2
	$\{r^2, r^3\}$	\mathbb{Z}_4	
r	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e
	$\{r^2, r^2, e, r\}$	\mathbb{Z}_4	

LEBRON VS BRET: GAME THREE


DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
	$\{r^2\}$	\mathbb{Z}_2	r^2
r^3	$\{r^2, r^3\}$	\mathbb{Z}_4	
	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e
r	$\{r^2, r^2, e, r\}$	\mathbb{Z}_4	

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
r^3	$\{r^2\}$	\mathbb{Z}_2	r^2
	$\{r^2, r^3\}$	\mathbb{Z}_4	
r	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e
	$\{r^2, r^2, e, r\}$	\mathbb{Z}_4	

- 1987: Harary and Anderson determine outcomes for abelian groups.

SOME HISTORY

- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.


SOME HISTORY

- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.
- 2014: Ernst and Sieben determine nim-numbers (and hence outcomes) for cyclic, dihedral, abelian.

SOME HISTORY


- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.
- 2014: Ernst and Sieben determine nim-numbers (and hence outcomes) for cyclic, dihedral, abelian.
- 2016: Benesh, Ernst, and Sieben establish **subgroup-based criteria** for the determination of nim-numbers (and hence outcomes) for DNG, characterize **spectrum of nim-numbers for DNG**, determine nim-numbers for GEN and DNG for a variety of groups including generalized dihedral, symmetric, and alternating groups.

REPRESENTATIVE GAME TREES




Representative game tree for $\text{GEN}(S_3) = *3$

STRUCTURE DIAGRAMS






$\text{DNG}(\mathbb{Z}_6)$



Structure diagram

SIMPLIFIED STRUCTURE DIAGRAMS




$DNG(\mathbb{Z}_6 \times \mathbb{Z}_3)$


SIMPLIFIED STRUCTURE DIAGRAMS




(a) $\text{DNG}(\mathbb{D}_{4k})$




(b) $\text{DNG}(\mathbb{D}_{2k+1})$




(c) $\text{DNG}(\mathbb{D}_{4k+2})$



(d) $\text{GEN}(\mathbb{D}_{4k})$



(e) $\text{GEN}(\mathbb{D}_{2k+1})$



(f) $\text{GEN}(\mathbb{D}_{4k+2})$

Simplified structure diagrams for dihedral groups

Theorem (Ernst, Sieben)

If $n \geq 2$, then $\text{nim}(\text{GEN}(\mathbb{Z}_n)) = \text{nim}(\text{DNG}(\mathbb{Z}_n)) + 1$.

Theorem (Ernst, Sieben)

If $n \geq 2$, then $\text{nim}(\text{GEN}(\mathbb{Z}_n)) = \text{nim}(\text{DNG}(\mathbb{Z}_n)) + 1$.

Theorem (Ernst, Sieben)

If $n \geq 2$, then

$$\text{DNG}(\mathbb{Z}_n) = \begin{cases} *1, & n = 2 \\ *1, & n \equiv_2 1 \\ *0, & n \equiv_4 0 \\ *3, & n \equiv_4 2 \end{cases}$$

and

$$\text{GEN}(\mathbb{Z}_n) = \begin{cases} *2, & n = 2 \\ *2, & n \equiv_2 1 \\ *1, & n \equiv_4 0 \\ *4, & n \equiv_4 2 \end{cases}$$

Theorem (Ernst, Sieben)

For $n \geq 3$, we have

$$\text{DNG}(\mathbb{D}_n) = \begin{cases} *3, & n \equiv_2 1 \\ *0, & n \equiv_2 0 \end{cases}$$

and

$$\text{GEN}(\mathbb{D}_n) = \begin{cases} *3, & n \equiv_2 1 \\ *0, & n \equiv_4 0 \\ *1, & n \equiv_4 2 \end{cases}$$

Theorem (Ernst, Sieben)

If G is a finite nontrivial abelian group, then

$$\text{DNG}(G) = \begin{cases} *1, & G \text{ is nontrivial of odd order} \\ *1, & G = \mathbb{Z}_2 \\ *3, & G = \mathbb{Z}_2 \times \mathbb{Z}_{2k+1} \text{ with } k \geq 1 \\ *0, & \text{else} \end{cases}$$

$$\text{GEN}(G) = \begin{cases} *2, & |G| \text{ is odd and } d(G) \leq 2 \\ *1, & |G| \text{ is odd and } d(G) \geq 3 \\ *2, & G = \mathbb{Z}_2 \\ *1, & G = \mathbb{Z}_{4k} \text{ with } k \geq 1 \\ *4, & G = \mathbb{Z}_{4k+2} \text{ with } k \geq 1 \\ *1, & G = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_m \times \mathbb{Z}_k \text{ for } m, k \text{ odd} \\ *0, & \text{else} \end{cases}$$

Theorem (Ernst, Sieben)

- If G is any finite nontrivial group, then $\text{DNG}(G)$ is $*0$, $*1$, or $*3$.

Theorem (Ernst, Sieben)

- If G is any finite nontrivial group, then $\text{DNG}(G)$ is $*0$, $*1$, or $*3$.
- If $|G|$ is odd and nontrivial, then $\text{GEN}(G)$ is $*1$ or $*2$.

Theorem (Ernst, Sieben)

- If G is any finite nontrivial group, then $\text{DNG}(G)$ is $*0$, $*1$, or $*3$.
- If $|G|$ is odd and nontrivial, then $\text{GEN}(G)$ is $*1$ or $*2$.

Conjecture (In Progress)

If $|G|$ is even, then $\text{GEN}(G)$ is one of $*0$, $*1$, $*2$, $*3$, $*4$.

Theorem (Benesh, Ernst, Sieben)

Let G be a finite nontrivial group.

- If all maximal subgroups are even, then $\text{DNG}(G) = *0$.
- If all maximal subgroups are odd, then $\text{DNG}(G) = *1$.
- If mixed maximal subgroups, then
 - If the even maximals cover G , then $\text{DNG}(G) = *0$.
 - If the even maximals do not cover G , then $\text{DNG}(G) = *3$.

Theorem (Benesh, Ernst, Sieben)

Let G be a finite nontrivial group.

- If all maximal subgroups are even, then $\text{DNG}(G) = *0$.
- If all maximal subgroups are odd, then $\text{DNG}(G) = *1$.
- If mixed maximal subgroups, then
 - If the even maximals cover G , then $\text{DNG}(G) = *0$.
 - If the even maximals do not cover G , then $\text{DNG}(G) = *3$.

Using our “checklist” criteria, we have completely characterized DNG for nilpotent, generalized dihedral, generalized quaternion, symmetric, Coxeter, alternating, and some Rubik’s cube groups.

Big Picture for DNG

- The players just race to fill up one maximal subgroup M .
- **The beginning of the game is a struggle to determine M .**
- α wants $|M|$ to be odd.
- β wants $|M|$ to be even.

Big Picture for DNG

- The players just race to fill up one maximal subgroup M .
- **The beginning of the game is a struggle to determine M .**
- α wants $|M|$ to be odd.
- β wants $|M|$ to be even.

Strategy

- α wants to pick an element not in any maximal subgroups of even order.
- β wants to pick an involution.

What's left to work on?

- Wrap up spectrum of GEN?
- Wrap up characterization of GEN for nilpotent groups?
- Are there nice results for products and quotients?
- Is it possible to characterize the nim-numbers of GEN in terms of covering conditions by maximal subgroups similar to what we did for DNG?
- What about other “closure systems”? We are currently tinkering with convex hulls of finitely many points in the plane.

Thanks!