

Course Notes for MAT 612: Abstract Algebra II

Dana C. Ernst, PhD
Northern Arizona University

Spring 2016

© 2016 Dana C. Ernst. Some Rights Reserved.

Contents

1	Ring Theory	3
1.1	Definitions and Examples	3
1.2	Ideals and Quotient Rings	7
1.3	Maximal and Prime Ideals	10
1.4	Rings of Fractions	12
1.5	Principal Ideal Domains	14
1.6	Euclidean Domains	16

Chapter 1

Ring Theory

1.1 Definitions and Examples

This section of notes roughly follows Sections 7.1–7.3 in Dummit and Foote.

Recall that a group is a set together with a single binary operation, which together satisfy a few modest properties. Loosely speaking, a ring is a set together with two binary operations (called addition and multiplication) that are related via a distributive property.

Definition 1.1. A **ring** R is a set together with two binary operations $+$ and \times (called **addition** and **multiplication**, respectively) satisfying the following:

- (i) $(R, +)$ is an abelian group.
- (ii) \times is associative: $(a \times b) \times c = a \times (b \times c)$ for all $a, b, c \in R$.
- (iii) The **distributive property** holds: $a \times (b + c) = (a \times b) + (a \times c)$ and $(a + b) \times c = (a \times c) + (b \times c)$ for all $a, b, c \in R$.

Note 1.2. We make a couple comments about notation.

- (1) We write ab in place $a \times b$.
- (2) The additive inverse of the ring element $a \in R$ is denoted $-a$.

Theorem 1.3. Let R be a ring. Then for all $a, b \in R$:

- (1) $0a = a0 = 0$
- (2) $(-a)b = a(-b) = -(ab)$
- (3) $(-a)(-b) = ab$

Definition 1.4. A ring R is called **commutative** if multiplication is commutative.

Definition 1.5. A ring R is said to have an **identity** (or called a **ring with 1**) if there is an element $1 \in R$ such that $1 \times a = a \times 1 = a$ for all $a \in R$.

Theorem 1.6. If R is a ring with 1, then the multiplicative identity is unique and $-a = (-1)a$.

Question 1.7. Requiring $(R, +)$ to be a group is fairly natural, but why require $(R, +)$ to be abelian? Here's one reason. Suppose R has a 1. Compute $(1 + 1)(a + b)$ in two different ways.

Definition 1.8. A ring R with 1 (with $1 \neq 0$) is called a **division ring** if every nonzero element in R has a multiplicative inverse: if $a \in R \setminus \{0\}$, then there exists $b \in R$ such that $ab = ba = 1$.

Definition 1.9. A commutative division ring is called a **field**.

Definition 1.10. A nonzero element a in a ring R is called a **zero divisor** if there is a nonzero element $b \in R$ such that either $ab = 0$ or $ba = 0$.

Theorem 1.11 (Cancellation Law). Assume $a, b, c \in R$ such that a is not a zero divisor. If $ab = ac$, then either $a = 0$ or $b = c$.

Definition 1.12. Assume R is a ring with 1 with $1 \neq 0$. An element $u \in R$ is called a **unit** in R if u has a multiplicative inverse (i.e., there exists $v \in R$ such that $uv = vu = 1$). The set of units in R is denoted R^\times .

Theorem 1.13. If $R^\times \neq \emptyset$, then R^\times forms a group under multiplication.

Note 1.14. We make a few observations.

- (1) A field is a commutative ring F with identity $1 \neq 0$ in which every nonzero element is a unit, i.e., $F^\times = F \setminus \{0\}$.
- (2) Zero divisors can never be units.
- (3) Fields never have zero divisors.

Definition 1.15. A commutative ring with identity $1 \neq 0$ is called an **integral domain** if it has no zero divisors.

Note 1.16. The Cancellation Law (Theorem 1.11) holds in integral domains for any three elements.

Corollary 1.17. Any finite integral domain is a field.

Proof. For any nonzero $a \in R$, define $f_a : R \rightarrow R$ via $f_a(x) = ax$. If R is an integral domain, the Cancellation Law forces f_a to be injective. If R is finite, then f_a is also surjective. In this case, there exists $b \in R$ such that $ab = 1$. □

Example 1.18. Here are some examples of rings. Details left as an exercise.

- (1) **Zero Ring:** If $R = \{0\}$, we can turn R into a ring in the obvious way. The zero ring is a finite commutative ring with 1. It is the only ring where the additive and multiplicative identities are equal. The zero ring is not a division ring, not a field, and not an integral domain.

- (2) **Trivial Ring:** Given any abelian group R , we can turn R into a ring by defining multiplication via $ab = 0$ for all $a, b \in R$. Trivial rings are commutative rings in which every nonzero element is a zero divisor. Hence a trivial ring is not a division ring, not a field, and not an integral domain.
- (3) The integers \mathbb{Z} form a ring under the usual operations of addition and multiplication. The integers form an integral domain, but \mathbb{Z} is not a division ring, and hence not a field.
- (4) The rational numbers \mathbb{Q} , the real numbers \mathbb{R} , and the complex numbers \mathbb{C} are fields under the usual operations of addition and multiplication.
- (5) For $n \geq 1$, the set \mathbb{Z}_n is a commutative ring with 1 under the operations of addition and multiplication mod n . The group of units \mathbb{Z}_n^\times is the set of elements in \mathbb{Z}_n that are relatively prime to n . All other nonzero elements are zero divisors. It turns out that \mathbb{Z}_n forms a finite field iff n is prime.
- (6) The set of even integers $2\mathbb{Z}$ forms a commutative ring under the usual operations of addition and multiplication. However, $2\mathbb{Z}$ does not have a 1, and hence cannot be a division ring nor a field nor an integral domain.
- (7) **Polynomial Ring:** Fix a commutative ring R . Let $R[x]$ denote the set of polynomials in the variable x with coefficients in R . Then $R[x]$ is a commutative ring with 1. The units of $R[x]$ are exactly the units of R (if there are any). So, $R[x]$ is never a division ring nor a field. However, if R is an integral domain, then so is $R[x]$.
- (8) **Matrix Ring:** Fix a ring R and let n be a positive integer. Let $M_n(R)$ be the set of $n \times n$ matrices with entries from R . Then $M_n(R)$ forms a ring under ordinary matrix addition and multiplication. If R is nontrivial and $n \geq 2$, then $M_n(R)$ always has zero divisors and $M_n(R)$ is not commutative even if R is. If R has a 1, then the matrix with 1's down the diagonal and 0's elsewhere is the multiplicative identity in $M_n(R)$. In this case, the group of units is the set of invertible $n \times n$ matrices, denoted $GL_n(R)$ and called the **general linear group of degree n over R** .
- (9) **Quadratic Field:** Define $\mathbb{Q}(\sqrt{2}) = \{a + b\sqrt{2} \mid a, b \in \mathbb{Q}\}$. It turns out that $\mathbb{Q}(\sqrt{2})$ is a field. In fact, we can replace 2 with any rational number that is not a perfect square in \mathbb{Q} .
- (10) **Hamilton Quaternions:** Define $\mathbb{H} = \{a + bi + cj + dk \mid a, b, c, d \in \mathbb{R}, i, j, k \in Q_8\}$. Then \mathbb{H} forms a ring, where addition is definite componentwise in i, j , and k and multiplication is defined by expanding products and the simplifying using the relations of Q_8 . It turns out that \mathbb{H} is a non-commutative ring with 1.

Definition 1.19. A **subring** of a ring R is a subgroup of R that is closed under multiplication.

Note 1.20. The property “is a subring” is clearly transitive. To show that a subset S of a ring R is a subring, it suffices to show that $S \neq \emptyset$, S is closed under subtraction, and S is closed under multiplication.

Example 1.21. Here are a few quick examples.

- (1) \mathbb{Z} is a subring of \mathbb{Q} , which is a subring of \mathbb{R} , which in turn is a subring of \mathbb{C} .
- (2) $2\mathbb{Z}$ is a subring of \mathbb{Z} .
- (3) The set $\mathbb{Z}(\sqrt{2}) = \{a + b\sqrt{2} \mid a, b \in \mathbb{Z}\}$ is a subring of $\mathbb{Q}(\sqrt{2})$.
- (4) The ring R is a subring of $R[x]$ if we identify R with set of constant functions.
- (5) The set of polynomials with zero constant term in $R[x]$ is a subring of $R[x]$.
- (6) $\mathbb{Z}[x]$ is a subring of $\mathbb{Q}[x]$.
- (7) \mathbb{Z}_n is *not* a subring of \mathbb{Z} .

Definition 1.22. Let R and S be rings. A **ring homomorphism** is a map $\phi : R \rightarrow S$ satisfying

- (i) $\phi(a + b) = \phi(a) + \phi(b)$
- (ii) $\phi(ab) = \phi(a)\phi(b)$

for all $a, b \in R$. The **kernel** of ϕ is defined via $\ker(\phi) = \{a \in R \mid \phi(a) = 0\}$. If ϕ is a bijection, then ϕ is called an **isomorphism**, in which case, we say that R and S are **isomorphic rings** and write $R \cong S$.

Example 1.23.

- (1) For $n \in \mathbb{Z}$, define $\phi_n : \mathbb{Z} \rightarrow \mathbb{Z}$ via $\phi_n(x) = nx$. We see that $\phi_n(x + y) = n(x + y) = nx + ny = \phi_n(x) + \phi_n(y)$. However, $\phi_n(xy) = n(xy)$ while $\phi_n(x)\phi_n(y) = (nx)(ny) = n^2xy$. It follows that ϕ_n is a ring homomorphism exactly when $n \in \{0, 1\}$.
- (2) Define $\phi : \mathbb{Q}[x] \rightarrow \mathbb{Q}$ via $\phi(p(x)) = p(0)$ (called **evaluation at 0**). It turns out that ϕ is a ring homomorphism, where $\ker(\phi)$ is the set of polynomials with 0 constant term.

Theorem 1.24. Let $\phi : R \rightarrow S$ be a ring homomorphism.

- (1) $\phi(R)$ is a subring of S .
- (2) $\ker(\phi)$ is a subring of R .

In fact, we can say something even stronger about the kernel of a ring homomorphism, which will lead us to the notion of an **ideal**.

Theorem 1.25. Let $\phi : R \rightarrow S$ be a ring homomorphism. If $\alpha \in \ker(\phi)$ and $r \in R$, then $\alpha r, r\alpha \in \ker(\phi)$. That is, $\ker(\phi)$ is closed under multiplication by elements of R .

1.2 Ideals and Quotient Rings

This section of notes roughly follows Sections 7.3–7.4 in Dummit and Foote.

Recall that in the case of a homomorphism ϕ of groups, the fibers of ϕ have the structure of a group (that happens to be isomorphic to the image of ϕ by the First Isomorphism Theorem). In this case, the kernel of ϕ is the identity of the associated quotient group. This naturally led to the notion of a normal subgroup (i.e., those groups that correspond to kernels of homomorphisms). Can we do the same sort of thing for rings?

Let $\phi : R \rightarrow S$ be a ring homomorphism with $\ker(\phi) = I$. Note that ϕ is also a group homomorphism of abelian groups and the fibers of ϕ are the cosets $r + I$. That is, if $\phi(r) = a$, then the fiber of ϕ over a is the coset $\phi^{-1}(a) = r + I$.

These cosets naturally have the structure of a ring isomorphic to the image of ϕ :

$$(r + I) + (s + I) = (r + s) + I \quad (1.1)$$

$$(r + I)(s + I) = (rs) + I \quad (1.2)$$

The reason for this is that if the fiber of $a \in S$ is $\phi^{-1}(a) = X$ and the fiber of $b \in S$ is $\phi^{-1}(b) = Y$, then the fibers of $a + b$ and ab are $X + Y$ and XY , respectively.

The corresponding ring of cosets is called the **quotient ring** of R by $I = \ker(\phi)$ and is denoted by R/I . The additive structure of the quotient ring R/I is exactly the additive quotient group of the additive abelian group R by the normal subgroup I (all subgroups are normal in abelian groups). When I is the kernel of some ring homomorphism ϕ , the additive abelian quotient group R/I also has a multiplicative structure defined in (2) above, making R/I into a ring.

Question 1.26. Can we make R/I into a ring for any subring I ?

The answer is “no” in general, just like in the situation with groups. But perhaps this isn’t obvious because if I is an arbitrary subring of R , then I is necessarily an additive subgroup of the abelian group R , which implies that I is an additive normal subgroup of the group R . It turns out that the multiplicative structure of R/I may not be well-defined if I is an arbitrary subring.

Let I be an arbitrary *subgroup* of the additive subgroup R . Let $r + I$ and $s + I$ be two arbitrary cosets. In order for multiplication of the cosets to be well-defined, the product of the two cosets must be independent of choice of representatives. Let $r + \alpha$ and $s + \beta$ be arbitrary representatives of $r + I$ and $s + I$, respectively ($\alpha, \beta \in I$), so that $r + I = (r + \alpha) + I$ and $s + I = (s + \beta) + I$. We must have

$$(r + \alpha)(s + \beta) + I = rs + I. \quad (1.3)$$

This needs to be true for all possible choices of $r, s \in R$ and $\alpha, \beta \in I$. In particular, it must be true when $r = s = 0$. In this case, we must have

$$\alpha\beta + I = I. \quad (1.4)$$

But this only happens when $\alpha\beta \in I$. That is, one requirement for multiplication of cosets to be well-defined is that I must be closed under multiplication, making I a *subring*.

Next, if we let $s = 0$ and let r be arbitrary, we see that we must have $r\beta \in I$ for every $r \in R$ and every $\beta \in I$. That is, it must be the case that I is closed under multiplication on the left by elements from R . Similarly, letting $r = 0$, we can conclude that we must have I closed under multiplication on the right by elements from R .

On the other hand, if I is closed under multiplication on the left and on the right by elements from R , then it is clear that relation (4) above is satisfied.

It is easy to verify that if the multiplication of cosets defined in (2) above is well-defined, then this multiplication makes the additive quotient group R/I into a ring (just check the axioms for being a ring).

We have shown that the quotient R/I of the ring R by a subgroup I has a natural ring structure iff I is closed under multiplication on the left and right by elements of R (which also forces I be a subring). Such subrings are called **ideals**.

Definition 1.27. Let R be a ring and let I be a subset of R .

- (1) I is a **left ideal** (respectively, **right ideal**) of R iff I is a subring and $rI \subseteq I$ (respectively, $Ir \subseteq I$) for all $r \in R$.
- (2) I is an **ideal** (or **two-sided ideal**) iff I is both a left and a right ideal.

Here's a summary of everything that just happened.

Theorem 1.28. Let R be a ring and let I be an ideal of R . Then the additive quotient group R/I is a ring under the binary operations:

$$(r + I) + (s + I) = (r + s) + I \tag{1.5}$$

$$(r + I)(s + I) = (rs) + I \tag{1.6}$$

for all $r, s \in R$. Conversely, if I is any subgroup such that the above operations are well-defined, then I is an ideal of R .

As you might expect, we have some isomorphism theorems.

Theorem 1.29 (First Isomorphism Theorem for Rings). If $\phi : R \rightarrow S$ is a ring homomorphism, then $\ker(\phi)$ is an ideal of R and $R/\ker(\phi) \cong \phi(R)$.

If I and J are ideals of R , then it is easy to verify that $I \cap J$, $I + J = \{a + b \mid a \in I, b \in J\}$, and $IJ = \{\text{finite sums of elements of the form } ab \text{ for } a \in I, b \in J\}$ are also ideals of R . We also have the expected Second, Third, and Fourth Isomorphism Theorems for rings.

The next theorem tells us that a subring is an ideal iff it is a kernel of a ring homomorphism.

Theorem 1.30. If I is any ideal of R , then the **natural projection** $\pi : R \rightarrow R/I$ defined via $\pi(r) = r + I$ is a surjective ring homomorphism with $\ker(\pi) = I$.

For the remainder of this section, assume that R is a ring with identity $1 \neq 0$.

Definition 1.31. Let A be any subset of R .

- (1) Let (A) denote the smallest ideal of R containing A , called the **ideal generated by A** . If A consists of a single element, say $A = \{a\}$, then $(a) := (\{a\})$ is called a **principal ideal**.
- (2) $RA := \{r_1 a_1 + \cdots + r_n a_n \mid r_i \in R, a_i \in A, n \in \mathbb{Z}^+\}$, $AR := \{a_1 r_1 + \cdots + a_n r_n \mid a_i \in A, r_i \in R, n \in \mathbb{Z}^+\}$, and $RAR := \{r_1 a_1 r'_1 + \cdots + r_n a_n r'_n \mid r_i, r'_i \in R, a_i \in A, n \in \mathbb{Z}^+\}$.

Note 1.32. The following facts are easily verified.

- (1) (A) is the intersection of all ideals containing A .
- (2) RA , AR , and RAR are the left, right, and two-sided ideals generated by A .
- (3) If R is commutative, then $RA = AR = RAR = (A)$.
- (4) If R is commutative, then $(a) = Ra = aR$.

Example 1.33. Here are a couple examples. The details are left as exercises.

- (1) In \mathbb{Z} , $n\mathbb{Z} = (n) = (-n)$. In fact, these are the only ideals in \mathbb{Z} (since these are the only subgroups). So, all the ideals in \mathbb{Z} are principal. If m and n are positive integers, then $n\mathbb{Z} \subseteq m\mathbb{Z}$ iff m divides n . Moreover, we have $(m, n) = (d)$, where d is the greatest common divisor of m and n .
- (2) Consider the ideal $(2, x)$ in $\mathbb{Z}[x]$. Note that $(2, x) = \{2p(x) + xq(x) \mid p(x), q(x) \in \mathbb{Z}[x]\}$. Then $(2, x)$ is the collection of polynomials from $\mathbb{Z}[x]$ that have even constant term. In particular, $2, x \in (2, x)$. However, there is no single polynomial in $\mathbb{Z}[x]$ that we can use to generate both 2 and x that only produces polynomials with even constant terms.

Theorem 1.34. Let I be an ideal of R .

- (1) $I = R$ iff I contains a unit.
- (2) Assume R is commutative. Then R is a field iff its only ideals are 0 and R .

Loosely speaking, the previous result says that fields are “like simple groups.”

Corollary 1.35. If R is a field, then every nonzero ring homomorphism from R into another ring is an injection.

1.3 Maximal and Prime Ideals

This section of notes roughly follows Section 7.4 in Dummit and Foote.

In this section of notes, we will study two important classes of ideals, namely **maximal** and **prime** ideals, and study the relationship between them.

Definition 1.36. An ideal M in a ring R is called a **maximal ideal** if $M \neq R$ and the only ideals containing M are M and R .

Example 1.37. Here are a few examples. Checking the details is left as an exercise.

- (1) In \mathbb{Z} , all the ideals are of the form $n\mathbb{Z}$ for $n \in \mathbb{Z}^+$. The maximal ideals correspond to the ideals $p\mathbb{Z}$, where p is prime.
- (2) Consider the integral domain $\mathbb{Z}[x]$. The ideals (x) (i.e., the subring containing polynomials with 0 constant term) and (2) (i.e., the set of polynomials with even coefficients) are not maximal since both are contained in the proper ideal $(2, x)$. However, as we shall see soon, $(2, x)$ is maximal in $\mathbb{Z}[x]$.
- (3) The zero ring has no maximal ideals.
- (4) Consider the abelian group \mathbb{Q} under addition. We can turn \mathbb{Q} into a trivial ring by defining $ab = 0$ for all $a, b \in \mathbb{Q}$. In this case, the ideals are exactly the additive subgroups of \mathbb{Q} . However, \mathbb{Q} has no maximal subgroups, and so \mathbb{Q} has no maximal ideals.

The next result states that rings with an identity $1 \neq 0$ always have maximal ideals. It turns out that we won't need this result going forward, so we'll skip its proof. However, it is worth noting that all known proofs make use of Zorn's Lemma (equivalent to the Axiom of Choice), which is also true for the proofs that a finitely generated group has maximal subgroups or that every vector space has a basis.

Theorem 1.38. In a ring with 1, every proper ideal is contained in a maximal ideal.

For commutative rings, there is a very nice characterization about maximal ideals in terms of the structure of their quotient rings.

Theorem 1.39 (Student Presentation 9). Assume R is commutative. The ideal M is maximal iff R/M is a field.

Example 1.40. We can use the previous theorem to verify whether an ideal is maximal.

- (1) Recall that $\mathbb{Z}/n\mathbb{Z} \cong \mathbb{Z}_n$ and that \mathbb{Z}_n is a field iff n is prime. We can conclude that $n\mathbb{Z}$ is a maximal ideal precisely when n is prime.
- (2) Define $\phi : \mathbb{Z}[x] \rightarrow \mathbb{Z}$ via $\phi(p(x)) = p(0)$. Then ϕ is surjective and $\ker(\phi) = (x)$. By the First Isomorphism Theorem for Rings, we see that $\mathbb{Z}[x]/(x) \cong \mathbb{Z}$. However, \mathbb{Z} is not a field. Hence (x) is not maximal in $\mathbb{Z}[x]$. Now, define $\psi : \mathbb{Z} \rightarrow \mathbb{Z}_2$ via $\psi(x) = x \bmod 2$ and consider the composite homomorphism $\psi \circ \phi : \mathbb{Z}[x] \rightarrow \mathbb{Z}_2$. It is clear that $\psi \circ \phi$ is onto and the kernel of $\psi \circ \phi$ is given by $\{p(x) \in \mathbb{Z}[x] \mid p(0) \in 2\mathbb{Z}\} = (2, x)$. Again by the First Isomorphism Theorem for Rings, $\mathbb{Z}[x]/(2, x) \cong \mathbb{Z}_2$. Since \mathbb{Z}_2 is a field, $(2, x)$ is a maximal ideal.

Definition 1.41. Assume R is commutative. An ideal P is called a **prime ideal** if $P \neq R$ and whenever the product $ab \in P$ for $a, b \in R$, then at least one of a or b is in P .

Example 1.42. In any integral domain, the 0 ideal (0) is a prime ideal. What if the ring is not an integral domain?

Note 1.43. The notion of a prime ideal is a generalization of “prime” in \mathbb{Z} . Suppose $n \in \mathbb{Z}^+ \setminus \{1\}$ such that n divides ab . In this case, n is guaranteed to divide either a or b exactly when n is prime. Now, let $n\mathbb{Z}$ be a proper ideal in \mathbb{Z} with $n > 1$ and suppose $ab \in n\mathbb{Z}$ for $a, b \in \mathbb{Z}$. In order for $n\mathbb{Z}$ to be a prime ideal, it must be true that n divides either a or b . However, this is only guaranteed to be true for all $a, b \in \mathbb{Z}$ when p is prime. That is, the nonzero prime ideals of \mathbb{Z} are of the form $p\mathbb{Z}$, where p is prime. Note that in the case of the integers, the maximal and nonzero prime ideals are the same.

Theorem 1.44. Assume R is a commutative ring. Then the ideal P is a prime ideal in R iff the quotient ring R/P is an integral domain.

Corollary 1.45. Assume R is a commutative ring. Every maximal ideal of R is a prime ideal.

Example 1.46. Recall that $\mathbb{Z}[x]/(x) \cong \mathbb{Z}$. Since \mathbb{Z} is an integral domain, it must be the case that (x) is a prime ideal in $\mathbb{Z}[x]$. However, as we saw in an earlier example, (x) is not maximal in $\mathbb{Z}[x]$ since \mathbb{Z} is not a field. This shows that the converse of the previous corollary is not true.

1.4 Rings of Fractions

This section of notes roughly follows Section 7.5 in Dummit and Foote.

Throughout this whole section, we assume that R is a commutative ring.

Note 1.47. We recall a few relevant facts.

- (1) Theorem 11 (Cancellation Law) tells us that if $ab = ac$ and a is neither 0 nor a zero divisor, then $b = c$.
- (2) Zero divisors are never units.

One upshot of the above is that ring elements that are not zero divisors possess some of the behavior of units. The goal of this section is to prove that every commutative ring R is always a subring of a larger ring Q in which every nonzero element of R that is not a zero divisor is a unit in Q . In particular, we can apply this to integral domains, in which case Q will be a field. This generalizes the construction of \mathbb{Q} from \mathbb{Z} .

Note 1.48. Recall that in \mathbb{Q} , the fraction $\frac{a}{b}$ is the equivalence class of order pairs (a, b) of integers with $b \neq 0$ under the equivalence relation:

$$(a, b) \sim (c, d) \text{ iff } \frac{a}{b} = \frac{c}{d} \text{ iff } ad = bc.$$

Also, every nonzero rational number $\frac{a}{b}$ has multiplicative inverse $\frac{b}{a}$. That is, every nonzero rational number is a unit, making \mathbb{Q} a field. The integers \mathbb{Z} are a subring of \mathbb{Q} . But \mathbb{Z} is an integral domain, not a field.

Theorem 1.49. Let R be a commutative ring. Let D be any nonempty subset of R that does not contain 0, does not contain any zero divisors, and is closed under multiplication. Then there exists a commutative ring Q with 1 such that Q contains R as a subring and every element of D is a unit in Q .

Theorem 1.50. Let R , D , and Q be as in Theorem 1.49. Then every element of Q is of the form rd^{-1} for some $r \in R$ and $d \in D$. In particular, if $D = R \setminus \{0\}$, then Q is a field.

Theorem 1.51. Let R , D , and Q be as in Theorem 1.49. Then Q is the smallest ring containing R in which all elements of D become units, in the following sense. Let S be any commutative ring with 1 and let $\phi : R \rightarrow S$ be any injective ring homomorphism such that $\phi(d)$ is a unit in S for every $d \in D$. Then there is an injective homomorphism $\Phi : Q \rightarrow S$ such that $\Phi|_R = \phi$.

Definition 1.52. Let R , D , and Q be as in Theorem 1.49.

- (1) The ring Q is called the **ring of fractions** of D with respect to R and is denoted $D^{-1}R$.
- (2) If R is an integral domain and $D = R \setminus \{0\}$, then Q is called the **field of fractions** (or **quotient field**) of R .

Corollary 1.53. Let R be an integral domain and let Q be the field of fractions of R . If a field F contains a subring R' isomorphic to R , then the subfield of F generated by R' (i.e., the intersection of all the subfields of F containing R') is isomorphic to Q .

Example 1.54. Here are a few quick examples.

- (1) If R is a field, then its field of fractions is R itself.
- (2) The field of fractions of \mathbb{Z} is \mathbb{Q} . The field of fractions of $2\mathbb{Z}$ is also \mathbb{Q} .
- (3) Consider the polynomial ring $\mathbb{Z}[x]$. Since \mathbb{Z} is an integral domain, so is $\mathbb{Z}[x]$. Then the field of fractions of $\mathbb{Z}[x]$ is the set of rational functions (i.e., functions of the form $p(x)/q(x)$, where $p(x)$ and $q(x)$ are polynomials with integer coefficients and $q(x)$ is not the zero polynomial). Notice that this field contains the field of fractions of \mathbb{Z} , namely \mathbb{Q} . However, it is interesting to point out that the field of fractions of $\mathbb{Q}[x]$ is the same as the field of fractions of $\mathbb{Z}[x]$.

1.5 Principal Ideal Domains

This section of notes roughly follows Sections 8.1-8.2 in Dummit and Foote. Throughout this whole section, we assume that R is a commutative ring.

Definition 1.55. Let R be a commutative ring and let $a, b \in R$ with $b \neq 0$.

- (1) a is said to be **multiple** of b if there exists an element $x \in R$ with $a = bx$. In this case, b is said to **divide** a or be a **divisor** of a , written $b \mid a$.
- (2) A **greatest common divisor** of a and b is a nonzero element d such that
 - (a) $d \mid a$ and $d \mid b$, and
 - (b) if $d' \mid a$ and $d' \mid b$, then $d' \mid d$.

A greatest common divisor of a and b will be denoted $\gcd(a, b)$ (or possibly (a, b)).

Note 1.56. Note that $b \mid a$ in a ring R iff $a \in (b)$ iff $(a) \subseteq (b)$. In particular, if d is any divisor of both a and b , then (d) must contain both a and b , and hence must contain (a, b) . Moreover, if $d = \gcd(a, b)$ iff $(a, b) \subseteq (d)$ and if (d') is any principal ideal containing (a, b) , then $(d) \subseteq (d')$.

The note above immediately proves the following result.

Theorem 1.57. If a and b are nonzero elements in the commutative ring R such that $(a, b) = (d)$, then $d = \gcd(a, b)$.

Note 1.58. It is important to point out that the theorem above is giving us a sufficient condition, but it is not necessary. For example, $(2, x)$ is a maximal ideal in $\mathbb{Z}[x]$ that is not principal. Then $\mathbb{Z}[x] = (1)$ is the unique principal ideal containing both 2 and x , and so $\gcd(2, x) = 1$.

Theorem 1.59. Let R be an integral domain. If $(d) = (d')$, then $d' = ud$ for some unit $u \in R$. In particular, if $d = \gcd(a, b) = d'$, then $d' = ud$ for some unit $u \in R$.

Proof. Easy exercise. □

Definition 1.60. A **principal ideal domain** (PID) is an integral domain in which every ideal is principal.

Example 1.61. Here are some short examples.

- (1) \mathbb{Z} is a PID.
- (2) $\mathbb{Z}[x]$ is not a PID since $(2, x)$ is not principal.

Theorem 1.62. Let R be a PID, $a, b \in R \setminus \{0\}$, and $(d) = (a, b)$. Then

- (1) $d = \gcd(a, b)$
- (2) $d = ax + by$ for some $x, y \in R$

(3) d is unique up to multiplication by a unit of R .

Proof. The result follows from Theorems 1.57 and 1.62. □

Theorem 1.63. Every nonzero prime ideal in a PID is a maximal ideal.

Corollary 1.64. If R is a commutative ring such that the polynomial ring $R[x]$ is a PID, then R is necessarily a field.

Example 1.65. Here are a few quick examples.

- (1) We already know that $\mathbb{Z}[x]$ is not a PID, but the above corollary tells us again that it isn't since \mathbb{Z} is not a field.
- (2) The polynomial ring $\mathbb{Q}[x]$ is an eligible PID and it turns out that it is. In fact, $F[x]$ ends up being a PID for every field F .
- (3) The polynomial ring $\mathbb{Q}[x, y]$ turns out not to be a PID. The reason for this is that $\mathbb{Q}[x, y] = (\mathbb{Q}[x])[y]$ and $\mathbb{Q}[x]$ is not a field.

1.6 Euclidean Domains

Coming soon...