

Section 11.10: Taylor Series

Goal

We will introduce Taylor and Maclaurin Series, which are special types of power series. Not every power series takes the form of a geometric series. So, we need a more general method.

Taylor & Maclaurin Series

Suppose f has a power series representation (with $|x - a| < R$):

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots$$

Since these functions are equal, their derivatives agree:

$$f'(x) = \sum_{n=1}^{\infty} n c_n(x-a)^{n-1} = c_1 + 2c_2(x-a) + 3c_3(x-a)^2 + \cdots$$

This implies that

$$f'(a) = c_1 + 0 + 0 + \cdots = c_1.$$

That is,

$c_1 = \underline{\hspace{2cm}}$

Let's repeat this process with the second derivative. We see that

$$f''(x) = \sum_{n=2}^{\infty} n(n-1)c_n(x-a)^{n-2} = \underline{\hspace{10cm}}$$

This implies that

$$f''(a) = \underline{\hspace{3cm}} = \underline{\hspace{3cm}}.$$

Then

$c_2 = \underline{\hspace{2cm}}$

And again using the third derivative:

$$f'''(x) = \sum_{n=3}^{\infty} n(n-1)(n-2)c_n(x-a)^{n-3} = \underline{\hspace{10cm}}$$

This implies that

$$f'''(a) = \underline{\hspace{3cm}} = \underline{\hspace{3cm}}.$$

Then

$$c_3 = \underline{\hspace{2cm}}$$

If we continue this way, we'll obtain the following

$$c_n = \underline{\hspace{2cm}}$$

Note 1. Here are a couple of conventions:

1. $0! = 1! = 1$
2. $f^{(0)}(a) = f(a)$

Theorem 2. If f has a power series representation at $x = a$, then it must be of the form

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for $|x - a| < R$, where R is the radius of convergence.

The above power series is called the *Taylor Series of f centered at $x = a$* . In the special case that $a = 0$, we get

$$f(x) = \underline{\hspace{3cm}},$$

which is called the *Maclaurin series of f* .

Important Note 3. We can always compute a Taylor/Maclaurin series, but that does not mean that it is equal to the given function. We only know that a Taylor/Maclaurin Series is equal to a given function *if* the given function can be represented by a power series. We will only deal with these types of functions.

Example 4.

- (a) Find Maclaurin series for $f(x) = e^x$ and its radius of convergence (given that f has a power series representation).

- (b) Find Maclaurin series for $f(x) = \sin x$ and its radius of convergence (given that f has a power series representation).

Common Taylor Series

Here are some common Taylor Series.

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n \qquad R = 1$$

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \qquad R = \infty$$

$$\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \qquad R = \infty$$

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \qquad R = \infty$$

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \qquad R = 1$$

Taylor Series approximation

If a function has a Taylor Series representation, then we can use a finite number of terms to approximate the function. We define the *kth degree Taylor polynomial of f at $x = a$* to be

$$T_k(x) = \sum_{n=0}^k \frac{f^{(n)}(a)}{n!} (x - a)^n$$

Let's take a look at the [Taylor Series and Polynomials Applet](http://calculusapplets.com/) available at <http://calculusapplets.com/>.

Example 5.

(a) Use the 9th degree Taylor polynomial for $\arctan x$ to approximate π .

(b) Approximate the following integral using a 5th degree Taylor polynomial for $\sin x$.

$$\int_0^1 x \sin(x^3) \, dx$$