

# IMPARTIAL ACHIEVEMENT & AVOIDANCE GAMES FOR GENERATING FINITE GROUPS

---

ACGT Seminar at NAU

Dana C. Ernst

Northern Arizona University

September 26 & October 3, 2017

Joint work with Bret Benesh and Nándor Sieben

## Intuitive Definition

**Combinatorial Game Theory** (CGT) is the study of two-person games satisfying:

- Two players alternate making moves.
- No hidden information.
- No random moves.

## Combinatorial games

- Chess
- Go
- Connect Four
- Nim
- Tic-Tac-Toe
- X-Only Tic-Tac-Toe

## Combinatorial games

- Chess
- Go
- Connect Four
- Nim
- Tic-Tac-Toe
- X-Only Tic-Tac-Toe

## Non-combinatorial games

- Battleship (hidden information)
- Rock-Paper-Scissors (non-alternating and random)
- Poker (hidden information and random)

## Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

## Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

## Partizan

- Chess
- Go
- Connect Four
- Tic-Tac-Toe

## Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

## Partizan

- Chess
- Go
- Connect Four
- Tic-Tac-Toe

## Impartial

- Nim
- X-Only Tic-Tac-Toe

## Comments

- We are interested in impartial games.
- We will require that game sequence is finite and there are no ties.
- Player that moves first is called  $\alpha$  and second player is called  $\beta$ .
- **Normal Play:** The last player to move wins.
- **Misère Play:** The last player to move loses.



## Single-pile Nim

Start with a pile of  $n$  stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted  $*n$  (called a **nimber**).



## Single-pile Nim

Start with a pile of  $n$  stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted  $*n$  (called a **nimber**).



## Question

Is there an optimal strategy for either player?

## Single-pile Nim

Start with a pile of  $n$  stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted  $*n$  (called a **nimber**).



## Question

Is there an optimal strategy for either player?

## Answer

Boring:  $\alpha$  always wins; just take the whole pile.

## Single-pile Nim

Start with a pile of  $n$  stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted  $*n$  (called a **nimber**).



## Question

Is there an optimal strategy for either player?

## Answer

Boring:  $\alpha$  always wins; just take the whole pile.

## Multi-pile Nim


Start with  $k$  piles consisting of  $n_1, \dots, n_k$  stones, respectively. Each player chooses at least one stone from a single pile. The player that takes the last stone wins. Denoted  $*n_1 + \dots + *n_k$ .

## Example

Let's play  $*1 + *2 + *2$ . Here's a possible sequence.


## Example

Let's play  $*1 + *2 + *2$ . Here's a possible sequence.




## Example

Let's play  $*1 + *2 + *2$ . Here's a possible sequence.



## Example


Let's play  $*1 + *2 + *2$ . Here's a possible sequence.





## Example

Let's play  $*1 + *2 + *2$ . Here's a possible sequence.




## Example

Let's play  $*1 + *2 + *2$ . Here's a possible sequence.



## Example


Let's play  $*1 + *2 + *2$ . Here's a possible sequence.



In this case,  $\alpha$  wins.

## Example

Let's play  $*1 + *2 + *2$ . Here's a possible sequence.



In this case,  $\alpha$  wins.

## Question

In general, is there an optimal strategy for either player?

## Example

Let's play  $*1 + *2 + *2$ . Here's a possible sequence.



In this case,  $\alpha$  wins.

## Question

In general, is there an optimal strategy for either player?

## Answer

Short answer is yes: write sizes of piles in binary, do binary addition without carry (XOR), and if possible, hand your opponent a sum of 0. If players make optimal moves, this is only possible for one of the players.

## Definition

An **impartial game** is a finite set  $X$  of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

## Definition

An **impartial game** is a finite set  $X$  of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

- Two players take turns choosing a single available option in  $\text{Opt}(Q)$  of current position  $Q$ .
- Player who encounters empty option set cannot move and loses.
- **P-position**: previous player (player that just moved) wins
- **N-position**: next player (player that is about to move) wins

## Definition

An **impartial game** is a finite set  $X$  of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

- Two players take turns choosing a single available option in  $\text{Opt}(Q)$  of current position  $Q$ .
- Player who encounters empty option set cannot move and loses.
- **P-position**: previous player (player that just moved) wins
- **N-position**: next player (player that is about to move) wins

From perspective of the player that is about to move, a **P-position** is a **losing position** while an **N-position** is a **winning position**.



## Examples

- $*n$  is an N-position

## Examples

- $*n$  is an N-position
- $*1 + *1$  is a P-position

## Examples

- $*n$  is an N-position
- $*1 + *1$  is a P-position
- $*1 + *2$  is an N-position

### Examples

- $*n$  is an N-position
- $*1 + *1$  is a P-position
- $*1 + *2$  is an N-position
- Empty X-Only Tic-Tac-Toe board is an N-position

### Examples

- $*n$  is an N-position
- $*1 + *1$  is a P-position
- $*1 + *2$  is an N-position
- Empty X-Only Tic-Tac-Toe board is an N-position

## Definition

If  $G$  and  $H$  are games, then  $G + H$  is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

## Definition

If  $G$  and  $H$  are games, then  $G + H$  is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

## Theorem

$G + G$  is a P-position.

## Definition

If  $G$  and  $H$  are games, then  $G + H$  is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

## Theorem

$G + G$  is a P-position.

## Proof



Copy cat.



## Definition

$G_1 = G_2$  if and only if  $G_1 + G_2$  is a P-position.

Intuition: something akin to “copy cat” works.

## Definition

$G_1 = G_2$  if and only if  $G_1 + G_2$  is a P-position.

Intuition: something akin to “copy cat” works.

## Examples

$$\cdot \ast 1 + \ast 1 = \ast 0$$

## Definition

$G_1 = G_2$  if and only if  $G_1 + G_2$  is a P-position.

Intuition: something akin to “copy cat” works.

## Examples

·  $*1 + *1 = *0$  since  $*1 + *1 + *0$  is a P-position.

## Definition

$G_1 = G_2$  if and only if  $G_1 + G_2$  is a P-position.

Intuition: something akin to “copy cat” works.

## Examples

- $*1 + *1 = *0$  since  $*1 + *1 + *0$  is a P-position.
- $*1 + *2 = *3$

## Definition

$G_1 = G_2$  if and only if  $G_1 + G_2$  is a P-position.

Intuition: something akin to “copy cat” works.

## Examples

- $*1 + *1 = *0$  since  $*1 + *1 + *0$  is a P-position.
- $*1 + *2 = *3$  since  $*1 + *2 + *3$  is a P-position.

## Definition

$G_1 = G_2$  if and only if  $G_1 + G_2$  is a P-position.

Intuition: something akin to “copy cat” works.

## Examples

- $*1 + *1 = *0$  since  $*1 + *1 + *0$  is a P-position.
- $*1 + *2 = *3$  since  $*1 + *2 + *3$  is a P-position.

## Theorem

$G_1 = G_2$  if and only if  $G_1 + H$  and  $G_2 + H$  have the same outcome for all  $H$ .

### Definition

If  $A$  is a finite subset of nonnegative integers, then  $\text{mex}(A)$  is the smallest nonnegative integer not in  $A$ .

## Definition

If  $A$  is a finite subset of nonnegative integers, then  $\text{mex}(A)$  is the smallest nonnegative integer not in  $A$ .

## Examples

- $\text{mex}(\{0, 1, 2, 4, 5\}) = 3$
- $\text{mex}(\{1, 3\}) = 0$
- $\text{mex}(\{0, 1\}) = 2$
- $\text{mex}(\emptyset) = 0$



### Definition

If  $G$  is a game, then

$$\text{nim}(G) := \text{mex}(\{\text{nim}(Q) \mid Q \in \text{Opt}(G)\}).$$

This is a recursive definition. We start computing with terminal positions (empty option set).

## Definition

If  $G$  is a game, then

$$\text{nim}(G) := \text{mex}(\{\text{nim}(Q) \mid Q \in \text{Opt}(G)\}).$$

This is a recursive definition. We start computing with terminal positions (empty option set).

## Examples

- $\text{nim}(*0) = \text{mex}(\emptyset) = 0$
- $\text{nim}(*1) = \text{mex}(\{\text{nim}(*0)\}) = \text{mex}(\{0\}) = 1$
- $\text{nim}(*2) = \text{mex}(\{\text{nim}(*0), \text{nim}(*1)\}) = \text{mex}(\{0, 1\}) = 2$
- $\text{nim}(*n) = n$
- $\text{nim}(*1 + *1) = \text{mex}(\{\text{nim}(*1)\}) = \text{mex}(\{1\}) = 0$
- $\text{nim}(*1 + *2) = \text{mex}(\{\text{nim}(*2), \text{nim}(*1), \text{nim}(*1 + *1)\})$   
 $= \text{mex}(\{2, 1, 0\}) = 3$

## Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile:  $G = * \text{nim}(G)$

## Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile:  $G = * \text{nim}(G)$

## Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

## Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile:  $G = * \text{nim}(G)$

## Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

Loosely speaking, we can think of nim-numbers as “isomorphism” classes of games.

## Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile:  $G = * \text{nim}(G)$

## Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

Loosely speaking, we can think of nim-numbers as “isomorphism” classes of games.

## Theorem

2nd player  $\beta$  wins  $G$  if and only if  $G = *0$ .

Let  $G$  be a finite (possibly trivial) group.

Let  $G$  be a finite (possibly trivial) group.

## Generate Game

For the achievement game  $\text{GEN}(G)$ :



Let  $G$  be a finite (possibly trivial) group.

## Generate Game

For the achievement game  $\text{GEN}(G)$ :

- 1st player chooses any  $g_1 \in G$ .
- At  $k$ th turn, designated player selects  $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$  to create position  $\{g_1, \dots, g_k\}$ .
- Player wins on the  $n$ th turn if  $\langle g_1, \dots, g_n \rangle = G$ .

Let  $G$  be a finite (possibly trivial) group.

## Generate Game

For the achievement game  $\text{GEN}(G)$ :

- 1st player chooses any  $g_1 \in G$ .
- At  $k$ th turn, designated player selects  $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$  to create position  $\{g_1, \dots, g_k\}$ .
- Player wins on the  $n$ th turn if  $\langle g_1, \dots, g_n \rangle = G$ .

Positions of  $\text{GEN}(G)$  are subsets of terminal positions, which are certain generating sets of  $G$ .

## MATCH-UP



---

Name: LeBron James

Bret Benesh

---

Height: 6'8"

6'5"

Weight: 260 lbs

180 lbs

Age: 32 years

>32 years

Salary: \$30.96 million/year

\$0 million/year

Accolades: 3x NBA Champion  
4x NBA MVP  
2x Olympic gold medalist  
11x NBA All-Star

Never had a cavity  
Sagittarius

---

# LEBRON VS BRET: GAME ONE

GEN on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



---

LeBron

$P$

$\langle P \rangle$

Bret

---

# LEBRON VS BRET: GAME ONE

GEN on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	

# LEBRON VS BRET: GAME ONE

GEN on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$

# LEBRON VS BRET: GAME ONE

GEN on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	$S_3$	

# LEBRON VS BRET: GAME ONE

GEN on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	$S_3$	



# LEBRON VS BRET: GAME ONE

GEN on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	$S_3$	

Let  $G$  be a finite nontrivial group.

## Do Not Generate Game

For the **avoidance game**  $\text{DNG}(G)$ :

- 1st player chooses  $g_1 \in G$  such that  $\langle g_1 \rangle \neq G$ .
- At the  $k$ th turn, designated player selects  $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$  such that  $\langle g_1, \dots, g_k \rangle \neq G$  to create position  $\{g_1, \dots, g_k\}$ .
- Player that cannot select an element without building a generating set is loser.

Positions of  $\text{DNG}(G)$  are exactly the non-generating subsets of  $G$  and terminal positions are the maximal subgroups of  $G$ .

## LEBRON VS BRET: GAME TWO

DNG on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



---

LeBron

$P$

$\langle P \rangle$

Bret

---

# LEBRON VS BRET: GAME TWO

DNG on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	

## LEBRON VS BRET: GAME TWO

DNG on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$

# LEBRON VS BRET: GAME TWO

DNG on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$
$e$	$\{(1, 2, 3), (1, 3, 2), e\}$	$\mathbb{Z}_3$	

## LEBRON VS BRET: GAME TWO

DNG on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$
$e$	$\{(1, 2, 3), (1, 3, 2), e\}$	$\mathbb{Z}_3$	

# LEBRON VS BRET: GAME TWO

DNG on  $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	$P$	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	$\mathbb{Z}_3$	
	$\{(1, 2, 3), (1, 3, 2)\}$	$\mathbb{Z}_3$	$(1, 3, 2)$
$e$	$\{(1, 2, 3), (1, 3, 2), e\}$	$\mathbb{Z}_3$	



# LEBRON VS BRET: GAME THREE

DNG on  $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



---

LeBron

$P$

$\langle P \rangle$

Bret

---

# LEBRON VS BRET: GAME THREE

DNG on  $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	$P$	$\langle P \rangle$	Bret
	$\{r^2\}$	$\mathbb{Z}_2$	$r^2$

# LEBRON VS BRET: GAME THREE

DNG on  $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	$P$	$\langle P \rangle$	Bret
	$\{r^2\}$	$\mathbb{Z}_2$	$r^2$
$r^3$	$\{r^2, r^3\}$	$\mathbb{Z}_4$	

# LEBRON VS BRET: GAME THREE

DNG on  $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	$P$	$\langle P \rangle$	Bret
$r^3$	$\{r^2\}$	$\mathbb{Z}_2$	$r^2$
	$\{r^2, r^3\}$	$\mathbb{Z}_4$	
	$\{r^2, r^3, e\}$	$\mathbb{Z}_4$	$e$

# LEBRON VS BRET: GAME THREE

DNG on  $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	$P$	$\langle P \rangle$	Bret
	$\{r^2\}$	$\mathbb{Z}_2$	$r^2$
$r^3$	$\{r^2, r^3\}$	$\mathbb{Z}_4$	
	$\{r^2, r^3, e\}$	$\mathbb{Z}_4$	$e$
$r$	$\{r^2, r^2, e, r\}$	$\mathbb{Z}_4$	

# LEBRON VS BRET: GAME THREE


DNG on  $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	$P$	$\langle P \rangle$	Bret
	$\{r^2\}$	$\mathbb{Z}_2$	$r^2$
$r^3$	$\{r^2, r^3\}$	$\mathbb{Z}_4$	
	$\{r^2, r^3, e\}$	$\mathbb{Z}_4$	$e$
$r$	$\{r^2, r^2, e, r\}$	$\mathbb{Z}_4$	

# LEBRON VS BRET: GAME THREE

DNG on  $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	$P$	$\langle P \rangle$	Bret
$r^3$	$\{r^2\}$	$\mathbb{Z}_2$	$r^2$
	$\{r^2, r^3\}$	$\mathbb{Z}_4$	
$r$	$\{r^2, r^3, e\}$	$\mathbb{Z}_4$	$e$
	$\{r^2, r^2, e, r\}$	$\mathbb{Z}_4$	

- 1987: Harary and Anderson determine outcomes for abelian groups.



## SOME HISTORY

- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.


## SOME HISTORY

- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.
- 2014: Ernst and Sieben determine nim-numbers (and hence outcomes) for cyclic, dihedral, abelian.

## SOME HISTORY


- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.
- 2014: Ernst and Sieben determine nim-numbers (and hence outcomes) for cyclic, dihedral, abelian.
- 2016: Benesh, Ernst, and Sieben establish **subgroup-based criteria** for the determination of nim-numbers (and hence outcomes) for DNG, characterize **spectrum of nim-numbers for DNG**, determine nim-numbers for GEN and DNG for a variety of groups including generalized dihedral, symmetric, and alternating groups.

# REPRESENTATIVE GAME TREES




Representative game tree for  $\text{GEN}(S_3) = *3$

# STRUCTURE DIAGRAMS






$\text{DNG}(\mathbb{Z}_6)$



Structure diagram

# SIMPLIFIED STRUCTURE DIAGRAMS




$DNG(\mathbb{Z}_6 \times \mathbb{Z}_3)$


# SIMPLIFIED STRUCTURE DIAGRAMS




(a)  $\text{DNG}(\mathbb{D}_{4k})$




(b)  $\text{DNG}(\mathbb{D}_{2k+1})$




(c)  $\text{DNG}(\mathbb{D}_{4k+2})$



(d)  $\text{GEN}(\mathbb{D}_{4k})$



(e)  $\text{GEN}(\mathbb{D}_{2k+1})$



(f)  $\text{GEN}(\mathbb{D}_{4k+2})$

Simplified structure diagrams for dihedral groups

## Theorem (Ernst, Sieben)

If  $n \geq 2$ , then  $\text{nim}(\text{GEN}(\mathbb{Z}_n)) = \text{nim}(\text{DNG}(\mathbb{Z}_n)) + 1$ .



## Theorem (Ernst, Sieben)

If  $n \geq 2$ , then  $\text{nim}(\text{GEN}(\mathbb{Z}_n)) = \text{nim}(\text{DNG}(\mathbb{Z}_n)) + 1$ .

## Theorem (Ernst, Sieben)

If  $n \geq 2$ , then

$$\text{DNG}(\mathbb{Z}_n) = \begin{cases} *1, & n = 2 \\ *1, & n \equiv_2 1 \\ *0, & n \equiv_4 0 \\ *3, & n \equiv_4 2 \end{cases}$$

and

$$\text{GEN}(\mathbb{Z}_n) = \begin{cases} *2, & n = 2 \\ *2, & n \equiv_2 1 \\ *1, & n \equiv_4 0 \\ *4, & n \equiv_4 2 \end{cases}$$

## Theorem (Ernst, Sieben)

For  $n \geq 3$ , we have

$$\text{DNG}(\mathbb{D}_n) = \begin{cases} *3, & n \equiv_2 1 \\ *0, & n \equiv_2 0 \end{cases}$$

and

$$\text{GEN}(\mathbb{D}_n) = \begin{cases} *3, & n \equiv_2 1 \\ *0, & n \equiv_4 0 \\ *1, & n \equiv_4 2 \end{cases}$$

## Theorem (Ernst, Sieben)

If  $G$  is a finite nontrivial abelian group, then

$$\text{DNG}(G) = \begin{cases} *1, & G \text{ is nontrivial of odd order} \\ *1, & G = \mathbb{Z}_2 \\ *3, & G = \mathbb{Z}_2 \times \mathbb{Z}_{2k+1} \text{ with } k \geq 1 \\ *0, & \text{else} \end{cases}$$

$$\text{GEN}(G) = \begin{cases} *2, & |G| \text{ is odd and } d(G) \leq 2 \\ *1, & |G| \text{ is odd and } d(G) \geq 3 \\ *2, & G = \mathbb{Z}_2 \\ *1, & G = \mathbb{Z}_{4k} \text{ with } k \geq 1 \\ *4, & G = \mathbb{Z}_{4k+2} \text{ with } k \geq 1 \\ *1, & G = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_m \times \mathbb{Z}_k \text{ for } m, k \text{ odd} \\ *0, & \text{else} \end{cases}$$

### Theorem (Ernst, Sieben)

- If  $G$  is any finite nontrivial group, then  $\text{DNG}(G)$  is  $*0$ ,  $*1$ , or  $*3$ .

### Theorem (Ernst, Sieben)

- If  $G$  is any finite nontrivial group, then  $\text{DNG}(G)$  is  $*0$ ,  $*1$ , or  $*3$ .
- If  $|G|$  is odd and nontrivial, then  $\text{GEN}(G)$  is  $*1$  or  $*2$ .

## Theorem (Ernst, Sieben)

- If  $G$  is any finite nontrivial group, then  $\text{DNG}(G)$  is  $*0$ ,  $*1$ , or  $*3$ .
- If  $|G|$  is odd and nontrivial, then  $\text{GEN}(G)$  is  $*1$  or  $*2$ .

## Conjecture (In Progress)

If  $|G|$  is even, then  $\text{GEN}(G)$  is one of  $*0$ ,  $*1$ ,  $*2$ ,  $*3$ ,  $*4$ .

### Theorem (Benesh, Ernst, Sieben)

Let  $G$  be a finite nontrivial group.

- If all maximal subgroups are even, then  $\text{DNG}(G) = *0$ .
- If all maximal subgroups are odd, then  $\text{DNG}(G) = *1$ .
- If mixed maximal subgroups, then
  - If the even maximals cover  $G$ , then  $\text{DNG}(G) = *0$ .
  - If the even maximals do not cover  $G$ , then  $\text{DNG}(G) = *3$ .

### Theorem (Benesh, Ernst, Sieben)

Let  $G$  be a finite nontrivial group.

- If all maximal subgroups are even, then  $\text{DNG}(G) = *0$ .
- If all maximal subgroups are odd, then  $\text{DNG}(G) = *1$ .
- If mixed maximal subgroups, then
  - If the even maximals cover  $G$ , then  $\text{DNG}(G) = *0$ .
  - If the even maximals do not cover  $G$ , then  $\text{DNG}(G) = *3$ .

Using our “checklist” criteria, we have completely characterized DNG for nilpotent, generalized dihedral, generalized quaternion, symmetric, Coxeter, alternating, and some Rubik’s cube groups.



## Big Picture for DNG

- The players just race to fill up one maximal subgroup  $M$ .
- **The beginning of the game is a struggle to determine  $M$ .**
- $\alpha$  wants  $|M|$  to be odd.
- $\beta$  wants  $|M|$  to be even.

## Big Picture for DNG

- The players just race to fill up one maximal subgroup  $M$ .
- **The beginning of the game is a struggle to determine  $M$ .**
- $\alpha$  wants  $|M|$  to be odd.
- $\beta$  wants  $|M|$  to be even.

## Strategy

- $\alpha$  wants to pick an element not in any maximal subgroups of even order.
- $\beta$  wants to pick an involution.

### What's left to work on?

- Wrap up spectrum of GEN?
- Wrap up characterization of GEN for nilpotent groups?
- Are there nice results for products and quotients?
- Is it possible to characterize the nim-numbers of GEN in terms of covering conditions by maximal subgroups similar to what we did for DNG?
- What about other “closure systems”? We are currently tinkering with convex hulls of finitely many points in the plane.

Thanks!