

IMPARTIAL ACHIEVEMENT & AVOIDANCE GAMES FOR GENERATING FINITE GROUPS

ACGT Seminar at NAU

Dana C. Ernst

Northern Arizona University

September 27, 2017

Joint work with Bret Benesh and Nándor Sieben

Intuitive Definition

Combinatorial Game Theory (CGT) is the study of two-person games satisfying:

- Two players alternate making moves.
- No hidden information.
- No random moves.

Combinatorial games

- Chess
- Go
- Connect Four
- Nim
- Tic-Tac-Toe
- X-Only Tic-Tac-Toe

Combinatorial games

- Chess
- Go
- Connect Four
- Nim
- Tic-Tac-Toe
- X-Only Tic-Tac-Toe

Non-combinatorial games

- Battleship (hidden information)
- Rock-Paper-Scissors (non-alternating and random)
- Poker (hidden information and random)

Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

Partizan

- Chess
- Go
- Connect Four
- Tic-Tac-Toe

Definition

A combinatorial game is called **impartial** if the move options are the same for both players. Otherwise, the game is called **partizan**.

Partizan

- Chess
- Go
- Connect Four
- Tic-Tac-Toe

Impartial

- Nim
- X-Only Tic-Tac-Toe

Comments

- We are interested in impartial games.
- We will require that game sequence is finite and there are no ties.
- Player that moves first is called α and second player is called β .
- **Normal Play:** The last player to move wins.
- **Misère Play:** The last player to move loses.

Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Question

Is there an optimal strategy for either player?

Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Question

Is there an optimal strategy for either player?

Answer

Boring: α always wins; just take the whole pile.

Single-pile Nim

Start with a pile of n stones. Each player chooses at least one stone from the pile. The player that takes the last stone wins. Game is denoted $*n$ (called a **nimber**).



Question

Is there an optimal strategy for either player?

Answer

Boring: α always wins; just take the whole pile.

Multi-pile Nim

Start with k piles consisting of n_1, \dots, n_k stones, respectively. Each player chooses at least one stone from a single pile. The player that takes the last stone wins. Denoted $*n_1 + \dots + *n_k$.

Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.


Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.




Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.




Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.




Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



Example


Let's play $*1 + *2 + *2$. Here's a possible sequence.



In this case, α wins.

Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



In this case, α wins.

Question

In general, is there an optimal strategy for either player?

Example

Let's play $*1 + *2 + *2$. Here's a possible sequence.



In this case, α wins.

Question

In general, is there an optimal strategy for either player?

Answer

Short answer is yes: write sizes of piles in binary, do binary addition without carry (XOR), and if possible, hand your opponent a sum of 0. If players make optimal moves, this is only possible for one of the players.

Definition

An **impartial game** is a finite set X of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

Definition

An **impartial game** is a finite set X of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

- Two players take turns choosing a single available option in $\text{Opt}(Q)$ of current position Q .
- Player who encounters empty option set cannot move and loses.
- **P-position**: previous player (player that just moved) wins
- **N-position**: next player (player that is about to move) wins

Definition

An **impartial game** is a finite set X of positions together with a starting position and a collection

$$\{\text{Opt}(Q) \subseteq X \mid Q \in X\}$$

of possible options.

- Two players take turns choosing a single available option in $\text{Opt}(Q)$ of current position Q .
- Player who encounters empty option set cannot move and loses.
- **P-position**: previous player (player that just moved) wins
- **N-position**: next player (player that is about to move) wins

From perspective of the player that is about to move, a **P-position** is a **losing position** while an **N-position** is a **winning position**.

Examples

- $*n$ is an N-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position
- $*1 + *2$ is an N-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position
- $*1 + *2$ is an N-position
- Empty X-Only Tic-Tac-Toe board is an N-position

Examples

- $*n$ is an N-position
- $*1 + *1$ is a P-position
- $*1 + *2$ is an N-position
- Empty X-Only Tic-Tac-Toe board is an N-position

Definition

If G and H are games, then $G + H$ is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

Definition

If G and H are games, then $G + H$ is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

Theorem

$G + G$ is a P-position.

Definition

If G and H are games, then $G + H$ is the game where each player makes a move in one of the games. Set of options:

$$\text{Opt}_{G+H}(S + T) := \{Q + T \mid Q \in \text{Opt}_G(S)\} \cup \{S + R \mid R \in \text{Opt}_H(T)\}$$

Theorem

$G + G$ is a P-position.

Proof



Copy cat.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

$$\cdot \ast 1 + \ast 1 = \ast 0$$

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

· $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

- $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.
- $*1 + *2 = *3$

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

- $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.
- $*1 + *2 = *3$ since $*1 + *2 + *3$ is a P-position.

Definition

$G_1 = G_2$ if and only if $G_1 + G_2$ is a P-position.

Intuition: something akin to “copy cat” works.

Examples

- $*1 + *1 = *0$ since $*1 + *1 + *0$ is a P-position.
- $*1 + *2 = *3$ since $*1 + *2 + *3$ is a P-position.

Theorem

$G_1 = G_2$ if and only if $G_1 + H$ and $G_2 + H$ have the same outcome for all H .

Definition

If A is a finite subset of nonnegative integers, then $\text{mex}(A)$ is the smallest nonnegative integer not in A .

Definition

If A is a finite subset of nonnegative integers, then $\text{mex}(A)$ is the smallest nonnegative integer not in A .

Examples

- $\text{mex}(\{0, 1, 2, 4, 5\}) = 3$
- $\text{mex}(\{1, 3\}) = 0$
- $\text{mex}(\{0, 1\}) = 2$
- $\text{mex}(\emptyset) = 0$

Definition

If G is a game, then

$$\text{nim}(G) := \text{mex}(\{\text{nim}(Q) \mid Q \in \text{Opt}(G)\}).$$

This is a recursive definition. We start computing with terminal positions (empty option set).

Definition

If G is a game, then

$$\text{nim}(G) := \text{mex}(\{\text{nim}(Q) \mid Q \in \text{Opt}(G)\}).$$

This is a recursive definition. We start computing with terminal positions (empty option set).

Examples

- $\text{nim}(*0) = \text{mex}(\emptyset) = 0$
- $\text{nim}(*1) = \text{mex}(\{\text{nim}(*0)\}) = \text{mex}(\{0\}) = 1$
- $\text{nim}(*2) = \text{mex}(\{\text{nim}(*0), \text{nim}(*1)\}) = \text{mex}(\{0, 1\}) = 2$
- $\text{nim}(*n) = n$
- $\text{nim}(*1 + *1) = \text{mex}(\{\text{nim}(*1)\}) = \text{mex}(\{1\}) = 0$
- $\text{nim}(*1 + *2) = \text{mex}(\{\text{nim}(*2), \text{nim}(*1), \text{nim}(*1 + *1)\})$
 $= \text{mex}(\{2, 1, 0\}) = 3$

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

Loosely speaking, we can think of nim-numbers as “isomorphism” classes of games.

Theorem (Sprague–Grundy)

Every game is equivalent to a single Nim pile: $G = * \text{nim}(G)$

Big Picture

Fundamental problem in the theory of impartial combinatorial games is the determination of the nim-number of the game.

Loosely speaking, we can think of nim-numbers as “isomorphism” classes of games.

Theorem

2nd player β wins G if and only if $G = *0$.

Let G be a finite (possibly trivial) group.

Let G be a finite (possibly trivial) group.

Generate Game

For the achievement game $\text{GEN}(G)$:

Let G be a finite (possibly trivial) group.

Generate Game

For the achievement game $\text{GEN}(G)$:

- 1st player chooses any $g_1 \in G$.
- At k th turn, designated player selects $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$ to create position $\{g_1, \dots, g_k\}$.
- Player wins on the n th turn if $\langle g_1, \dots, g_n \rangle = G$.

Let G be a finite (possibly trivial) group.

Generate Game

For the achievement game $\text{GEN}(G)$:

- 1st player chooses any $g_1 \in G$.
- At k th turn, designated player selects $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$ to create position $\{g_1, \dots, g_k\}$.
- Player wins on the n th turn if $\langle g_1, \dots, g_n \rangle = G$.

Positions of $\text{GEN}(G)$ are subsets of terminal positions, which are certain generating sets of G .

MATCH-UP



Name: LeBron James

Bret Benesh

Height: 6'8"

6'5"

Weight: 260 lbs

180 lbs

Age: 32 years

>32 years

Salary: \$30.96 million/year

\$0 million/year

Accolades: 3x NBA Champion
4x NBA MVP
2x Olympic gold medalist
11x NBA All-Star

Never had a cavity
Sagittarius

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron

P

$\langle P \rangle$

Bret

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	S_3	

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	$(1, 3, 2)$
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	S_3	

LEBRON VS BRET: GAME ONE

GEN on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	$(1, 3, 2)$
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	
$(1, 2)$	$\{(1, 2, 3), (1, 3, 2), (1, 2)\}$	S_3	

Let G be a finite nontrivial group.

Do Not Generate Game

For the **avoidance game** $\text{DNG}(G)$:

- 1st player chooses $g_1 \in G$ such that $\langle g_1 \rangle \neq G$.
- At the k th turn, designated player selects $g_k \in G \setminus \{g_1, \dots, g_{k-1}\}$ such that $\langle g_1, \dots, g_k \rangle \neq G$ to create position $\{g_1, \dots, g_k\}$.
- Player that cannot select an element without building a generating set is loser.

Positions of $\text{DNG}(G)$ are exactly the non-generating subsets of G and terminal positions are the maximal subgroups of G .

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron

P

$\langle P \rangle$

Bret

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
e	$\{(1, 2, 3), (1, 3, 2), e\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
e	$\{(1, 2, 3), (1, 3, 2), e\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME TWO

DNG on $S_3 = \{e, (1, 2), (1, 3), (2, 3), (1, 2, 3), (1, 3, 2)\}$



LeBron	P	$\langle P \rangle$	Bret
$(1, 2, 3)$	$\{(1, 2, 3)\}$	\mathbb{Z}_3	
	$\{(1, 2, 3), (1, 3, 2)\}$	\mathbb{Z}_3	$(1, 3, 2)$
e	$\{(1, 2, 3), (1, 3, 2), e\}$	\mathbb{Z}_3	

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron

P

$\langle P \rangle$

Bret

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
	$\{r^2\}$	\mathbb{Z}_2	r^2

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
	$\{r^2\}$	\mathbb{Z}_2	r^2
r^3	$\{r^2, r^3\}$	\mathbb{Z}_4	

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
r^3	$\{r^2\}$	\mathbb{Z}_2	r^2
	$\{r^2, r^3\}$	\mathbb{Z}_4	
	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
	$\{r^2\}$	\mathbb{Z}_2	r^2
r^3	$\{r^2, r^3\}$	\mathbb{Z}_4	
	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e
r	$\{r^2, r^2, e, r\}$	\mathbb{Z}_4	

LEBRON VS BRET: GAME THREE


DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
	$\{r^2\}$	\mathbb{Z}_2	r^2
r^3	$\{r^2, r^3\}$	\mathbb{Z}_4	
	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e
r	$\{r^2, r^2, e, r\}$	\mathbb{Z}_4	

LEBRON VS BRET: GAME THREE

DNG on $D_8 = \langle r, s \rangle = \{e, r, r^2, r^3, s, rs, r^2s, r^3s\}$



LeBron	P	$\langle P \rangle$	Bret
r^3	$\{r^2\}$	\mathbb{Z}_2	r^2
	$\{r^2, r^3\}$	\mathbb{Z}_4	
r	$\{r^2, r^3, e\}$	\mathbb{Z}_4	e
	$\{r^2, r^2, e, r\}$	\mathbb{Z}_4	

- 1987: Harary and Anderson determine outcomes for abelian groups.

SOME HISTORY

- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.


SOME HISTORY

- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.
- 2014: Ernst and Sieben determine nim-numbers (and hence outcomes) for cyclic, dihedral, abelian.

SOME HISTORY


- 1987: Harary and Anderson determine outcomes for abelian groups.
- 1988: Barnes establishes **element-based criteria** for who wins DNG, assorted GEN results.
- 2014: Ernst and Sieben determine nim-numbers (and hence outcomes) for cyclic, dihedral, abelian.
- 2016: Benesh, Ernst, and Sieben establish **subgroup-based criteria** for the determination of nim-numbers (and hence outcomes) for DNG, characterize **spectrum of nim-numbers for DNG**, determine nim-numbers for GEN and DNG for a variety of groups including generalized dihedral, symmetric, and alternating groups.

REPRESENTATIVE GAME TREES




Representative game tree for $\text{GEN}(S_3) = *3$

STRUCTURE DIAGRAMS





$\text{DNG}(\mathbb{Z}_6)$




Structure diagram

SIMPLIFIED STRUCTURE DIAGRAMS




$\text{DNG}(\mathbb{Z}_6 \times \mathbb{Z}_3)$




SIMPLIFIED STRUCTURE DIAGRAMS




(a) $\text{DNG}(\mathbb{D}_{4k})$




(b) $\text{DNG}(\mathbb{D}_{2k+1})$




(c) $\text{DNG}(\mathbb{D}_{4k+2})$



(d) $\text{GEN}(\mathbb{D}_{4k})$



(e) $\text{GEN}(\mathbb{D}_{2k+1})$



(f) $\text{GEN}(\mathbb{D}_{4k+2})$

Simplified structure diagrams for dihedral groups

Theorem (Ernst, Sieben)

If $n \geq 2$, then $\text{nim}(\text{GEN}(\mathbb{Z}_n)) = \text{nim}(\text{DNG}(\mathbb{Z}_n)) + 1$.

Theorem (Ernst, Sieben)

If $n \geq 2$, then $\text{nim}(\text{GEN}(\mathbb{Z}_n)) = \text{nim}(\text{DNG}(\mathbb{Z}_n)) + 1$.

Theorem (Ernst, Sieben)

If $n \geq 2$, then

$$\text{DNG}(\mathbb{Z}_n) = \begin{cases} *1, & n = 2 \\ *1, & n \equiv_2 1 \\ *0, & n \equiv_4 0 \\ *3, & n \equiv_4 2 \end{cases}$$

and

$$\text{GEN}(\mathbb{Z}_n) = \begin{cases} *2, & n = 2 \\ *2, & n \equiv_2 1 \\ *1, & n \equiv_4 0 \\ *4, & n \equiv_4 2 \end{cases}$$

Theorem (Ernst, Sieben)

For $n \geq 3$, we have

$$\text{DNG}(\mathbb{D}_n) = \begin{cases} *3, & n \equiv_2 1 \\ *0, & n \equiv_2 0 \end{cases}$$

and

$$\text{GEN}(\mathbb{D}_n) = \begin{cases} *3, & n \equiv_2 1 \\ *0, & n \equiv_4 0 \\ *1, & n \equiv_4 2 \end{cases}$$

Theorem (Ernst, Sieben)

If G is a finite nontrivial abelian group, then

$$\text{DNG}(G) = \begin{cases} *1, & G \text{ is nontrivial of odd order} \\ *1, & G = \mathbb{Z}_2 \\ *3, & G = \mathbb{Z}_2 \times \mathbb{Z}_{2k+1} \text{ with } k \geq 1 \\ *0, & \text{else} \end{cases}$$

$$\text{GEN}(G) = \begin{cases} *2, & |G| \text{ is odd and } d(G) \leq 2 \\ *1, & |G| \text{ is odd and } d(G) \geq 3 \\ *2, & G = \mathbb{Z}_2 \\ *1, & G = \mathbb{Z}_{4k} \text{ with } k \geq 1 \\ *4, & G = \mathbb{Z}_{4k+2} \text{ with } k \geq 1 \\ *1, & G = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_m \times \mathbb{Z}_k \text{ for } m, k \text{ odd} \\ *0, & \text{else} \end{cases}$$

Theorem (Ernst, Sieben)

- If G is any finite nontrivial group, then $\text{DNG}(G)$ is $*0$, $*1$, or $*3$.

Theorem (Ernst, Sieben)

- If G is any finite nontrivial group, then $\text{DNG}(G)$ is $*0$, $*1$, or $*3$.
- If $|G|$ is odd and nontrivial, then $\text{GEN}(G)$ is $*1$ or $*2$.

Theorem (Ernst, Sieben)

- If G is any finite nontrivial group, then $\text{DNG}(G)$ is $*0$, $*1$, or $*3$.
- If $|G|$ is odd and nontrivial, then $\text{GEN}(G)$ is $*1$ or $*2$.

Conjecture (In Progress)

If $|G|$ is even, then $\text{GEN}(G)$ is one of $*0$, $*1$, $*2$, $*3$, $*4$.

Theorem (Benesh, Ernst, Sieben)

Let G be a finite nontrivial group.

- If all maximal subgroups are even, then $\text{DNG}(G) = *0$.
- If all maximal subgroups are odd, then $\text{DNG}(G) = *1$.
- If mixed maximal subgroups, then
 - If the even maximals cover G , then $\text{DNG}(G) = *0$.
 - If the even maximals do not cover G , then $\text{DNG}(G) = *3$.

Theorem (Benesh, Ernst, Sieben)

Let G be a finite nontrivial group.

- If all maximal subgroups are even, then $\text{DNG}(G) = *0$.
- If all maximal subgroups are odd, then $\text{DNG}(G) = *1$.
- If mixed maximal subgroups, then
 - If the even maximals cover G , then $\text{DNG}(G) = *0$.
 - If the even maximals do not cover G , then $\text{DNG}(G) = *3$.

Using our “checklist” criteria, we have completely characterized DNG for nilpotent, generalized dihedral, generalized quaternion, symmetric, Coxeter, alternating, and some Rubik’s cube groups.

Big Picture for DNG

- The players just race to fill up one maximal subgroup M .
- **The beginning of the game is a struggle to determine M .**
- α wants $|M|$ to be odd.
- β wants $|M|$ to be even.

Big Picture for DNG

- The players just race to fill up one maximal subgroup M .
- **The beginning of the game is a struggle to determine M .**
- α wants $|M|$ to be odd.
- β wants $|M|$ to be even.

Strategy

- α wants to pick an element not in any maximal subgroups of even order.
- β wants to pick an involution.

What's left to work on?

- Wrap up spectrum of GEN?
- Wrap up characterization of GEN for nilpotent groups?
- Are there nice results for products and quotients?
- Is it possible to characterize the nim-numbers of GEN in terms of covering conditions by maximal subgroups similar to what we did for DNG?
- What about other “closure systems”? We are currently tinkering with convex hulls of finitely many points in the plane.

Thanks!