

MTH 302: Linear Algebra and Differential Equations

Parts of these activities will be assigned to Application/Analysis 10. Those parts will be announced later.

1: Solving a second-order equation

1. Go back to the Class Prep and consider $y'' - y' - 12y = 0$.
 - (a) Make the guess $y = e^{rt}$ for a solution. Find y' and y'' and then substitute into the left side.
 - (b) Find the values of r that solve the equation. (There will be two of them.)
 - (c) Replace those values of r back into the guess $y = e^{rt}$. Then check that both functions individually solve the differential equation.
 - (d) Form a linear combination of those two functions using arbitrary constants c_1 and c_2 as weights, and check that the linear combination is also a solution to the DE.
2. Mimic this process to find the general solution to each of the following second-order DEs. Each group will be responsible for one of these and putting the work on the board. But do more for practice.
 - (a) $y'' + y' - 2y = 0$
 - (b) $y'' - y = 0$
 - (c) $y'' + 3y' = 0$
 - (d) $y'' = 0$
 - (e) $y'' + 4y' + 3y = 0$
 - (f) $y'' + y' - y = 0$

2: Application to spring-mass systems

In a spring-mass system, the displacement $y(t)$ of the mass from its natural equilibrium is governed by the equation

$$y'' + \frac{c}{m}y' + \frac{k}{m}y = \frac{1}{m}F(t)$$

where c is a damping constant (coming from friction within the system), k is the spring constant, m is the mass of the suspended object, and F is a forcing function (some force from outside the system acting upon it).

For an *unforced* system with $c = 3$, $k = 2$, and $m = 1$, determine the displacement of the mass at time t is the system is set in motion with initial displacement $y(0) = 2$ and initial velocity $y'(0) = 1$.

3: What if the roots aren't real numbers?

Solve the second-order DE:

$$y'' + 2y' + 10y = 0$$

using the method you've learned.

1. Set up and solve the characteristic equation. What's different this time?
2. Let r be one of the characteristic roots. Go ahead and set up $y = e^{rt}$ as a solution, as usual. Then simplify this using Euler's formula.
3. The result of the previous part will give you *two* functions: One attached to the real part, and a second attached to the imaginary part. Check that *both* of these solve the original DE.
4. Using the results of the previous question, state the general solution to the DE.