

Your Paper

You

August 16, 2023

Abstract

Your abstract.

1 Introduction

Your introduction goes here! Simply start writing your document and use the Recompile button to view the updated PDF preview. Examples of commonly used commands and features are listed below, to help you get started.

Once you're familiar with the editor, you can find various project settings in the Overleaf menu, accessed via the button in the very top left of the editor. To view tutorials, user guides, and further documentation, please visit our [help library](#), or head to our plans page to [choose your plan](#).

2 Derivation

2.1 τ_{ij} Transnational Search Time

$$J_{ij} = k_{ij}c_i c_j$$

where

$$k_{ij} = 4\pi(i+j)D_{ij}c_i c_j$$

where i, j the radius of two interacting particles we are considering, D_{ij} is the effective transnational diffusion constant defined as

$$D_{ij} = D_i + D_j \tag{1}$$

$$= \frac{k_B T}{6\pi i \mu} + \frac{k_B T}{6\pi j \mu} \tag{2}$$

Thus,

$$k_{ij} = \frac{2k_B T}{3\mu} \frac{(i+j)^2}{ij}$$
$$J_{ij} = \frac{2k_B T}{3\mu} \frac{(i+j)^2}{ij} c_i c_j$$

this is number of collision per sec per square meter, to find the time τ_{ij} for each collision (time for origami to meet within their interactive region), take inverse

$$\tau_{ij} = \frac{3ij\mu}{2k_B T(i+j)^2} \frac{1}{c_i c_j}$$

2.2 τ_s Rotational Search Time

$$\tau_s = \frac{1}{D_\theta} \frac{\ln 4}{\alpha} \times \frac{1}{6}$$

$\frac{1}{6}$ account for 6 sticky ends, this is not accurate, just a rough approximation. $\alpha = \frac{\pi r_c^2}{2\pi r^2}$ is the relative capture surface, in our case, it can be approximated as following

$$\alpha = \frac{\frac{1}{\sqrt{2}}(l + \frac{Rg}{2})R_g}{(l + \frac{Rg}{2})^2 2\pi} \quad (3)$$

$$\approx \frac{2}{3\sqrt{2}} \quad (4)$$

D_θ is the effective rotational diffusion constant, which can be computed as

$$D_\theta = D_{\theta i} + D_{\theta j} \quad (5)$$

$$= \frac{k_B T}{8\pi\mu} \left(\frac{1}{i^3} + \frac{1}{j^3} \right) \quad (6)$$

$$= \frac{k_B T}{8\pi\mu} \frac{i^3 + j^3}{i^3 j^3} \quad (7)$$

Therefore

$$\tau_s \approx \frac{1}{D_\theta} \frac{\ln 4 \sqrt{2}}{3} \frac{1}{6} \quad (8)$$

$$= \frac{4\sqrt{2} \ln 4 \pi \mu}{9k_B T} \frac{i^3 j^3}{i^3 + j^3} \quad (9)$$

2.3 τ_c Collision Time

This is the time for two origami to remain in active range. Simply using

$$x = \sqrt{6D_{ij}\tau_c}$$

and set distance $x = 5nm$, which is approximately two third of the length of the sticky end plus Ts.

$$\tau_c = \frac{H\pi\mu}{k_B T} \frac{ij}{i+j}$$

where $H \approx 2.5 \times 10^{-17} m^2$.

2.4 τ Total Time For Two Origami To Combine in $1m^3$

$$\tau = \tau_{ij} \left(1 + \frac{\tau_s}{\tau_c} \right) \quad (10)$$

$$= \frac{3\mu}{2k_B T} \left(\frac{ij}{(i+j)^2} + \frac{4\sqrt{2} \ln 4}{9H} \frac{i^3 j^3}{(i^3 + j^3)(i+j)} \right) \frac{1}{c_i c_j} \quad (11)$$

$$= \frac{1}{M} \left(\frac{ij}{(i+j)^2} + N \frac{i^3 j^3}{(i^3 + j^3)(i+j)} \right) \frac{1}{c_i c_j} \quad (12)$$

where $M = \frac{2k_B T}{3\mu}$, $N = \frac{4\sqrt{2} \ln 4}{9H}$ are constant. Then, we can obtain the collision rate per meter squared per second by simply take reciprocal of τ

$$n(i, j) = \left(\frac{M(i+j)^2}{ij + N \frac{i^3 j^3 (i+j)}{(i^3 + j^3)}} \right) c_i c_j$$

where

$$k(i, j) = \frac{M(i+j)^2}{ij + N \frac{i^3 j^3 (i+j)}{(i^3 + j^3)}}$$

is the kernel we want.

References