

SELECT

DIAGNOSTIC IMAGING

2350 OCEAN AVENUE, SUITE 4
[REDACTED] NY 11229
TEL: (718) 676-4550 FAX: (718) 676-4557

Farzam Kashanian, M.D.
MEDICAL DIRECTOR

Report Details

Patient [REDACTED]
Patient [REDACTED]
Date of Birth: [REDACTED]
Gender: F
Accession: [REDACTED]
Referring Physician: [REDACTED]

[REDACTED]
[REDACTED]
[REDACTED]

Report Text

EXAM: ULTRASOUND GUIDED CORE NEEDLE BIOPSY OF TWO SITES IN THE RIGHT BREAST AND ONE SITE IN THE LEFT BREAST FOLLOWED BY CLIP PLACEMENT
EXAM: DIAGNOSTIC BILATERAL MAMMOGRAM WITH ICAD

History: This is followup to the mammogram and sonogram of [REDACTED]

Technique: An initial targeted ultrasound was performed using a high frequency linear transducer.

Within the right retroareolar region, a lobulated irregular mass measures 1.8 cm. At the right 11 o'clock axis periareolar region, a lobulated soft tissue nodule measures 1.0 cm. At the left 2 o'clock axis, 5 cm from the nipple, a hypoechoic nodule measures 0.6 cm.

Consent: The patient [REDACTED] and alternative to the procedure. The risks include infection, bleeding. Informed consent was given by the patient.

Procedure: Both breasts was prepped and draped in the usual sterile fashion. 1% lidocaine was used to infiltrate the skin and subcutaneous tissues.

Under ultrasound guidance, a core needle biopsy was performed upon the right retroareolar nodule. A marker was deployed.

Under ultrasound guidance, a core needle biopsy was performed upon the right 11 o'clock nodule. Marker was then deployed.

Under ultrasound guidance, fine needle aspiration/core needle biopsy was performed upon the 0.6 left 2 o'clock nodule. A marker was then deployed.

The patient [REDACTED] did not reveal a hematoma. The samples were submitted to CBL pathology laboratories for further evaluation.

Bilateral diagnostic MLO and Craniocaudal views were performed that demonstrate bilateral markers with postbiopsy changes.

IMPRESSION:

STATUS POST ULTRASOUND GUIDED BIOPSY OF RIGHT RETROAREOLAR IRREGULAR NODULE, RIGHT 11 O'CLOCK NODULE AND LEFT 2 O'CLOCK NODULE.

RESULTS: AS PER THE BREAST PATHOLOGY REPORT, THE DIAGNOSIS FOR THE RIGHT RETROAREOLAR NODULE AND RIGHT 11 O'CLOCK NODULE ARE INFILTRATING DUCTAL CARCINOMA, ARCHITECTURAL GRADE 3. THE DIAGNOSIS FOR THE LEFT 2 O'CLOCK NODULE IS BENIGN BREAST TISSUE AND STROMAL FIBROSIS. THESE FINDINGS ARE CONCORDANT. FOLLOWUP WITH SURGICAL CONSULTATION IS ADVISED. BILATERAL BREAST MRI WITH CONTRAST ENHANCEMENT IS RECOMMENDED.

BI-RADS CATEGORY 6(KNOWN BIOPSY-PROVEN MALIGNANCY-APPROPRIATE ACTION SHOULD BE TAKEN)

THE RESULTS WERE REPORTED TO IRADA AT DR TRAKHTENBERG'S OFFICE ON

FARZAM KASHANIAN MD

Signed by: Kashanian, Farzam
Transcribed By: TRANSCRIPTIONIST, EXCEL

Date Signed:
Date Transcribed: