



Results

MR BREAST W/WO CON W/CAD BILAT (Order)

MR BREAST W/WO CON W/CAD BILAT

Order:

Impression

Biopsy-proven DCIS, ultrasound underestimates the extent of disease which reaches up to 4.6 centimeters on MRI. The non mass enhancement extends within 0.7 centimeters from the nipple areola complex.

RECOMMENDATION: Surgical Consultation
Side: Right

BI-RADS MRI: 6: Known Malignancy

Breast Cancer Risk Assessment
The American College of Radiology recommends that all women undergo breast cancer risk assessment by age 25 so that those at higher-than-average risk can be identified and appropriate screening initiated. Black, Asian, Hispanic and Ashkenazi Jewish women are at higher risk in general. One widely used risk assessment tool is the Tyrer-Cuzick model at ibis.ikonopedia.com. The American Cancer Society, American College of Radiology and Society of Breast Imaging recommend screening breast MRI for women with 20% or greater lifetime risk of breast cancer.

Electronically Signed By: Roberto Lo Gullo on November 14, 2025 07:52 PM

I, Roberto Lo Gullo, have personally reviewed this report and concur with its findings and conclusions, as affirmed by my electronic signature.

Narrative

EXAMINATION:
BILATERAL BREAST MRI WITH AND WITHOUT INTRAVENOUS CONTRAST WITH COMPUTER-AIDED DETECTION

CLINICAL INFORMATION:
Biopsy-proven DCIS, staging

TECHNIQUE:
Magnetic resonance imaging of the breasts was performed using multisequence, multiplanar technique with and without intravenous contrast. Subtraction images were also archived.

Computer-Aided Detection: CAD analysis facilitated the interpretation of the contrast enhanced (DCE) series. CAD analysis was performed and 3D post-processing images were reconstructed and interpreted on an independent workstation by the radiologist,

Contrast: GADOPICLENOL (ELUCIREM) 0.5 MMOL/ML IV SOLN 6.6 mL

COMPARISON:
None

FINDINGS:
BREASTS: Fibroglandular tissue: Heterogeneous fibroglandular tissue with moderate background parenchymal enhancement

RIGHT BREAST: In the superior breast there is heterogeneous non mass enhancement measuring 4.6 x 2.7 x 4.3

centimeters consistent with biopsy-proven DCIS. The suspicious enhancement appears to be 0.7 centimeters distant from the nipple-areolar complex.

LEFT BREAST:No suspicious findings

OTHER: None.

Exam End: 12:07

Specimen Collected: 00:00

Received From: Weill Cornell Medicine, Columbia
Physicians, and New York Presbyterian Hospital

Last Resulted: 19:56

Result Received: 07:24

 View Encounter  Received Information