



Results

US BREAST DIAGNOSTIC BILAT (Order)

US BREAST DIAGNOSTIC BILAT

Order:

Impression

Indeterminate ultrasound finding in the upper right breast area of palpable concern for which representative ultrasound-guided biopsy is recommended. No

RECOMMENDATION: Ultrasound Biopsy
Side: Right
Follow-Up: Prompt follow up

The patient

BI-RADS mammogram: 2: Benign
BI-RADS sonogram: 4: Suspicious

Breast Cancer Risk Assessment
The American College of Radiology recommends that all women undergo breast cancer risk assessment by age 25 so that those at higher-than-average risk can be identified and appropriate screening initiated. Black, Asian, Hispanic and Ashkenazi Jewish women are at higher risk in general. One widely used risk assessment tool is the Tyrer-Cuzick model at ibis.ikonopedia.com. The American Cancer Society, American College of Radiology and Society of Breast Imaging recommend screening breast MRI for women with 20% or greater lifetime risk of breast cancer.

Electronically Signed By: Mark Francescone on October 09, 2025 05:22 PM

I, , have personally reviewed this report and concur with its findings and conclusions, as affirmed by my electronic signature.

Narrative

EXAMINATION:
BILATERAL DIAGNOSTIC DIGITAL MAMMOGRAM WITH TOMOSYNTHESIS AND COMPUTER-AIDED DETECTION and BILATERAL BREAST SONOGRAM COMPLETE

CLINICAL INFORMATION:
70-year-old female who presents for screening mammogram and also complains of a palpable lump in the right periareolar region. Date of last clinical breast exam: The patient

TECHNIQUE:
Mammogram: CC and MLO views including tomosynthesis and 2D synthetic views were obtained. This examination was evaluated with computer-aided detection following primary review.

Breast Ultrasound: Doppler and grayscale imaging of both breasts, complete: four quadrants, retroareolar

COMPARISON:

FINDINGS:

MAMMOGRAM:

Density: The breasts are heterogeneously dense, which may obscure small masses.

There is no . A was placed over the right superior periareolar region in the area of palpable concern. Additional imaging of this area with a spot tangential view demonstrates no

BILATERAL BREAST SONOGRAM:

Breast composition: Heterogeneous background echotexture.

Right breast: The entire breast was imaged.

Along 2 o'clock 2 cm from nipple in the area of palpable concern there is a hypoechoic finding that extends over 24 mm. This appears different from the surrounding tissue; ultrasound-guided biopsy is warranted.

Small inspissated cysts/fibrocystic changes were demonstrated along 12 o'clock.

Left breast: The entire breast was imaged.

No

Small cysts and inspissated cysts were demonstrated, as on prior exams.

Exam End: 16:44

Specimen Collected: 00:00

Last Resulted: 17:26

Received From: Weill Cornell Medicine, Columbia

Result Received: 10:25

Physicians, and New York Presbyterian Hospital

 View Encounter  Received Information