

Media Information

08/25/2025 09:34 AM

FROM:

Page: 3

SELECT
DIAGNOSTIC
IMAGING2350 OCEAN AVENUE, SUITE 4
NY 11229
TEL: (718) 676-4350 FAX: (718) 676-4357Farzam Kashanian, M.D.
MEDICAL DIRECTOR**Report Details**

Patient Name:

Patient

Date of Birth:

Gender:

F

Accession:

Referring Physician:

Report Text**EXAM: DIGITAL BILATERAL DIAGNOSTIC MAMMOGRAM**
EXAM: BILATERAL BREAST ULTRASOUND

History: Palpable lump in the upper outer right breast. No

Technique: Bilateral MLO and Craniocaudal mammographic views were performed using digital technique.

Findings: The breast parenchyma is heterogeneously dense, 50 to 75% fibroglandular, lowering the sensitivity of the study. Palpable nodule in the upper outer periareolar right breast was marked with a marker. Spot compression over this region demonstrates several ill defined nodular densities measuring 1.5 cm and smaller. These are irregular. Nodularity is also seen in the upper outer right breast. Within the left breast, mild nodularity is seen in the upper outer left breast. No

Sonographic evaluation of both breasts was performed using a high frequency linear transducer. All four quadrants, retroareolar regions and axillae were evaluated bilaterally.

The breast parenchyma is heterogeneously dense. Within the right breast, at the 10 o'clock retroareolar region, a lobulated irregular mass measures 1.6 x 1.4 x 1.8 cm corresponding to the palpable lump. An adjacent irregular nodule measures 0.7 x 0.7 x 0.6 cm. At the right 11 o'clock periareolar region, a lobulated nodule measures 1.0 x 1.0 x 0.8 cm. Right axillary lymph nodes are seen demonstrating thickened cortex. Within the left breast, an ill defined nodule at the 2 o'clock axis measures 0.7 x 0.6 x 0.6 cm. Benign appearing lymph nodes are seen in the left axilla.

IMPRESSION:

PALPABLE LUMP IN THE RIGHT RETROAREOLAR REGION CORRESPONDS TO 1.8 CM IRREGULAR SOLID MASS. AN ADJACENT 0.7 CM SOLID MASS IS ALSO SEEN AS WELL AS RIGHT 11 O'CLOCK 1.0 CM SOLID MASS. THESE ARE ALL SUSPICIOUS FOR CANCER. 0.7 CM ILL DEFINED NODULE IS ALSO SEEN AT THE LEFT BREAST 2 O'CLOCK POSITION THAT IS ALSO SUSPICIOUS. LYMPH NODES IN THE RIGHT AXILLA ALSO SEEN THAT ARE INDETERMINATE. RECOMMEND FURTHER EVALUATION WITH ULTRASOUND GUIDED CORE NEEDLE BIOPSY OF BOTH BREASTS. BILATERAL BREAST MRI WITH CONTRAST ENHANCEMENT IS ALSO RECOMMENDED.

BI-RADS CATEGORY 5(HIGHLY SUGGESTIVE OF MALIGNANCY-APPROPRIATE ACTION SHOULD BE TAKEN)

THE RESULTS WERE REPORTED TO RADA AT DR TRAKHTENBERG'S OFFICE ON

Please be advised that the false negative rate of mammography is approximately 10% and management of a palpable mass should be based on clinical grounds.

As per the FDA requirements, a layman's letter has been generated and sent to your patient results and recommendations of this breast imaging study. The patient system and well be notified when they are due for their next breast imaging exam.

FARZAM KASHANIAN MD

Signed by:

Kashanian, Farzam

Date Signed:

8/15/2025

Transcribed By:

TRANSCRIPTIONIST, EXCEL

Date Transcribed:

00:00

Attached To:

OUTSIDE RADIOLOGY ORDER

Orders Only on with Historical Provider,

Source Information

Megan Thomas | Msk Pas