

Media Information

9:46 AM

Fax Services

→

pg 16 of 34

10/6/25, 2:25 PM

Atlantic Medical Imaging - Synapse EIS



Accurate answers. Fast.

6529 Black Horse Pike, Egg Harbor Township, NJ 08234 • P: • F:

Name:
MRN: 930411641
DOB:
Acc #:
Sec Acc#:
Exam Date: 9/9/2025 12:52 PM

Address: 829 SHORE ROAD
Referrer Phone#: (609) 926-6353
Referrer Fax#:
cc:
cc:

BILATERAL BREAST MRI WITHOUT AND WITH CONTRAST WITH COMPUTER-AIDED DETECTION:
12:00 AM

INDICATION: 63 years old presenting for short interval follow-up for right upper outer breast enhancement and for correlation to prior mammographic finding on diagnostic mammogram dated

COMPARISON: 2/2/2024, and 11/19/2023. Correlation is made to prior ultrasound biopsy dated and diagnostic mammogram/breast ultrasound dated 8/4/2025.

PERSONAL HISTORY/RISK FACTORS: Benign right breast ultrasound-guided biopsy in 2025.

FAMILY HISTORY OF BREAST CANCER: No

RISK ASSESSMENT: The patient's lifetime breast cancer risk according to the Tyrer-Cuzick model is 7.7%. (Based on a revised TC model (version 8) which now incorporates breast density).

TECHNIQUE

Multisequence, multiplanar images of both breasts were obtained without and with contrast. Subtracted MIP images were performed using the CAD stream system. The patient received:

IV CONTRAST VIALS USED: Dotarem one 15cc vial.
IV CONTRAST INJECTED: 15 cc.
WASTED CONTRAST: 0 cc.

The study was reviewed on the PACS workstation and in conjunction with the data supplied from computer aided detection software, CADstream. Available pathology was reviewed.

FINDINGS

RIGHT BREAST:

10/6/25, 2:25 PM

Atlantic Medical Imaging - Synapse EIS

RATIO OF FIBROGLANDULAR TISSUE: Heterogeneously dense fibroglandular tissue.

BACKGROUND PARENCHYMAL ENHANCEMENT: Marked, which limits sensitivity of this study.

In the upper outer right breast, 8 cm from the nipple, there is an enhancing 7 mm mass (series 10 image 141, series 5 image 33, and CAD 36), likely correlate for previously identified mammographic asymmetry identified on study dated [REDACTED]. Stereotactic guided biopsy is recommended.

There are scattered complicated oil cysts some which demonstrate rim enhancement.

Visualized portion of the axilla is unremarkable. There is no [REDACTED] [REDACTED] [REDACTED] [REDACTED] [REDACTED]

LEFT BREAST:

RATIO OF FIBROGLANDULAR TISSUE: Heterogeneously dense fibroglandular tissue.

BACKGROUND PARENCHYMAL ENHANCEMENT: Marked, which limits sensitivity of this study.

There is no suspicious enhancement.

There are scattered complicated oil cysts some which demonstrate rim enhancement.

Visualized portion of the axilla is unremarkable. There is no [REDACTED] [REDACTED] [REDACTED] [REDACTED] [REDACTED]

IMPRESSION:

1. Right upper outer breast, 8 cm from the nipple, enhancing 7 mm mass, likely correlate for previously identified mammographic asymmetry identified on study dated 8/4/2025. Stereotactic guided biopsy is recommended.

2. No [REDACTED] [REDACTED] [REDACTED] [REDACTED] [REDACTED]

BI-RADS: 4: Suspicious

MANAGEMENT RECOMMENDATIONS: Right breast stereotactic guided biopsy is recommended in the absence of clinical contraindication. If stereotactic guided biopsy is unsuccessful, MRI guided biopsy would then be recommended.

If you are the referring physician [REDACTED] [REDACTED] [REDACTED] [REDACTED] [REDACTED] (609)650-0004 to reach a radiologist.

Date Of Signature: 9/10/2025 8:53 AM

Report Electronically Signed by: Kahn, D.O., Julia

Report Electronically Signed on: [REDACTED] 08:53 AM

Patient Name: [REDACTED] [REDACTED]

[REDACTED] [REDACTED] [REDACTED] [REDACTED] [REDACTED]

Patient [REDACTED] [REDACTED]

anirix/WebForms/Patient/PatientId [REDACTED]

[REDACTED]

Document Information

Misc Administration: Outside Radiology, Breast

MRI BREAST

00:00

Attached To:

OUTSIDE RADIOLOGY ORDER

Orders Only on with Historical Provider,

Source Information

Priscilla P. Hazel | Msk Pas