

May 05, 2025 10:52

To: [REDACTED]

Page: 1/2

From: Pathline [REDACTED]



LABORATORY MEDICINE ASSOCIATES, LLP
Montefiore Nyack Hospital Laboratory
160 North Midland Ave, Nyack NY [REDACTED]
Tel: 845-348-2770 Fax: 845-348-3430

Surgical Pathology Report

PATIENT

Patterson, Nadia L

185 First Street

Apt 1

Newburgh, NY [REDACTED]

DOB: [REDACTED]

AGE: 46 SEX: Female

Patient [REDACTED]

ST25-010494

COLLECTED: [REDACTED]

RECEIVED: [REDACTED]

REPORTED: [REDACTED]

Steven Reichard, DO

Hudson Valley Radiology Associates
of Westchester

505 Route 208

Monroe, NY [REDACTED]

CC: Anderson, Melissa

Final Diagnosis

A. Left breast, medial calcifications, stereotactic biopsy, suros assisted:

- Benign breast tissue.

B. Left breast, medial tissue, stereotactic biopsy, suros assisted:

- Benign breast tissue with a rare calcification within lobules.

- Focal fibrocystic changes.

C. Left breast, lateral calcifications, stereotactic biopsy, suros assisted:

-Ductal carcinoma in situ, intermediate grade, cribriform type, with associated calcifications.

-Radial sclerosing lesion.

-Fibrocystic changes with calcifications.

Breast Cancer Biomarker Results (Block C):

Estrogen Receptor (ER): Positive(98%, strong)

Progesterone Receptor (PR): Positive(100%, strong)

Status of internal controls (for ER and PR Immunohistochemistry):

☒ Internal control cells present and stain as expected.

Methods: Immunohistochemistry (IHC) for ER and PR was performed at Pathline - Histopathology Services, Ramsey, NJ, and interpreted at Montefiore Nyack Hospital, Nyack, NY. ER and PR IHC are laboratory-developed tests. Cold ischemia and fixation times meet requirements specified in the latest version of the ASCO/CAP guidelines. The fixative used is 10% buffered formalin. The following primary antibodies were used on the Leica Bond III system: ER (SP1); PR (312).

D. Left breast, lateral tissue, stereotactic biopsy, suros assisted:

-Ductal carcinoma in situ, intermediate grade. cribriform type,

-Radial sclerosing lesion.

-Fibrocystic changes with calcifications.

Electronically Signed By:

Vandana Hoon, MD On [REDACTED] 01:46 PM EST

ACCESSION [REDACTED]

Clinical History:

Left breast stereotactic biopsy, suros assisted, 2 sites.

Gross Description:

- A) Received in formalin labeled with patient's name, [REDACTED] single tan-yellow to white cylindrical fragment of fibrofatty soft tissue core measuring 2.7 cm in length by 0.2 to 0.3 cm in diameter. Entirely submitted in one cassette.
- B) Received in formalin labeled with patient's name, [REDACTED] tan-yellow to white irregular to cylindrical fragments of fibrofatty soft tissue cores measuring and 2.7 x 2.6 x 0.2 cm in aggregate. Entirely submitted in two cassettes.
- C) Received in formalin labeled with patient's name, [REDACTED] tan-yellow to white irregular to cylindrical fragments of fibrofatty soft tissue cores measuring 2.8 x 1.5 x 0.2 cm in aggregate. Entirely submitted in one cassette.
- D) Received in formalin labeled with patient's name, [REDACTED] tan-yellow to white irregular to cylindrical fragments of fibrofatty soft tissue cores measuring 1.8 x 1.5 x 0.2 cm in aggregate. Entirely submitted in one cassette.

Time to fixation: Presumed Immediate;

Time of fixation: Approximately 13-14 hours.

Previous Cases:

None



LABORATORY MEDICINE ASSOCIATES, LLP
Montefiore Nyack Hospital Laboratory
160 North Midland Ave, Nyack NY [REDACTED]
Tel: 845 348 2270 Fax: 845 348 8430

Surgical Pathology Report

PATIENT

Patterson, Nadia L

185 First Street

Apt 1

Newburgh, NY [REDACTED]

DOB: [REDACTED]

AGE: 46 SEX: Female

Patient [REDACTED]

ST25-010494

COLLECTED: [REDACTED]

RECEIVED: [REDACTED]

REPORTED: [REDACTED]

Steven Reichard, DO

Hudson Valley Radiology Associates

of Westchester

505 Route 208

Monroe, NY 10950

CC: Anderson, Melissa

Final Diagnosis

A. Left breast, medial calcifications, stereotactic biopsy, suros assisted:

- Benign breast tissue.

B. Left breast, medial tissue, stereotactic biopsy, suros assisted:

- Benign breast tissue with a rare calcification within lobules.

- Focal fibrocystic changes.

C. Left breast, lateral calcifications, stereotactic biopsy, suros assisted:

-Ductal carcinoma in situ, intermediate grade, cribriform type, with associated calcifications.

-Radial sclerosing lesion.

-Fibrocystic changes with calcifications.

Breast Cancer Biomarker Results (Block C):

Estrogen Receptor (ER): Positive(98%, strong)

Progesterone Receptor (PR): Positive(100%, strong)

Status of internal controls (for ER and PR immunohistochemistry):

☒ Internal control cells present and stain as expected.

Methods: Immunohistochemistry (IHC) for ER and PR was performed at Pathline - Histopathology Services, Ramsey, NJ, and interpreted at Montefiore Nyack Hospital, Nyack, NY. ER and PR IHC are laboratory-developed tests. Cold Ischemia and fixation times meet requirements specified in the latest version of the ASCO/CAP guidelines. The fixative used is 10% buffered formalin. The following primary antibodies were used on the Leica Bond III system: ER (SP1); PR (312).

D. Left breast, lateral tissue, stereotactic biopsy, suros assisted:

-Ductal carcinoma in situ, intermediate grade. cribriform type,

-Radial sclerosing lesion.

-Fibrocystic changes with calcifications.

Electronically Signed By:

Vandana Hoon

Vandana Hoon, MD On [REDACTED] 25 01:46 PM EST

Patient: [REDACTED] 2 Sites [MABX05] - Breast - L] Scheduled:

ACCESSION [REDACTED]

Clinical History:

Left breast stereotactic biopsy, suros assisted, 2 sites.

Gross Description:

- A) Received in formalin labeled with patient's name, [REDACTED] single tan-yellow to white cylindrical fragment of fibrofatty soft tissue core measuring 2.7 cm in length by 0.2 to 0.3 cm in diameter. Entirely submitted in one cassette.
- B) Received in formalin labeled with patient's name, [REDACTED] tan-yellow to white irregular to cylindrical fragments of fibrofatty soft tissue cores measuring and 2.7 x 2.6 x 0.2 cm in aggregate. Entirely submitted in two cassettes.
- C) Received in formalin labeled with patient's name, [REDACTED] tan-yellow to white irregular to cylindrical fragments of fibrofatty soft tissue cores measuring 2.8 x 1.5 x 0.2 cm in aggregate. Entirely submitted in one cassette.
- D) Received in formalin labeled with patient's name, [REDACTED] tan-yellow to white irregular to cylindrical fragments of fibrofatty soft tissue cores measuring 1.8 x 1.5 x 0.2 cm in aggregate. Entirely submitted in one cassette.

Time to fixation: Presumed Immediate;

Time of fixation: Approximately 13-14 hours.

Previous Cases:

None

Technical component performed at Pathline LLC, 535 E. Crescent Ave., Ramsey, NJ [REDACTED]

Patterson, Nadia L
ST25-010494

Page 2 of 2