

Module 5. Bivariate analysis: qualitative - quantitative

Data Science & AI

Sabine De Vreese Stijn Lievens Lieven Smits Bert Van Vreckem
2021-2022

**HO
GENT**

Contents

Data visualization

Two-sample t-test

Paired samples

Effect size

**HO
GENT**

Learning goals

- Apply the t -test for two samples
- Calculate effect size using Cohen's d
- Visualization

Bivariate analysis: overview

Independent	Dependent	Test/Metric
Qualitative	Qualitative	χ^2 -test Cramér's V
Qualitative	Quantitative	two-sample t -test Cohen's d
Quantitative	Quantitative	— Regression, correlation

Example research questions

- Are male penguins larger than females?
- Do men get a higher salary than women?
- Does a new vaccine protect against a disease?
- Does “retrieval practice” improve learning outcomes (i.e. student grades)?
- ...

In these examples, what is the independent/dependent variable?

Data visualization

**HO
GENT**

Data visualization

- Chart types for quantitative data
- Grouped by qualitative variable

Data visualization


- Chart types for quantitative data
- Grouped by qualitative variable

Suitable chart types:

- Grouped boxplot
- Grouped density plot
- Bar chart with error bars
- ...


Grouped boxplot

(source code: see `demo-2sample-t.ipynb`)




**HO
GENT**

Grouped violin plot




**HO
GENT**

Grouped density plot



**HO
GENT**

Beware! Bar chart of group means



**HO
GENT**


Beware! Bar chart of group means



**HO
GENT**

Bar chart with error bars

- Always add error bars!
- Only makes sense for normally distributed data
- Example: 1 standard deviation:



**HO
GENT**

Two-sample t-test

**HO
GENT**

Comparing two samples

Is the sample mean of two samples significantly different?

- Independent samples
- Paired samples

Independent samples

In a clinical study, the aim is to determine whether a new drug has a delayed (i.e. higher) reaction speed as a side effect

- Control group: 6 participants receive placebo
- Intervention group: 6 participants receive medicine

Independent samples

In a clinical study, the aim is to determine whether a new drug has a delayed (i.e. higher) reaction speed as a side effect

- Control group: 6 participants receive placebo
- Intervention group: 6 participants receive medicine

Next, the reaction speed is measured:

- Control group: 91, 87, 99, 77, 88, 91 ($\bar{x} = 88.83$)
- Intervention group: 101, 110, 103, 93, 99, 104 ($\bar{y} = 101.67$)

Are there significant differences between the intervention and control group?

HO
GENT

Independent samples

Testing procedure

1. Hypotheses:
 - $H_0: \mu_1 - \mu_2 = 0$
 - $H_1: \mu_1 - \mu_2 < 0$
2. Significance level: $\alpha = 0.05$
3. Test statistic:
 - $\bar{x} - \bar{y} = -12.833$
 - \bar{x} = estimation for μ_1 (control group)
 - \bar{y} = estimation for μ_2 (intervention group)
4. Calculate p

Independent samples

Calculation in Python

```
control = np.array([91, 87, 99, 77, 88, 91])
treatment = np.array([101, 110, 103, 93, 99, 104])
stats.ttest_ind(a=control, b=treatment,
                 alternative='less', equal_var=False)
```

Result:

```
Ttest_indResult(statistic=-3.445612673536487,
                 pvalue=0.003391230079206901)
```

**HO
GENT**

Independent samples

Testing procedure (continued)

5. $p \approx 0.00339 < \alpha = 0.05$. We reject the null hypothesis. In this sample, there is reason to assume that the drug does indeed have an impact on reaction speed.

**HO
GENT**

Paired samples

**HO
GENT**

Paired samples

A study examined whether cars that run on petrol with additives have a lower consumption.

For 10 cars, the consumption was measured (expressed in miles per gallon) for both fuel types:

Car	1	2	3	4	5	6	7	8	9	10
Regular petrol	16	20	21	22	23	22	27	25	27	28
With additives	19	22	24	24	25	25	25	26	28	32

Paired samples

Testing procedure

1. Hypotheses:
 - $H_0: \overline{x - y} = 0$
 - $H_1: \overline{x - y} > 0$
2. Significance level: $\alpha = 0.05$
3. Test statistic:
 - $\overline{x - y}$
 - x = miles per gallon with additives ($\bar{x} = 25.1$)
 - y = miles per gallon regular petrol ($\bar{y} = 23.1$)
4. Calculate p

**HO
GENT**

Paired samples

Calculation in Python

```
regular = np.array([16, 20, 21, 22, 23, 22, 27, 25, 27, 28])  
additives = np.array([19, 22, 24, 24, 25, 25, 26, 26, 28, 32])  
stats.ttest_rel(regular, additives, alternative='less')
```

Result:

```
Ttest_relResult(statistic=-4.47213595499958,  
                 pvalue=0.00077494295585091)
```

**HO
GENT**

Paired samples

5. $p \approx 0.0007749 < \alpha = 0.05$. We reject the null hypothesis. In this sample, there is reason to assume that the fuel with additives leads to lower fuel consumption.

Effect size

**HO
GENT**

Effect size

Effect size

The **effect size** is a metric which expresses how great the difference between two groups is

- Control group vs. intervention group
- Can be used in addition to hypothesis test
- Often used in educational sciences
- There are several definitions, here: *Cohen's d*

Cohen's d

$$d = \frac{\overline{x}_1 - \overline{x}_2}{s}$$

with $\overline{x}_1, \overline{x}_2$ the sample means
and s a standard deviation of both groups combined:

$$s = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}}$$

with n_1, n_2 the sample sizes, s_1, s_2 the standard deviation of both groups

**HO
GENT**

Interpretation Cohen's d

$ d $	Effect Size
0.01	Very small
0.2	Small
0.5	Average
0.8	Large
1.2	Very large
2.0	Huge

In educational sciences (John Hattie):

- 0,4 = tipping point for desired effects
- effect size $d = 1$: process material from 1y in 6m!

E.g. <https://visible-learning.org/backup-hattie-ranking-256-effects-2017/>

**HO
GENT**

Typical approach research in education

- Research question: Is X a good learning strategy, in other words, does this have a positive effect on final results?
- Control group uses “traditional” approach
- Intervention group uses X
- Followed by an evaluation moment
- Determine scores, calculate d