

AI

AI

Search Problem

Uninformed search strategies

Breadth-first search

Depth-first search

Iterative deepening search

Heuristic Search

Greedy best-first search

A* search

Admissible heuristics

Constraint satisfaction problem (CSP)

Which variable should be assigned next?

***Most constrained variable** (选之后的点用这个)

Most constraining variable (如果是选第一个点，用这个)

Least constraining value (选涂什么颜色)

Logic

Propositional Logic

language

Axiomatisation

Proof

Valuation

Semantic consequence

Equivalent Formulae

Conjunctive Normal Form

Resolution

Deduction Theorem

Probabilistic

Boolean algebras of set

Finitely Additive Probability Measures

Conditional Probability

Independent

Bayes' Rule

Search Problem

The trade-off usually between:

- Speed (Time complexity)
- Memory (Space complexity)
- 'Optimality' (Optimal)

Four characteristics:

- Time complexity: number of nodes generated
- Space complexity: maximum number of nodes in memory
- Optimality: does it always find a least-cost solution?
- completeness: does it always find a solution if one exists?

b: maximum branching factor of the search tree

d: depth of the least-cost solution

m: maximum depth of the state space (may be ∞)

Uninformed search strategies

Breadth-first search

- Complete: Yes
- Time: $O(b^{d+1})$
- Space: $O(b^{d+1})$
- Optimal: Yes
- **Space** is the bigger problem (More than time)

Depth-first search

- Complete: No
- Time: $O(b^m)$, terrible if m is much larger than d
- Space: $O(bm)$, linear space
- Optimal: No

Iterative deepening search

- Complete: Yes
- Time: $O(b^d)$
- Space: $O(bd)$ linear space
- Optimal: Yes

Heuristic Search

Best first search:

Idea: Use an **evaluation function** $f(n)$ for each node

special case: Greedy Best First Search, A* Search

Greedy best-first search

- Evaluation function $f(n) = h(n)$ (**heuristic**) = estimate of cost from n to goal
- Complete: No
- Time: $O(b^m)$
- Space: $O(b^m)$
- Optimal: No


A* search

- Idea: avoid expending paths that are already expensive
- Evaluation function $f(n) = g(n) + h(n)$
 - $g(n)$: cost so far (to reach n)
 - $h(n)$: estimated cost from n to goal
 - $f(n)$: estimated total cost of path through n to goal
- **Admissible heuristics** for A*:

- A heuristic $h(n)$ is **admissible** if for every node n , $h(n) \leq h^*(n)$, where $h^*(n)$ is the **true** cost to reach the goal state from n .
- An admissible heuristic never overestimates the cost to reach the goal
- **Theorem:** If $h(n)$ is admissible, A* using TREE-SEARCH is optimal
- Complete: Yes
- Time: Exponential (in length of optimal solution)
- Space: Keeps all (expanded) nodes in memory
- Optimal: Yes

Optimality of A* (proof)


- ▶ Suppose some suboptimal goal state G_2 has been generated and is in the fringe. Let n be an unexpanded node in the fringe such that n is on a shortest path to an optimal goal G with true cost C^* .



Admissible heuristics

E.g., for the 8-puzzle:

- ▶ $h_1(n)$ = number of misplaced tiles
- ▶ $h_2(n)$ = total Manhattan distance
(i.e., no. of squares from desired location of each tile)



▶ $h_1(S) = ?$ 8

▶ $h_2(S) = ?$ $3 + 1 + 2 + 2 + 2 + 3 + 3 + 2 = 18$

Dominance

- ▶ If $h_2(n) \geq h_1(n)$ for all n (both admissible)
- ▶ then h_2 **dominates** h_1
- ▶ h_2 is better for search
- ▶ Typical search costs (average number of nodes expanded = effective branching factor):
 - ▶ $d=12$ IDS = 3,644,035 nodes
 $A^*(h_1) = 227$ nodes
 $A^*(h_2) = 73$ nodes
 - ▶ $d=24$ IDS = too many nodes
 $A^*(h_1) = 39,135$ nodes
 $A^*(h_2) = 1,641$ nodes

Constraint satisfaction problem (CSP)

- **state** is defined by **variables** X_i with **values** from **domain** D_i
- **goal test** is a set of **constraints** specifying allowable combinations of values for subsets of variables

Which variable should be assigned next?

*Most constrained variable (选之后的点用这个)

- choose the variable with the fewest legal values,
- as known as minimum remaining values (MRV) heuristic
- Because these are the variables that are most likely to prune the search tree

Most constraining variable (如果是选第一个点, 用这个)

- Most constraining variable = the variable with the most constraints on remaining variables
- Because the variable involved in the most constraints, so it is most likely to cause prune the search tree

Least constraining value (选涂什么颜色)

- Given a variable, choose the least constraining value:
 - the one that rules out the fewest values in the neighbouring variables – to leave other variables as open as possible

Summary

- ▶ CSPs are a special kind of problem:
 - states defined by values of a fixed set of variables
 - goal test defined by constraints on variable values
 -
- ▶ Backtracking = depth-first search with one variable assigned per node
- ▶ Variable ordering and value selection heuristics help significantly
- ▶
- ▶ Forward checking prevents assignments that guarantee later failure
- ▶
- ▶ Constraint propagation (e.g., arc consistency) does additional work to constrain values and detect inconsistencies
- ▶
- ▶ Iterative min-conflicts is usually effective in practice
- ▶

Logic

(Not important)

The central component of a knowledge-based agent is their **knowledge base**, or KB.

The agent uses logical reasoning to make decisions.

The agent draws conclusions from the available information.

Propositional Logic

(Just know the concept)

language

To define a knowledge base and define sentences, we need a language

The language of propositional logic is built from

true			
false			
\wedge	and	conjunction	(& or .)
\vee	or	disjunction	(or +)
\neg	not	negation	(~)
\Rightarrow	if . . . then	implication	(\rightarrow)
\Leftrightarrow	if and only if	equivalence	(\leftrightarrow)

Formulas of propositional logic are defined as follows:

- Each propositional variable P, Q, R, \dots is a formula.
- **T** and **F** are formulas
- If ϕ is a formula, $\neg\phi$ is a formula
- If ϕ and ψ are formulas so are

$$\phi \wedge \psi, \quad \phi \vee \psi, \quad \phi \Rightarrow \psi, \quad \phi \Leftrightarrow \psi.$$

Propositional variables and constants are called **atomic formulas**. The remaining cases are called **compound (or complex) formulas**.

Axiomatisation

(Important)

Propositional logic is defined by a set of axioms and rules.

Propositional logic has the following axiomatisation:

- $A \Rightarrow (B \Rightarrow A)$
- $(A \Rightarrow (B \Rightarrow C)) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C))$
- $(\neg A \Rightarrow \neg B) \Rightarrow (B \Rightarrow A)$
- From A and $A \Rightarrow B$, infer B (**Modus Ponens**)
 - which means if A and $A \Rightarrow B$ are **true**, then we could infer B is also **true**
 -

$$\frac{A \Rightarrow B, A}{B}$$

Or, $A \Rightarrow B, A \vdash B$

If the antecedent of an implication (a conditional claim) is true, then the consequent must also be true.

Modus Ponens Example

A \Rightarrow B: "If today is Tuesday, then I will go to AI lecture."

A: "Today is Tuesday."

B: "I will go to AI lecture."

A \Rightarrow B: "If it is raining, then there are clouds in the sky."

A: "It is raining."

B: "There are clouds in the sky."

A \Rightarrow B: "If Yankees win today's game, they will be champions."

A: "Yankees win today's game."

B: Yankees are champions.

A \Rightarrow B: "If the weather is good, we can go to the beach."

A: The weather is good.

B: We can go to the beach.

Proof

Given a KB, a **proof** is a finite sequence of formulas

$$\phi_1, \dots, \phi_n$$

where each ϕ_i is

- a formula from KB, or
- an instance of an axiom (A1-A3), or
- derived from ϕ_j, ϕ_k , with $j, k < i$ by using modus ponens.

Given a KB and a formula ϕ , we say that ϕ is a **logical consequence** of KB, denoted

$$KB \vdash \phi$$

if there exists a proof of ϕ from KB.

Example: how to prove that

$$\phi \Rightarrow \phi$$

is a theorem.

① $(\phi \Rightarrow ((\psi \Rightarrow \phi) \Rightarrow \phi)) \Rightarrow ((\phi \Rightarrow (\psi \Rightarrow \phi)) \Rightarrow (\phi \Rightarrow \phi))$

Instance of A2

② $\phi \Rightarrow ((\psi \Rightarrow \phi) \Rightarrow \phi)$

Instance of A1

③ $(\phi \Rightarrow (\psi \Rightarrow \phi)) \Rightarrow (\phi \Rightarrow \phi)$

From (1) and (2) with modus ponens

④ $\phi \Rightarrow (\psi \Rightarrow \phi)$

Instance of A1

⑤ $\phi \Rightarrow \phi$

From (3) and (4) with modus ponens.

Example

From $p \Rightarrow q$, $(\neg r \vee q) \Rightarrow (s \vee p)$, q , prove $s \vee q$.

1. $p \Rightarrow q$	[Given]
2. $(\neg r \vee q) \Rightarrow (s \vee p)$	[Given]
3. q	[Given]
4. $s \vee q$	[3, \vee introduction]

Exercise

Show r from $p \Rightarrow (q \Rightarrow r)$ and $p \wedge q$ using the rules we have been given so far. That is, prove

$$p \Rightarrow (q \Rightarrow r), p \wedge q \vdash r.$$

1. $p \Rightarrow (q \Rightarrow r)$	[Given]
2. $p \wedge q$	[Given]
3. q	[2, \wedge -Introduction]
4. p	[2, \wedge -Introduction]
5. $q \Rightarrow r$	[1,4, modus ponens]
6. r	[3, modus ponens]

Let *Prop.* be a set of propositional variables. A valuation v is a mapping :

$$v : Prop \rightarrow \{0, 1\}$$

A valuation assigns to each propositional variable either 1 (true) or 0 (false)

Negations:

$$v(\neg\phi) = \begin{cases} 1 & v(\phi) = 0 \\ 0 & \text{otherwise} \end{cases}$$

$\neg\phi$ is true IFF ϕ is false.

negation \neg

p	$\neg p$
T	F
F	T

Conjunctions:

$$v(\phi \wedge \psi) = \begin{cases} 1 & v(\phi) = v(\psi) = 1 \\ 0 & \text{otherwise} \end{cases}$$

$\phi \wedge \psi$ is true IFF both ϕ and ψ are true

Conjunction \wedge

p	q	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

Disjunction:

$$v(\phi \vee \psi) = \begin{cases} 0 & v(\phi) = v(\psi) = 0 \\ 1 & \text{otherwise} \end{cases}$$

$\phi \vee \psi$ is true IFF at least one formula in $\{\phi, \psi\}$ is true.

Disjunction \vee

p	q	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

Implications:

$$v(\phi \Rightarrow \psi) = \begin{cases} 0 & v(\phi) = 1, v(\psi) = 0 \\ 1 & \text{otherwise} \end{cases}$$

$\phi \Rightarrow \psi$ is true IFF either ϕ is false or both ϕ and ψ are true.

Implication \Rightarrow (If...then...)

p	q	$p \Rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

$p \Rightarrow q$:

converse: $q \Rightarrow p$

contrapositive: $\neg q \Rightarrow \neg p$

inverse: $\neg p \Rightarrow \neg q$

Double Implication:

$$v(\phi \Leftrightarrow \psi) = \begin{cases} 1 & v(\phi) = v(\psi) \\ 0 & \text{otherwise} \end{cases}$$

$\phi \Leftrightarrow \psi$ is true IFF ϕ and ψ have the same truth-value.

Double Implication \Leftrightarrow (if and only if...)

p	q	$p \Leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

Truth Table

A truth-table for a formula is just a representation of all the possible assignments of values and the corresponding outputs.

ϕ	ψ	$\neg\phi$	$\phi \wedge \psi$	$\phi \vee \psi$	$\phi \Rightarrow \psi$	$\phi \Leftrightarrow \psi$
1	1	0	1	1	1	1
1	0	0	0	1	0	0
0	1	1	0	1	1	0
0	0	1	0	0	1	1

Example

p
(my breakfast is) eggs.
q
(my breakfast is) cereal.
r
(my breakfast is) toast.

p	q	r	$p \vee q$	$r \wedge (p \vee q)$
T	T	T	T	T
T	T	F	T	F
T	F	T	T	T
T	F	F	T	F
F	T	T	T	T
F	T	F	T	F
F	F	T	F	F
F	F	F	F	F

The statement 'my breakfast is toast and either eggs or cereal' in symbolic form as $r \wedge (p \vee q)$

Valuation

We can check whether a formula is true under a given valuation by looking at its truth-table.

- A formula ϕ is called **satisfiable** if there exists a valuation v such that $v(\phi) = 1$.
 - "a sentence is satisfiable if it is **True in some** model"
- A formula ϕ is called a **tautology** if for all possible valuations v , $v(\phi) = 1$.
 - a formula is said to be tautology if and only if it is **true under every interpretation**.
- A formula ϕ is called a **contradiction** if for all possible valuations v , $v(\phi) = 0$.
 - a sentence is contradiction if it is **false in all** models
- Two formulas are **logically equivalent** if they have the **same truth-table**.
- Given a knowledge base KB, we denote by $Mod(KB)$, the set of all valuations that make all the formulas in KB true.

Examples:

The following truth-table shows that $\phi \Rightarrow \psi$ and $\neg\phi \vee \psi$ are logically equivalent:

ϕ	ψ	$\neg\phi$	$\phi \Rightarrow \psi$	$\neg\phi \vee \psi$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

The following truth-table shows that axiom A1 is a tautology: (easy could prove axioms A2 and A3 are also tautologies)

ϕ	ψ	$\psi \Rightarrow \phi$	$\phi \Rightarrow (\psi \Rightarrow \phi)$
1	1	1	1
1	0	1	1
0	1	0	1
0	0	1	1

Semantic consequence

Given a knowledge base KB and a formula ϕ , we say that ϕ is **semantic consequence** of KB, or that KB **entails** ϕ denoted

$$KB \models \phi$$

if every valuation (model) that makes all the formulas in KB true also makes ϕ true.

Which means:

$$Mod(KB) \subseteq Mod(\phi)$$

the set of models that make all the formulas in KB true is a subset of the set of valuations that make ϕ true.

\vdash is about deductive inference and proof. It is a syntactical notion.

\models is about relationship between models. It is a semantical notion.

Propositional logic is **sound**:

- If $KB \vdash \phi$ then $KB \models \phi$
- This means that logical inference preserves truth, i.e. from true premises we can only derive true conclusions.

Propositional logic is also **complete**:

- If $KB \models \phi$ then $KB \vdash \phi$
- Everything we prove from true premises is true, and every truth can be proven.

The simple algorithm enumerates all the models of KB and ϕ and checks if $Mod(KB) \subseteq Mod(\phi)$.

If KB and ϕ are built from n propositional variables, this means we have to check **2^n models**.

Equivalent Formulae

Examples of inference rules and theorems (which can be derived from A1-A3 and modus ponens)

And-elimination:

From $\phi \wedge \psi$, derive ϕ .

Double negation:

$$\neg\neg\phi \Leftrightarrow \phi$$

De Morgan's Laws:

$$\neg(\phi \wedge \psi) \Leftrightarrow (\neg\phi \vee \neg\psi)$$

$$\neg(\phi \vee \psi) \Leftrightarrow (\neg\phi \wedge \neg\psi)$$

Distributivity Laws:

$$(\phi \wedge (\psi \vee \gamma)) \Leftrightarrow ((\phi \wedge \psi) \vee (\phi \wedge \gamma))$$

$$(\phi \vee (\psi \wedge \gamma)) \Leftrightarrow ((\phi \vee \psi) \wedge (\phi \vee \gamma))$$

Two formulae A and B are equivalent, written $A \equiv B$ if and only if A and B have the **same truth values** for **every interpretation**. (A 为 true, B 也为 true; A 为 false, B 也为 false) (这里的 \equiv 和课件里面的 \Leftrightarrow 同义)

\wedge - elimination

$\frac{A \wedge B}{A}$	or	$\frac{A \wedge B}{B}$	Or,	$\frac{A \wedge B \vdash B}{A \wedge B \vdash A}$
------------------------	----	------------------------	-----	---

From a conjunction you can infer any of the conjuncts

\wedge - introduction

$\frac{A, B}{A \wedge B}$	$A, B \vdash A \wedge B$
---------------------------	--------------------------

If A holds (true), and B holds, then $A \wedge B$ must also hold.

\vee - introduction

$\frac{A}{A \vee B}$	or	$\frac{A}{B \vee A}$	Or	$A \vdash A \vee B$
----------------------	----	----------------------	----	---------------------

If A holds (or are provable or true) then $A \vee B$ must hold.

$$\frac{A \vee B, \neg B \vee C}{A \vee C}$$

It says that if you know “A or B”, and you know “not be B or C”, then you're allowed to conclude “A or C”.

Contraposition

$$(A \Rightarrow B = \neg B \Rightarrow \neg A)$$

Associative laws

$$(A \wedge B) \wedge C \equiv A \wedge (B \wedge C)$$

$$(A \vee B) \vee C \equiv A \vee (B \vee C)$$

Commutative laws

$$A \wedge B \equiv B \wedge A$$

$$A \vee B \equiv B \vee A$$

Distributive laws

$$A \wedge (B \vee C) \equiv (A \wedge B) \vee (A \wedge C)$$

$$A \vee (B \wedge C) \equiv (A \vee B) \wedge (A \vee C)$$

Complement laws

$$A \wedge \neg A \equiv F$$

$$A \vee \neg A \equiv T$$

$$\neg(\neg A) \equiv A$$

de Morgan's laws

$$\neg(A \wedge B) \equiv \neg A \vee \neg B$$

$$\neg(A \vee B) \equiv \neg A \wedge \neg B$$

laws for \Rightarrow and \Leftrightarrow :

$$A \Rightarrow B \equiv \neg A \vee B$$

$$A \Leftrightarrow B \equiv (A \Rightarrow B) \wedge (B \Rightarrow A)$$

Example

Prove the following equivalence

$$\neg(\neg(P \wedge Q) \vee P) \equiv F$$

$$\neg(A \wedge B) \equiv \neg A \vee \neg B$$

$\neg(\neg(P \wedge Q) \vee P)$	[given]
$\equiv \neg((\neg P \vee \neg Q) \vee P)$	[de Morgan's laws]
$\equiv \neg((\neg Q \vee \neg P) \vee P)$	[Commutative laws]
$\equiv \neg(\neg Q \vee (\neg P \vee P))$	[Associative laws]
$\equiv \neg(\neg Q \vee T)$	[Complement laws]
$\equiv \neg T$	[Complement laws]
$\equiv F$	[Complement laws]

(课件中的example)

Example:

R1: $\neg P_{1,1}$, R2: $B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})$, R3: $B_{2,1} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{3,1})$, R4: $\neg B_{1,1}$, R5: $B_{2,1}$

KB = {R1,R2,R3,R4,R5}

prove $KB \models \neg P_{1,2}$, which also means prove $KB \models \neg \neg P_{1,2}$

We prove that $\neg P_{1,2}$ is a consequence of KB syntactically.

- 1 $((B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow (B_{1,1})))$
Equivalent to $((B_{1,1}) \Leftrightarrow (P_{1,2} \vee P_{2,1}))$
- 2 $((P_{1,2} \vee P_{2,1}) \Rightarrow (B_{1,1}))$
From (1) by And-elimination
- 3 $(\neg(B_{1,1}) \Rightarrow \neg(P_{1,2} \vee P_{2,1}))$
From (2) and by modus ponens, A3 and double negation.
- 4 $\neg(P_{1,2} \vee P_{2,1})$
From $\neg B_{1,1}$ and modus ponens.
- 5 $\neg P_{1,2} \wedge \neg P_{2,1}$
From (4), by De Morgan's laws.
- 6 $\neg P_{1,2}$
by And-elimination

Conjunctive Normal Form

A formula is in **Conjunctive Normal Form** if it is of the form

$$A_1 \wedge A_2 \wedge \dots \wedge A_k$$

where each A_i is a **disjunction of propositions or their negations**.

Example

$(p \vee q) \wedge r \wedge (\neg p \vee \neg r \vee s)$ is in CNF.

$\neg(p \vee q) \wedge r \wedge (\neg p \vee \neg r \vee s)$ is not in CNF.

$(p \vee q) \wedge r \wedge (p \Rightarrow (\neg r \vee s))$ is not in CNF.

Translation into CNF

To translate into CNF, first translate into NNF. Apply **distribution laws** or **commutativity laws** until in the correct form.

Example

$$A \Rightarrow B \equiv \neg A \vee B$$

Translate $(\neg(p \vee \neg q) \vee r) \Rightarrow p$ into CNF.

We first translate into NNF and obtain

$$((p \vee \neg q) \wedge \neg r) \vee p.$$

$$\neg(A \vee B) \equiv \neg A \wedge \neg B$$

$$((p \vee \neg q) \wedge \neg r) \vee p$$

$$((p \vee \neg q) \vee p) \wedge (\neg r \vee p)$$

$$\neg(A \wedge B) \equiv \neg A \vee \neg B$$

$$(p \vee \neg q \vee p) \wedge (\neg r \vee p)$$

Note the latter could be transformed into

$$(p \vee \neg q) \wedge (\neg r \vee p)$$

We show how to rewrite $B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})$ in CNF:

$$\textcircled{1} (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1})$$

By definition of \Leftrightarrow

$$\textcircled{2} (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg(P_{1,2} \vee P_{2,1}) \vee B_{1,1})$$

By definition of \vee

$$\textcircled{3} (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge ((\neg P_{1,2} \wedge \neg P_{2,1}) \vee B_{1,1})$$

By De Morgan laws and double negation

$$\textcircled{4} (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (B_{1,1} \vee \neg P_{1,2}) \wedge (B_{1,1} \vee \neg P_{2,1})$$

By distributivity

Resolution

Resolution is a proof method for classical **propositional** and **first-order logic**.

Given a formula ϕ , resolution will decide whether the formula is **contradiction** or not.

Propositional resolution works only on expressions in **clausal form**

A **literal** is either an atomic sentence or a negation of an atomic sentence, such as p or $\neg p$

A **clausal sentence** is either a literal or a disjunction of literals.

- p
- $\neg p$
- $p \vee q$

Resolution method involves:

- translation to a normal form (CNF);
- At each step, a new clause is derived from two clauses you already have
- Proof steps all use the same rule **resolution rule**
- repeat until false is derived or no new formulae can be derived.

resolution rule

If A or p is true, and B is true or p is false, Then either A or B is true.

$A \vee p$
$B \vee \neg p$
<hr/>
$A \vee B$

$A \vee B$ is called the **resolvent**

$A \vee p$ and $B \vee \neg p$ are called **parents of the resolvent**.

p and $\neg p$ are called complementary literals.

(Note in the above A or B can be empty)

The resolution algorithm shows that

$$KB \models \phi$$

by showing that $(KB \wedge \neg \phi)$ is a contradiction and so is unsatisfiable, i.e. there are no models that make $(KB \wedge \neg \phi)$ true.

The algorithm starts by converting $(KB \wedge \neg \phi)$ into a CNF.

Then, the resolution rule is applied to the resulting clauses.

Example: $\{\}$ means a contradiction

Given

$$P \vee Q$$

$$P \Rightarrow R$$

$$Q \Rightarrow R$$

Prove R

1. $P \vee Q$

2. $\neg P \vee R$

3. $\neg Q \vee R$

4. $\neg R$

negated conclusion

5. $Q \vee R$

1, 2

6. $\neg P$

2, 4

7. $\neg Q$

3, 4

8. R

5, 7

9. $\{\}$

4, 8

Using resolution rule, prove $\neg(p \Rightarrow (q \Rightarrow p))$

Step 1: translate to CNF:

$$\neg(\neg p \vee (\neg q \vee p))$$

I

$$\neg\neg p \wedge \neg(\neg q \vee p)$$

N

$$p \wedge (\neg\neg q \wedge \neg p)$$

D

$$p \wedge q \wedge \neg p$$

$$\{p\}$$

Operators out

$$\{q\}$$

$$\{-p\}$$

Step 2: apply the resolution rule

1. $\{p\}$

Premise

2. $\{q\}$

Premise

3. $\{-p\}$

Premise

4. $\{\}$

1,3

Example:

- We want to prove that

$$((B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})) \wedge \neg B_{1,1}) \models \neg P_{1,2}.$$

- So, we show that

$$((B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})) \wedge \neg B_{1,1}) \wedge P_{1,2}$$

is a contradiction (recall that $P_{1,2}$ is equivalent to $\neg\neg P_{1,2}$).


- We convert the above formula in CNF (see page 50):

$$\textcircled{1} (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg(P_{1,2} \vee P_{2,1}) \vee B_{1,1}) \wedge \neg B_{1,1} \wedge P_{1,2}$$

$$\textcircled{2} (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge ((\neg P_{1,2} \wedge \neg P_{2,1}) \vee B_{1,1}) \wedge \neg B_{1,1} \wedge P_{1,2}$$

$$\textcircled{3} (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg P_{1,2} \vee B_{1,1}) \wedge (\neg P_{2,1} \vee B_{1,1}) \wedge \neg B_{1,1} \wedge P_{1,2}$$

- So, we obtain the following clauses:



- The clauses we obtained are in the first row above.
- The second row shows clauses obtained by resolving pairs in the first row.
- When $P_{1,2}$ is resolved with $\neg P_{1,2}$ we obtain the empty clause.
- This means that

$$((B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})) \wedge \neg B_{1,1}) \wedge P_{1,2}$$

is a contradiction.

- So, we have proved

$$((B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})) \wedge \neg B_{1,1}) \models \neg P_{1,2}.$$

Deduction Theorem

Validity is connected to inference via the **Deduction Theorem**:

$KB \models a$ if and only if $KB \Rightarrow a$

The Deduction Theorem tells us that **a** is a logical consequence of KB if and only if $KB \Rightarrow a$ is a tautology.

$KB \models a$ if and only if $\neg(KB \wedge \neg a)$

This means that **a** is a logical consequence of KB if and only if $KB \wedge \neg a$ is a contradiction.

Probabilistic

Boolean algebras of set


Let W be a non-empty set and let 2^W be the powerset of W , i.e. the set of all subsets of W . A

Boolean algebra of sets \mathcal{F} is a subset of 2^W that is


- closed under intersection, i.e. for all $A, B \in \mathcal{F}$, $A \cap B \in \mathcal{F}$
- closed under union, i.e. for all $A, B \in \mathcal{F}$, $A \cup B \in \mathcal{F}$
- closed under complementation, i.e. for all $A \in \mathcal{F}$, $A^c \in \mathcal{F}$

Events:


- Let W be a set of possible worlds and let $A, B \subseteq W$ be events.
 - $A \cup B$ is the event that occurs if at least one of A and B occurs.
 - $A \cap B$ is the event that occurs if both A and B occur.
 - A^c is the event that occurs if A does not occur.
 - W is the certain event, i.e. the event that always occurs.
- Pick a card from a standard deck of 52 cards.
 - The sample space W is the set of all 52 cards.
 - Consider the following four events:
 - A: card is an ace.




- B: card has a black suit.



- D: card is a diamond.




- H: card is a heart:



- $A \cap H$:



- $A \cap B$:



Finitely Additive Probability Measures

Let W be a finite set and let \mathcal{F} be a Boolean algebra of sets. A **finitely additive probability measure** over \mathcal{F} is a function

$$\mu : \mathcal{F} \rightarrow [0, 1]$$

such that, for all $A, B \in \mathcal{F}$

- $\mu(W) = 1$
- $\mu(A \cup B) = \mu(A) + \mu(B) - \mu(A \cap B)$
- $\mu(\emptyset) = 0$
- $\mu(A^c) = 1 - \mu(A)$
- if $A \subseteq B$, then $\mu(A) \leq \mu(B)$

Definition

Given a (finite) set W , a probability distribution over W is a function

$$\pi : W \rightarrow [0, 1]$$

such that

$$\sum_{x \in W} \pi(x) = 1.$$

- For every probability distribution over a finite set it is possible to define a corresponding probability measure.
- Given a finite set W along with a Boolean algebra \mathcal{F} and a probability distribution π over W , for all $A \in \mathcal{F}$, the function

$$\mu(A) = \sum_{x \in A} \pi(x)$$

is a probability measure over \mathcal{F} .


Conditional Probability

If A and B are events, with $\mu(B) > 0$, the conditional probability of A given B , denoted by $\mu(A|B)$ is given by

$$\mu(A|B) = \frac{\mu(A \cap B)}{\mu(B)}$$

- A is the event we want to update and B is the evidence.
- $\mu(A)$ is the **prior** probability of A .
- $\mu(A|B)$ is the **posterior** probability of A .

($\mu(A|B)$ 等于 A 和 B 同时发生的部分的占总体的概率除以 B 占总体的概率)



- Take a finite set of possible worlds and an event A .
- We learn that B occurred.
- We can obtain $\mu(A | B)$ by eliminating the worlds in B^c and renormalising the probability of A over B .

Suppose I have 2 cats.

What is the probability that both are female, given that you know that at least one is a girl? What is the probability that both are female, given that you know that the younger is a girl?

$$\mu(\text{both girls} \mid \text{at least one girl}) = \frac{\mu(\text{both girls} \cap \text{at least one girl})}{\mu(\text{at least one girl})} = \frac{1/4}{3/4} = \frac{1}{3}$$

$$\mu(\text{both girls} \mid \text{younger is a girl}) = \frac{\mu(\text{both girls} \cap \text{younger is a girl})}{\mu(\text{younger is a girl})} = \frac{1/4}{1/2} = \frac{1}{2}$$

Independent

Events A and B are **independent** if

$$\mu(A \cap B) = \mu(A) \cdot \mu(B).$$

If $\mu(A) > 0$ and $\mu(B) > 0$, then this is equivalent to

$$\mu(A|B) = \mu(A)$$

$$\mu(B|A) = \mu(B)$$

Independence is completely different from **disjointness**.

- If A and B are disjoint, then $\mu(A \cap B) = 0$, so they are independent only if $\mu(A) = \mu(B) = 0$.
- **Disjoin**: A occurs tells us that B definitely did not occur, so A clearly conveys information about B, meaning the two events are not independent

Bayes' Rule

- For any events A,B with positive probabilities,

$$\mu(A \cap B) = \mu(B) \cdot \mu(A|B) = \mu(A) \cdot \mu(B|A)$$

- For any events A_1, \dots, A_n with positive probabilities,


$$\mu(\bigcap_{i=1}^n A_i) = \mu(A_1) \cdot \mu(A_2|A_1) \cdot \mu(A_3|A_1 \cap A_2) \cdots \mu(A_n|\bigcap_{j=1}^{n-1} A_j)$$

- Bayes' Rule

$$\mu(A|B) = \frac{\mu(B|A) \cdot \mu(A)}{\mu(B)}$$

- Let A_1, \dots, A_n be a partition of the set of possible worlds, with $\mu(A_i) > 0$ for all i. Then

$$\mu(B) = \sum_{i=1}^n \mu(B|A_i) \cdot \mu(A_i)$$



Since the A_i form a partition, then

$$B = (B \cap A_1) \cup (B \cap A_2) \cup \dots \cup (B \cap A_n).$$

Then, since the pieces are disjoint

$$\mu(B) = \mu(B \cap A_1) + \mu(B \cap A_2) + \dots + \mu(B \cap A_n).$$

Finally

$$\mu(B) = \mu(B \mid A_1) \cdot \mu(A_1) + \dots + \mu(B \mid A_n) \cdot \mu(A_n).$$

MINI-Max
