



Problem 1 (25 %)

Optimization problem:

$$\min x_1 + 2x_2 \quad \text{s.t.} \quad 2 - x_1^2 - x_2^2 \geq 0, \quad x_2 \geq 0 \quad (1)$$

a) The optimal point is $x^* = (-\sqrt{2}, 0)^\top$, this can be found by inspection.

b) The Lagrangean for the problem is

$$\mathcal{L}(x, \lambda) = x_1 + 2x_2 - \lambda_1 c_1(x) - \lambda_2 c_2(x) \quad (2)$$

with

$$c_1(x) = 2 - x_1^2 - x_2^2 \quad (3a)$$

$$c_2(x) = x_2 \quad (3b)$$

$$\mathcal{I} = \{1, 2\} \quad (3c)$$

Hence,

$$\nabla_x \mathcal{L}(x^*, \lambda^*) = \begin{bmatrix} 1 + 2\lambda_1^* x_1^* \\ 2 + 2\lambda_1^* x_2^* - \lambda_2^* \end{bmatrix} = 0 \Rightarrow \lambda^* = \begin{bmatrix} \frac{1}{2\sqrt{2}} \\ 2 \end{bmatrix} \quad (4)$$

and all KKT conditions are satisfied.

- c) The gradients of the active constraints and the objective function at the solution are illustrated in Figure 1.
- d) A positive number means any small change of x results in an increase of the objective function.
- e) This is a convex problem, since the objective function is convex (all linear functions are linear) and the feasible set is convex. For a problem with only inequality constraints, the feasible set is convex if all inequalities are concave functions. Here, both $c_1(x)$ (a paraboloid with a unique maximizer) and $c_2(x)$ (a linear function) are concave functions. (Note that a linear function is both concave and convex).




Figure 1: Gradients at the optimal point in Problem 1.

Problem 2 (30 %)

Optimization problem:

$$\min 2x_1 + x_2 \quad \text{s.t.} \quad x_1^2 + x_2^2 - 2 = 0 \quad (5)$$

a) The extreme points are

$$x = \left(-2\sqrt{\frac{2}{5}}, -\sqrt{\frac{2}{5}} \right)^\top \quad (6a)$$

and

$$x = \left(2\sqrt{\frac{2}{5}}, \sqrt{\frac{2}{5}} \right)^\top \quad (6b)$$

(Use, for instance, $\nabla f = \lambda_1 \nabla c_1$ and $x_1^2 = 2 - x_2^2$ to find these.)

b) The Lagrangian for the problem is

$$\mathcal{L}(x, \lambda) = 2x_1 + x_2 - \lambda_1 c_1(x) \quad (7)$$

with

$$c_1(x) = x_1^2 + x_2^2 - 2, \quad \mathcal{E} = \{1\} \quad (8)$$

Hence,

$$\nabla_x \mathcal{L}(x^*, \lambda^*) = \begin{bmatrix} 2 - 2\lambda_1^* x_1^* \\ 1 - 2\lambda_1^* x_2^* \end{bmatrix} = 0 \Rightarrow \lambda^* = \pm \frac{\sqrt{10}}{4} \quad (9)$$

- c) The values of the multiplier above mean that the KKT conditions are satisfied at both extreme points.
- d) The gradients of the active constraint and the objective function at the optimal point are illustrated in Figure 2.




Figure 2: Gradients at the optimal point in Problem 2.

- e) We have that

$$\nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*) = \begin{bmatrix} -2\lambda^* & 0 \\ 0 & -2\lambda^* \end{bmatrix} > 0 \quad (10)$$

holds for $\lambda^* = -\frac{\sqrt{10}}{4}$. Then,

$$w^\top \nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*) w > 0, \quad \text{for all } w \in \mathcal{C}(x^*, \lambda^*), \quad w \neq 0 \quad (11)$$

which means that both the necessary and sufficient second-order conditions in Chapter 12.5 hold (Theorems 12.5 and 12.6, respectively).

- f) The problem is nonconvex due to the nonlinear equality constraint.

Problem 3 (20 %)

Optimization problem:

$$\min_{x \in \mathbb{R}^2} f(x) = -2x_1 + x_2 \quad \text{s.t.} \quad \begin{cases} c_1(x) = (1 - x_1)^3 - x_2 \geq 0 \\ c_2(x) = x_2 + 0.25x_1^2 - 1 \geq 0 \end{cases} \quad (12)$$

a) The set of active constraint gradients at the solution,

$$\nabla c_1(x^*) = \begin{bmatrix} -3 \\ -1 \end{bmatrix}, \quad \nabla c_2(x^*) = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad (13)$$

is linearly independent. Hence, LICQ hold.

b) The Lagrangian for the problem is

$$\mathcal{L}(x, \lambda) = -2x_1 + x_2 - \lambda_1 c_1(x) - \lambda_2 c_2(x) \quad (14)$$

with

$$c_1(x) = (1 - x_1)^3 - x_2 \quad (15a)$$

$$c_2(x) = x_2 + 0.25x_1^2 - 1 \quad (15b)$$

$$\mathcal{I} = \{1, 2\} \quad (15c)$$

Hence,

$$\nabla_x \mathcal{L}(x^*, \lambda^*) = \begin{bmatrix} -2 + 3\lambda_1^*(1 - x_1^*)^2 - 0.5\lambda_2^*x_1^* \\ 1 + \lambda_1 - \lambda_2 \end{bmatrix} = 0 \Rightarrow \lambda^* = \begin{bmatrix} 2/3 \\ 5/3 \end{bmatrix} \quad (16)$$

and all KKT conditions are satisfied.

d) The Hessian of the Lagrangian at the solution,

$$\nabla_{xx}^2 \mathcal{L}(x^*, \lambda^*) = \begin{bmatrix} -6\lambda_1^*(1 - x_1^*) - 0.5\lambda_2^* & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} -29/6 & 0 \\ 0 & 0 \end{bmatrix} \quad (17)$$

is negative semidefinite. However, the critical cone $\mathcal{C}(x^*, \lambda^*)$ contains only the vector $w^\top = [0, 0]$. Hence, the second-order necessary conditions are satisfied, whereas the second-order sufficient conditions are not.

Problem 4 (25 %)

Finding the maximizer for $f(x) = x_1x_2$ is equivalent to finding the minimizer for the function $\bar{f} = -x_1x_2$. We therefore state the optimization problem as

$$\min -x_1x_2 \quad \text{s.t.} \quad 1 - x_1^2 - x_2^2 \geq 0 \quad (18)$$

Note that the objective function represents a saddle, and it is therefore clear that the minimizer(s) exist(s) on the boundary of the unit disk. The Lagrangian is given by

$$\mathcal{L}(x, \lambda) = -x_1x_2 - \lambda_1(1 - x_1^2 - x_2^2) \quad (19)$$

From the KKT conditions,

$$\nabla_x \mathcal{L}(x^*, \lambda^*) = \begin{bmatrix} -x_2^* + 2\lambda_1^*x_1^* \\ -x_1^* + 2\lambda_1^*x_2^* \end{bmatrix} = 0 \Rightarrow \lambda_1^* = \pm \frac{1}{2} \quad (20)$$

Since λ_1^* has to be nonnegative, we have

$$x_1^* = x_2^* = \pm \frac{1}{\sqrt{2}} \quad (21)$$

Hence, $f(x) = x_1x_2$ has two maximizers, $x^* = \left[-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right]$ and $x^* = \left[\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right]$, both with $\lambda_1^* = +\frac{1}{2}$. The gradients at the optimal point $x^* = \left[-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right]$ are illustrated in Figure 3, while the gradients at the optimal point $x^* = \left[\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right]$ are illustrated in Figure 4.




Figure 3: Gradients at the solution $x^* = \left[-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right]$ of Problem 4.




Figure 4: Gradients at the solution $x^* = \left[\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right]$ of Problem 4.