



# **national carbon accounting system**

## **Developing a National Forest Productivity Model**

**Jenny Kesteven,  
Joe Landsberg and  
URS Australia**



**Australian Government**  
**Australian Greenhouse Office**

**technical report no. 23**

## ***The National Carbon Accounting System:***

- Supports Australia's position in the international development of policy and guidelines on sinks activity and greenhouse gas emissions mitigation from land based systems.
- Reduces the scientific uncertainties that surround estimates of land based greenhouse gas emissions and sequestration in the Australian context.
- Provides monitoring capabilities for existing land based emissions and sinks, and scenario development and modelling capabilities that support greenhouse gas mitigation and the sinks development agenda through to 2012 and beyond.
- Provides the scientific and technical basis for international negotiations and promotes Australia's national interests in international fora.

<http://www.greenhouse.gov.au/ncas>

For additional copies of this report phone 1300 130 606

# **DEVELOPING A NATIONAL FOREST PRODUCTIVITY MODEL**

---

**Jenny Kesteven<sup>+</sup>, Joe Landsberg<sup>#</sup> and URS Australia<sup>o</sup>**

**<sup>+</sup>Centre for Resource and Environmental Studies,  
Australian National University, Canberra**

**<sup>#</sup>Landsberg Consulting Pty Ltd**

**<sup>o</sup>URS Australia**

**National Carbon Accounting System Technical Report No. 23**

**May 2004**



**Australian Government**

**Australian Greenhouse Office**

---

Printed in Australia for the Australian Greenhouse Office

© Commonwealth of Australia 2004

This work is copyright. It may be reproduced in whole or part for study or training purposes subject to the inclusion of an acknowledgement of the source and no commercial usage or sale results. Reproduction for purposes other than those listed above requires the written permission of the Communications Team, Australian Greenhouse Office. Requests and enquiries concerning reproduction and rights should be addressed to the Communications Team, Australian Greenhouse Office, GPO Box 621, CANBERRA ACT 2601.

For additional copies of this document please contact the Australian Greenhouse Office Publications Hotline on 1300 130 606.

For further information please contact the National Carbon Accounting System at <http://www.greenhouse.gov.au/ncas/>

Neither the Australian Government nor the Consultants responsible for undertaking this project accepts liability for the accuracy of or inferences from the material contained in this publication, or for any action as a result of any person's or group's interpretations, deductions, conclusions or actions in reliance on this material.

May 2004

*Environment Australia Cataloguing-in-Publication*

Kesteven, Jennifer L.

Developing a national forest productivity model / Jenny Kesteven, Joe Landsberg and URS Australia.

p. cm.

(National Carbon Accounting System technical report; No. 23)

Bibliography

ISSN: 1442 6838

1. Forest productivity-Australia-Simulation methods. 2. Forest productivity-Australia-Forecasting. I. Landsberg, J.J. (Joseph John), 1938- . II. URS Australia. III. Australian Greenhouse Office.

IV. Series

333.75'0994'011-dc22

## SUMMARY

Knowledge of the spatial and temporal patterns of forest growth is fundamental to estimating the carbon stocks (and biomass) of mature forests and rates of carbon accumulation in any forest regrowth.

As part of the National Carbon Accounting System developed by the Australian Greenhouse Office, indices of forest growth were developed and used to predict potential biomass at maturity (forest productivity) and rates for biomass increment across the Australian continent over time.

This report documents the application of these forest growth indices in developing a National Forest Productivity Model.



# TABLE OF CONTENTS

|   | Page No. |
|---|----------|
| Summary   | iii      |
| List of Symbols and Acronyms  | vii      |
| 1. Introduction   | 1        |
| 2. Input Data   | 2        |
| 2.1 Soils   | 2        |
| 2.1.1 Soil Water Holding Capacity ( $S_C$ )   | 2        |
| 2.1.2 Soil Nutrient Status ( $S_N$ )  | 4        |
| 2.2 Climate   | 5        |
| 2.2.1 Fitting the Climate Surfaces  | 5        |
| 2.2.1.1 Errors in the Model   | 7        |
| 2.2.1.2 From Surface Coefficient Files to Maps  | 8        |
| 2.2.2 Digital Elevation Model (DEM)   | 8        |
| 2.2.3 Station Dictionary  | 10       |
| 2.2.4 Long-Term Average Climate Variables   | 11       |
| 2.2.4.1 Average Temperature   | 12       |
| 2.2.4.2 Number of Frost Days  | 12       |
| 2.2.4.3 Radiation   | 12       |
| 2.2.4.4 Vapour Pressure Deficit ( $VPD$ )   | 15       |
| 2.2.5 Monthly Average Climate Variables   | 15       |
| 2.2.5.1 Average Temperature   | 16       |
| 2.2.5.2 Maximum Temperature   | 17       |
| 2.2.5.3 Minimum Temperature   | 19       |
| 2.2.5.4 Rainfall  | 20       |
| 2.2.5.5 Frost Days  | 23       |
| 2.2.5.6 Radiation   | 24       |
| 2.3 Normalised Difference Vegetation Index (NDVI)   | 24       |
| 3. Calculation of Plant Productivity  | 26       |
| 3.1 Temperature Modifier  | 26       |
| 3.2 Frost Modifier  | 27       |
| 3.3 Vapour Pressure Deficit ( $VPD$ ) Modifier  | 27       |
| 3.4 Soil Water Content ( $S_W$ )  | 28       |
| 3.5 Absorbed Photosynthetically Active Radiation (APAR)   | 32       |
| 3.6 Plant Productivity Index  | 33       |
| 4. Select References  | 36       |
| Appendix 1 – Preliminary Investigations: Spatial Estimation of Plant Productivity and Classification of Non-Commercial Vegetation Types | 39       |

# LIST OF FIGURES

|  | Page No. |
|--|----------|
| Figure 1. Potential available soil water holding capacity ( $S_C$ ).   | 3        |
| Figure 2. Soil nutrient status ( $S_N$ ).  | 4        |
| Figure 3. The Australian DEM.  | 9        |
| Figure 4. A screen capture for station 061121 – Lostock Post Office.   | 11       |
| Figure 5. Long-term monthly average temperature.   | 13       |
| Figure 6. Long-term average number of frost days per month.  | 14       |
| Figure 7. Annual average temperature derived from monthly data grids for 1968-2002 and examples of low (1976), average (1982) and high (1996) temperature years. | 16       |
| Figure 8. Australian average maximum temperature ( $T_{max}$ ) and the 12 month running mean.  | 17       |
| Figure 9. Maximum temperature grids for high and low continental average values for July and December.   | 18       |
| Figure 10. Australian average minimum temperature ( $T_{min}$ ) and the 12 month running mean.   | 19       |
| Figure 11. Minimum temperature grids for high and low continental average values for July and February.  | 20       |
| Figure 12. Australian continental average monthly rainfall.  | 21       |
| Figure 13. Annual rainfall 1970–2002.  | 22       |
| Figure 14. Australian continental average annual number of frost days.   | 23       |
| Figure 15. Example grids of low (1991, average = 6.1 days), medium (1987, average = 14.0 days) and high (1982, average = 21.1 days) number of days frost.        | 24       |
| Figure 16. Temperature modifier spatial equations (from Equation 11) for January and July long-term averages.  | 26       |
| Figure 17. The frost modifier equation for the July long-term average.   | 27       |
| Figure 18. VPD modifier spatial equations for January and July long-term averages.   | 28       |
| Figure 19. Spatial equation for equilibrium evaporation, January 1995.   | 29       |
| Figure 20. Spatial equation for transpiration, January 1995.   | 29       |
| Figure 21. Spatial equation for interception, January long term average.   | 29       |
| Figure 22. Spatial equation for water balance, January 1995.   | 30       |
| Figure 23. Spatial equation for soil water content, January 1995.  | 30       |
| Figure 24. Spatial equation for moisture ratio, January 1995.  | 31       |
| Figure 25. Spatial equation for soil water modifier, January 1995.   | 32       |
| Figure 26. Long-term average APAR for January and July.  | 33       |
| Figure 27. Spatial equations for productivity indices, January and July long-term averages.  | 33       |
| Figure 28. Derived long-term average plant productivity index for Australia.   | 34       |
| Figure 29. Derived annual continental productivity indices 1970-2002.  | 35       |



## **LIST OF TABLES**

|                 | Page No.   |
|-----------------|--|
| <b>Table 1</b>  | ANUCLIM variables used in the plant productivity modelling.                |
| <b>Table 2.</b> | ANUSPLIN variables created for the long-term plant productivity modelling. |
| <b>Table 3.</b> | Monthly climate variables.   |

# LIST OF SYMBOLS AND ACRONYMS

|                |  |                      |   |
|----------------|--|----------------------|---|
| AGO            | Australian Greenhouse Office   | $P$                  | Rainfall (mm)   |
| APAR           | Absorbed Photosynthetically Active Radiation                               | $R_A$                | Adjusted direct radiation   |
| BoM            | Bureau of Meteorology (Australia)  | $R_C$                | Slope and aspect modified radiation                                       |
| CRES           | Centre for Resource and Environmental Studies                              | $R_D$                | Direct radiation  |
| CSIRO          | Commonwealth Scientific and Industrial Research Organization               | $R_F$                | Diffuse radiation   |
| DEM            | Digital Elevation Map  | $R_G$                | Global radiation  |
| $e^{\circ}(T)$ | Saturation vapour pressure at air temperature $T$ (kPa)                    | $R_N$                | Net radiation ( $\text{MJ m}^{-2} \text{ day}^{-1}$ )                     |
| $E_{eq}$       | Equilibrium evaporation  | $R_P$                | Radiation with rainfall   |
| $exp(x)$       | 2.7183 (base of natural logarithm) raised to the power $x$                 | $S_N$                | Soil nutrient status  |
| $F$            | Frost (days per month)   | $S_C$                | Soil water holding capacity   |
| $F_{mod}$      | Frost modifier   | $S_W$                | Soil water  |
| $fPAR$         | Fraction of Photosynthetically Active Radiation absorbed                   | $S_{Wmod}$           | Soil water modifier   |
| GCV            | Generalised Cross Validation   | $T$                  | Temperature ( $^{\circ}\text{C}$ )  |
| GPP            | Gross Primary Productivity   | $T_{avg}$            | Average monthly temperature / mean air temperature ( $^{\circ}\text{C}$ ) |
| I              | Interception intercept   | $T_{dew9}, T_{dew3}$ | Dew point temperature (9 am and 3 pm)                                     |
| J              | Month  | $T_{dry9}, T_{dry3}$ | Dry bulb temperature (9 am and 3 pm)                                      |
| LAI            | Leaf Area Index ( $\text{m}^2$ (leaf area) $\text{m}^{-2}$ (soil surface)) | $T_{high}$           | Monthly average temperature above which plant growth stops                |
| M              | Moisture ratio   | $T_{low}$            | Monthly average temperature below which plant growth stops                |
| NDVI           | Normalised Difference Vegetation Index                                     | $T_{max}$            | Maximum air temperature ( $^{\circ}\text{C}$ )                            |
| $\lambda_I$    | Near infra-red electromagnetic wavelengths                                 | $T_{min}$            | Minimum air temperature ( $^{\circ}\text{C}$ )                            |
| $\lambda_R$    | Low micron range electromagnetic wavelengths                               | $T_{mod}$            | Temperature modifier  |
| NPP            | Net Primary Production   | $T_{opt}$            | Optimum temperature for growth  |
|                |  | $Trans$              | Transpiration   |
|                |  | $VPD$                | Vapour Pressure Deficit   |
|                |  | $VPD_{mod}$          | Vapour Pressure Deficit modifier  |
|                |  | $W$                  | Water balance   |

## 1. INTRODUCTION

Net primary productivity (NPP) is the rate at which chemical or solar energy is converted to biomass. The main primary producers are the green plants, which convert solar energy, carbon dioxide and water to glucose, and eventually to plant tissue.

Estimation of productivity can be obtained through different methods. Direct measurement methods include destructive sampling of the above-ground and below-ground plant biomass, and the recording of carbon dioxide fluxes at the vegetation-atmosphere interface.

Available direct measurements of NPP have some limitations for mapping productivity across the Australian continent. The reliability of available data is variable – the data is limited in number and not evenly distributed among the various types of ecosystems. To overcome these deficiencies in measured data several model types have been developed. These are either physiological models simulating ecosystem fluxes from environmental variables, remote sensing-based models interpreting the light spectrum reflected by the land surface, or inverse models deducing fluxes from time and space variations in atmospheric CO<sub>2</sub> and <sup>13</sup>C data. Process models with physiological functions (i.e. those that acknowledge temporal variability due to changing biophysical conditions such as rainfall and temperature) offer the opportunity to simulate past and future changes in NPP according to environmental changes.

A truncated spatial version of the 3-PG (process) model (after Landsberg and Waring, 1997) was developed as part of the National Carbon Accounting System (NCAS) to predict forest productivity across the Australian continent. The resultant 'productivity index' model as documented here retains the essential features of biomass NPP estimation, without the biomass fixation (Gross Primary Productivity (GPP) minus NPP) and carbon partitioning procedures.

A relative index of plant productivity was mapped for Australia using the 'productivity index' model – based on the relationship between the amount of photosynthetically active radiation absorbed by plant canopies (APAR) and the various productivity modifiers that affect plant growth (e.g. temperature, soil water content, frost). The model used a monthly time step to derive both a long-term average monthly productivity index (~250 m resolution) and a monthly productivity index for 1970 to 2002 (1 km resolution).

Factors converting APAR to productivity indices were reduced from presumed optimum values by modifiers dependent on soil fertility, atmospheric vapour pressure deficits, soil water content and temperature. The ANUCLIM and ANUSPLIN programs were used to generate climate surfaces for the continent. Soil fertility and water holding capacity values were obtained from the CSIRO using the digital Soil Atlas of Australia. Leaf Area Index, essential for the calculation of APAR, was estimated from 10-year mean values of Normalised Difference Vegetation Indices (NDVI), for 1 km pixels, for the entire country. Incoming short-wave radiation – and hence APAR – was corrected for slope and aspect using a Digital Elevation Map (DEM) for the long-term average, but with 1 km pixels being used in the monthly productivity maps derived as this made no significant difference to the results. Analyses were, therefore, carried out on estimates based on the assumption of flat terrain.

The resultant maps (as digital grids) of plant productivity index, for each month for Australia, provide relative indices of plant productivity (reflecting the spatial and temporal patterns of plant growth) in any region. The data generated is integral to the estimation of carbon stocks under the NCAS and provides a framework for assessing carbon accumulation by various vegetation types.

## 2. INPUT DATA

The production of mapped productivity indices for Australia required integration and analysis of a number of spatially represented environmental variables. These included: soil water holding capacity (to estimate water balance), soil nutrient status, Advanced Very High Resolution Radiometer (AVHRR) data (to estimate the Normalised Difference Vegetative Index (NDVI)), and various climate variables (temperature, rainfall and number of frost days).

### 2.1 SOILS

The Atlas of Australian Soils (Northcote *et. al.* 1960–1968) was integrated with a set of interpreted soil variables from CSIRO (McKenzie and Hook 1992) to produce continental maps of potential available water holding capacity (PAWHC) and nutrient status of soils.

The Atlas of Australian Soils completed in 1968 (Northcote *et. al.* 1960–1968) and made available in digital form in 1990 provides the only consistent source of spatial information for the continent. While large areas have been surveyed in more detail during the last 30 years, those surveys have not been compiled to produce an updated continental map.

McKenzie *et. al.* 2000 provided a set of interpreted soil variables (from McKenzie and Hook 1992) that can be used with the Digital Atlas. The interpretations for each soil type were based, wherever possible, on data held within the CSIRO National Soil Database. Summaries from the database were used with other sources of information to assign an interpreted value for each variable.

Soil fertility and water holding capacity ( $S_c$ ) were obtained by merging the Atlas of Australian soils (Northcote *et. al.* 1960–1968) and data provided by McKenzie *et. al.* 2000. The data provided included soil fertility and water holding capacity for both the

dominant soil type (by area) and the average of the top five dominant soils. McKenzie (pers. comm.) advised using the properties of the dominant soil type (by area) to characterise the region. Caveats on the use of the Digital Atlas of Australian Soils are presented in McKenzie *et. al.* (2000).

The resultant data had limitations, as a large proportion of soil variation within a region occurred over short distances which could not be resolved at the mapping scale of the Atlas. The qualitative nature of the Digital Atlas and restrictions associated with the classification scheme and structure of the soil-landscape model imposed further constraints.

The Atlas of Australian Soils uses 725 soil profile classes, normally at the level of Principle Profile Form (PPF) (e.g. Ug5.15). The legend of the Atlas defines 3,060 map unit types. The map unit types have various combinations of the 725 soil profile classes. The map unit type descriptions identify dominant and subdominant soil profile classes. Many of the map unit types occur more than once and the Digital Atlas has 22,560 mapped polygons.

McKenzie and Hook (1992) prepared interpretations of the 725 soil profile classes. The dominant soil in each map unit type was identified and interpreted values for each soil profile class were ascribed to each map unit type. Lookup tables were created from the accompanying spreadsheets. Some soil polygons had PPFs containing 'Undefined' or 'NS' - their soil attributes were coded so as to represent no data.

#### 2.1.1 Soil Water Holding Capacity ( $S_c$ ).


Available water capacity is presented on a per unit depth basis, as a total for each horizon, and as a total for the solum. The total solum was recommended (McKenzie pers. comm.) for use. The available water capacities were calculated as the difference in volumetric water content at matric potentials of  $-0.1$  bar and  $-15$  bar for a specified depth increment.

It should be noted that the total available water capacity for the solum is constrained by limitations associated with the estimate of solum thickness of the actual plant available water capacity (see Hillel 1980). Despite these limitations, McKenzie *et. al.* 2000 suggest that it provides a reasonable first approximation of the water storage capacity of a soil.

In the resultant  $S_C$  maps derived, PPFs with an  $S_C$  equal to 0 were assigned the value of the A Horizon  $S_C$ , except for PPF Dy5.91 (comprising 2 spatial units in Kangaroo Island SA) which was assigned the PPF2  $S_C$  as there was no PPF1 A Horizon value.

Mapped values (with a 250 m resolution) ranged from 21 to 270 mm with a mean value of 104 mm (see Figure 1).

Initial soil water content was set to 75% of the potential available water holding capacity. As this is not a true representation of the January soil water content for Australia, the water balance routine was run for two years to derive a reliable estimation of the actual soil water content to initialise the model for the long-term average. The water balance routine was run for the years 1968 and 1969 to estimate initial soil water holding capacity for the 1970 to 2002 monthly productivity model.



**Figure 1. Potential available soil water holding capacity ( $S_C$ ).**



### 2.1.2 Soil Nutrient Status ( $S_N$ )

Because of natural variation and the considerable uncertainty surrounding soil fertility values, only three levels of fertility were used:

- high (effective modifier = 1),
- medium (effective modifier = 0.8) and
- low (effective modifier = 0.6)

giving  $S_N$  values of 1.25, 1 and 0.75, respectively. These were applied to each polygon representing a soil type in the Atlas of Australian Soils (Figure 2).

The rating system for gross nutrient status prepared by McKenzie and Hook (1992) relates to the behaviour of profiles under agricultural development. Profiles with a low status (class 1)

exhibited major responses to nitrogen, phosphorus and potassium along with most micronutrients. Profiles with a moderate nutrient status (class 2) responded to nitrogen and phosphorus with occasional responses to some micronutrients. It was uncommon for profiles with a high nutrient status (class 3) to respond to nitrogen and phosphorus except after intensive farming. The main sources of information for the assessment of nutrient status were Stace *et. al.* (1968) and Northcote *et. al.* (1975).

In the resultant map derived for application in the productivity model, soil polygons with no nutrient value were assigned the lowest class i.e.  $S_N = 0.75$ . The average value for Australia was 0.81 (see Figure 2).




Figure 2. Soil nutrient status ( $S_N$ ).

## 2.2 CLIMATE

Climate data was needed to produce the long-term average productivity maps and the 1968 to 2002 monthly productivity maps. Spatial coverage of long-term average climate surfaces for several climate variables were available from the ANUCLIM package. ANUCLIM (incorporating ESOCLIM, BIOCLIM and GROCLIM) is used to store climate surfaces derived by ANUSPLIN and to enable systematic interrogation of these surfaces, in point and grid form, for biophysical applications. Most surfaces need a Digital Elevation Model (DEM) to provide climate values in grid form (see <http://cres.anu.edu.au>). These climate variables include:

- Long-term average rainfall ( $P$ )
- Long-term average radiation ( $Rad$ )
- Long-term average minimum temperature ( $T_{min}$ )
- Long-term average maximum temperature ( $T_{max}$ )
- Long-term average 9am and 3pm dry bulb temperature ( $T_{dry}$ )
- Long-term average 9am and 3pm dew point temperature ( $T_{dew}$ )

Data layers that were required as inputs for the productivity model but not available in the ANUCLIM package included:

- Long-term average temperature ( $T_{avg}$ )
- January 1968 to December 2002 monthly minimum temperature ( $T_{min}$ )
- January 1968 to December 2002 monthly maximum temperature ( $T_{max}$ )
- January 1968 to December 2002 monthly average temperature ( $T_{avg}$ )

- January 1968 to December 2002 monthly rainfall ( $P$ )
- Long-term average number of days with frost ( $F$ )
- January 1968 to December 2002 monthly number of days with frost ( $F$ ).

### 2.2.1 Fitting The Climate Surfaces

---

Climate surfaces for long-term average temperature and frosts and for all the monthly climate variables from 1968 to 2002 were created using the ANUSPLIN software package (see <http://cres.anu.edu.au>). The ANUSPLIN package provides a facility for transparent analysis and interpolation of noisy multi-variate data using thin plate smoothing splines. The package supports this aim by providing comprehensive statistical analyses, data diagnostics and spatially distributed standard errors. It also supports flexible data input and surface interrogation procedures. The climate surfaces available in the ANUCLIM package were created using the ANUSPLIN package.

A brief description of the surface fitting techniques is given below but for a more comprehensive discussion see Kesteven (1998), Hutchinson (1991a, 1991b), Kesteven and Hutchinson (1996) and Wahba (1990).

The original thin plate (formerly Laplacian) surface fitting technique was described by Wahba (1979), with modifications for larger data sets due to Bates and Wahba (1982), Elden (1984) and Hutchinson (1984). Thin plate smoothing spline interpolation is a global method as it uses all the given data for prediction at each point (Laslett *et. al.* 1987). The method is also non-parametric and so is relatively insensitive to the distribution of the parent population.

Thin plate smoothing splines can be viewed as a generalisation of standard multi-variate linear regression, in which the parametric model is replaced by a suitably smooth non-parametric function. The degree of smoothness, or inversely the degree of complexity, of the fitted function is usually determined automatically from the data by minimising a measure of predictive error of the fitted surface given by the generalised cross validation (GCV).

The main idea is to consider a spatial variable as a realisation of a spatially autocorrelated random function that accounts for the complicated behaviour of natural spatial data:

$$z_i = f(x_i) + \varepsilon_i \quad (i = 1, \dots, n) \quad (1)$$

where  $f$  is a function to be estimated from the observations  $z_i$ , which include a zero mean, usually spatially discontinuous, error term  $\varepsilon_i$ . The  $x_i$  are commonly assumed to represent coordinates in two- or three-dimensional Euclidean space.

The value of the variable over a region is interpolated by estimating the true underlying function. This function is estimated to be as smooth as possible without significantly distorting the given data values (Hutchinson 1991a). The observational model for a thin plate spline with three independent spline variables is that the observed monthly mean  $z_i$  at the position  $x_i, y_i, z_i$  of the  $i^{th}$  station is given by Equation 1, where  $f$  is an unknown smooth function and the  $\varepsilon_i$  are independent random errors with zero mean and variance  $d_i s^2$ , where  $d_i$  is the (local) relative variance and  $s^2$  is the common (normally unknown) variance.

The unknown smooth function is estimated by finding the function  $f$  (a thin plate spline) which minimises

$$\sum_{i=1}^n \left[ \frac{z_i - f(x_i, y_i, h_i)}{d_i} \right]^2 + \lambda J_m(f) \quad (2)$$

where  $J_m(f)$  is a measure of the roughness of  $f$  in terms of the  $m^{th}$  order derivatives of  $f$  (usually second order), and  $\lambda$  is a positive smoothing parameter.

In meteorological and climatological applications the weights,  $d_i$ , are used when the data do not have equal length records. The smoothing parameter  $\lambda$  determines a trade-off between data fidelity and surface roughness. It is usually calculated by minimising the generalised cross validation (GCV). This is a measure of the predictive error of the fitted surface which is calculated by removing each data point in turn and summing, with appropriate weighting, the square of the discrepancy of each omitted data point from a surface fitted to all the other points. The value of  $\lambda$  is chosen to minimise this sum, that is, to minimise the predictive error as ascertained by the performance of the fitted function in predicting omitted data points.

It is possible to calculate the GCV implicitly, and hence efficiently, because the fitted values depend linearly on the data. Intuitively the GCV is a good measure of the predictive power of the fitted surface, as has been verified both theoretically and in applications to real and simulated data (Wahba 1990 and Hutchinson 1991a).

Statistical interpretation proceeds by way of analogy with least squares regression analysis. Thin plate splines can be viewed as a non-parametric generalisation of linear regression analysis. A comprehensive introduction to the technique of thin plate smoothing splines, with various extensions, is given in Wahba (1990). A summary of the basic methodology, with climate interpolation principally in mind, can be found in Hutchinson (1991a). More comprehensive discussion of the algorithms and the associated statistical analyses are given in Hutchinson and Gessler (1994) and Hutchinson (1993, 1995). Comparisons with geostatistical (kriging) methods have been presented by Hutchinson (1993), Hutchinson and Gessler (1994) and Laslett (1994).



The surface fitting procedure was primarily developed for the task of fitting climate data so that there are normally at least two independent spline variables, longitude and latitude in units of decimal degrees. A third independent variable, elevation above sea-level, is often appropriate when fitting surfaces to temperature or precipitation. This is normally included as a third independent spline variable and was scaled to be in units of kilometres. Minor improvements can sometimes be made by slightly altering this scaling of elevation. This scaling was originally determined by Hutchinson and Bischof (1983) and has been verified by Hutchinson (1995, 1996). Extension to real time simulation and surface fitting is discussed by Hutchinson (1995) and Kesteven and Hutchinson (1996).

Over restricted areas, superior performance can sometimes be achieved by including elevation, not as an independent spline variable but as an independent covariate. Thus, in the case of fitting a temperature surface, the coefficient of an elevation covariate would be an empirically determined temperature lapse rate (Kesteven 1998, Hutchinson, 1991a). Other factors which influence the climate variable may be included as additional covariates if appropriate parameterisations can be determined and the relevant data are available. These might include, for example, topographic effects other than elevation above sea-level such as slope and aspect. Other applications to climate interpolation have been described by Hutchinson *et. al.* (1984) and Hutchinson (1989, 1991a, 1991b). Applications of fitted spline climate surfaces to global agroclimatic classifications and to the assessment of biodiversity are described by Hutchinson *et. al.* (1992).

To fit multi-variate climate surfaces, the values of the independent variables need to be known at the data points. Thus meteorological stations should be accurately located in position and elevation. Errors in these locations are often indicated by large values in the output ranked residual list. A number of applications have examined the utility of using

elevation and slope and aspect obtained from digital elevation models of various horizontal resolutions (Hutchinson 1995, 1998b).

### 2.2.1.1 Errors in the Model

In applying data smoothing to climate means, two components of the covariance structure of the  $\varepsilon_i$  in the model given by Equation 1 should be recognised. The first component allows for deficiencies in the model being fitted. These are essentially due to local effects, including measurement error, which are below the spatial resolution of the observed point data network. The magnitude of these effects depends on the adequacy of the model represented by the function  $z_i$  and on the spatial density of the data. These effects can be reasonably assumed to be independent between different locations.

The second component arises when using serially incomplete data to interpolate monthly means for a standard period. This component can be ignored for some variables such as temperature. Monthly means of temperature stabilise after a few years. However, this component is significant for rainfall which normally exhibits large variation from year to year. This variability gives rise to strong correlations between the error components of rainfall means at stations with common periods of record.

Hutchinson (1995) shows how the resulting correlated error structure can be incorporated into the interpolation process. The model permits the rigorous estimation of this smooth function by a thin plate smoothing spline in two ways. The statistical error structure of the data can either be accommodated directly, by using the corresponding non-diagonal error covariance matrix, or observed means can first be standardised to long-term estimates using linear regression. Both methods give similar interpolation accuracy and error estimates of the fitted surfaces were in good agreement with residuals from withheld data (Hutchinson 1995).

There are several useful output statistics in the diagnostics file generated by the SPLINA program, which can be used to check for data homogeneity. The square root of the generalised cross validation (RTGCV) is a good measure of the predictive power of the fitted surface, especially when comparing the values from the same climate variable fitted for different times. In combination with the ranked residuals listing, which allows for each station to be assessed in relation to its deviation from the calculated surface, the RTGCV provides a powerful tool for investigation of station homogeneity. This process of assessing each station in relation to its deviation from the calculated surface and correcting errors may be repeated many times, steadily reducing overall surface error. Once the statistics reach a level where there is very little change with the removal of additional stations and residuals are within expected ranges, the fitted surface coefficients are used to create the monthly surfaces. Coefficients of the surfaces fitted to the monthly values were stored by month. Values for any month at any site or for grids can be calculated using the LAPPNT or LAPGRD programs in the ANUSPLIN package. A visual representation for each month was created using ARCInfo although they could be displayed by any of a number of commonly available computer programs which can display raster images.

### **2.2.1.2 From Surface Coefficient Files to Maps**

The LAPGRD program can be used to calculate regular grids of fitted climate values and their standard errors, for mapping and other purposes, provided a regular grid of values of each independent variable, additional to longitude and latitude, is supplied. This usually means that a regular grid digital elevation model (DEM) is required.

Coefficients of the surfaces fitted to the monthly values are stored by year. They can be used to calculate values of the climate variable for any month at any site using the LAPPNT program in the


ANUSPLIN package. Alternatively, regular grids across Australia of monthly climate variables for any month can be calculated using the LAPGRD program in ANUSPLIN and displayed or used by any of a number of commonly available computer programs which can display raster images. The system can be updated to incorporate later years as new data become available.

### **2.2.2 Digital Elevation Model (DEM)**

A Digital Elevation Model (DEM) is a representation of the terrain using point elevation information where each grid point represents the approximate elevation of the centre of the corresponding grid cell. A DEM was required for the production of the climate surfaces and for the estimation of slope and aspect corrected radiation grids.

The GEODATA 9 Second DEM Version 2 is a grid of elevation points covering the whole of Australia with a grid spacing of 9 seconds in longitude and latitude (~250 m resolution) in the GDA94 coordinate system. Version 2 is based on the ANUDEM 5.0 elevation gridding program developed by the Centre for Resource and Environmental Studies (CRES) at the Australian National University.

The ANUDEM program has been designed to produce accurate digital elevation models with sensible drainage properties from point elevations, stream lines, contour lines and cliff lines (Hutchinson 2000). The major data sources for the product were revised national spot height elevation data taken from 1:100,000 scale topographic mapping and revised river information from 1:250,000 scale topographic mapping. These datasets are components of AUSLIG's GEODATA TOPO-250K digital map product (AUSLIG 1994). All revisions to the source data were made by CRES. The data was augmented by national trigonometric data supplied by AUSLIG from the National Geodetic Data Base (NGDB).



**Figure 3. The Australian DEM.**

The source elevation data has a standard deviation of around 7.5 m. Errors in the DEM are closely related to terrain complexity. Theoretical estimates and tests of the DEM against trigonometric data distributed evenly across the continent indicate that the standard elevation error of the DEM varies between about 7.5 m and 20 m for most of the continent. Standard errors are larger in highland areas with steep and complex terrain where the largest errors can exceed 200 metres (Hutchinson *et. al.* 2000).

The density of source data points used to create the grid and the horizontal resolution of the grid warrant that the final grid be considered as having a scale of approximately 1:250,000. This makes the DEM useful for national, statewide and regional applications.

A bilinear interpolation technique was used to resample the 250 m elevation grid to produce data at a 1 km resolution to derive subsequent monthly productivity index map grids.

### 2.2.3 Station Dictionary

The task of spatial interpolation is made difficult by a number of factors. Climate data are often sparse, measured for a limited duration and of varying quality. More significantly, the density of the data may be much less than the resolution of the required interpolated grid. In spite of this, the size of the available data set may be sufficient to cause computational difficulties, particularly when dealing with continental or global data sets.

The iterative process of ‘fitting’ the statistical surface demonstrates a vitally important characteristic of the spline technique. Verification of climate data poses many challenges and one tool to assist in that verification is placing a climate station in its spatial context. The results of the spline interpolation provided feedback about the meteorological station database by providing summary statistics and the deviation for each station relative to the fitted surface. Examination of data residuals is a powerful aid for detecting and correcting the data errors.

To fit multi-variate climate surfaces, the values of the independent variables need to be known at the data points. Thus meteorological stations should be accurately located in position and elevation. The Bureau of Meteorology (BoM) provided a base station dictionary (23/05/01) with the data. The dictionary provided basic station information including:


- BoM site number – Australian mainland stations range from 001000 to 099999 where the first three digits represent the meteorological region.
- Latitude and longitude – in decimal degrees

- Elevation – in metres (both station and barometer heights may be given – station height is used for all climate variables as barometer heights are only used for pressure data)
- Station name – descriptive
- Start and finish year – gives years of operation of the station.

Climate data from 1968 to 2002 was required to produce the 1970 to 2002 plant productivity map grids for the NCAS, stations with no climate data for this period were removed from the data base. This version of the BoM dictionary was compared with both the CRES (1998) and AGO (2001) dictionary. Any station with a difference greater than 0.1° in position or 50 m in elevation was checked. The BoM base dictionary is updated monthly and comparisons were done in May 2001, September 2001, January 2002 and August 2002.

Stations which showed up as high residuals in the production of the climate surface were checked for correct locations. The final station dictionary which was used consisted of 11,714 stations, 2,511 of which were checked/geo-coded using the digital maps of Australia and the 9 second DEM. The digital maps provided a complete coverage of Australia at the 1:100,000 scale and at 1:250,000. The 1:100,000 scale was used where possible but in areas where only dyelines were available the 1:250,000 maps were also used. The 9 second DEM was used as a general check on elevation. A screen capture of each station checked was also taken (see Figure 4).





#### 2.2.4 Long-Term Average Climate Variables

The production of a productivity index map for Australia required a number of climate variables. Long-term average climate surfaces from the ANUCLIM software package were used for the climate variables available (Table 1). The ANUSPLIN software package and methods described in the previous section were used to produce a gridded set of long-term means for average temperature and number of frost days (Table 2).

The concept of a 'long-term average' climate surface has problems associated with the assumption of stationarity of the time series. The earth's climate will probably never settle reliably into an equilibrium with average long-term behaviour. Climatic averages are best given for either the full length of the record or for a standard period (for example a set thirty year period). The long-term averages used to produce productivity index maps used the full length of the record although records prior to 1920 were not used.

**Table 1. ANUCLIM variables used in the plant productivity modelling.**

| ANUCLIM Climate Variable  | Data points (knots) | Fitted variables  | Order of spline | Transformation |
|---------------------------|---------------------|---|-----------------|----------------|
| Maximum temperature       | 1200                | 3 independent (long,lat,elev)                           | 2               | none           |
| Minimum temperature       | 1200                | 3 independent (long,lat,elev)                           | 2               | none           |
| Rainfall                  | 13000 (3500)        | 3 independent (long,lat,elev)                           | 2               | Square root    |
| 9am Dry bulb temperature  | 185                 | 3 independent (long,lat,elev)                           | 2               | none           |
| 9am Dew point temperature | 176                 | 3 independent (long,lat,elev)                           | 2               | none           |
| 3pm Dry bulb temperature  | 190                 | 3 independent (long,lat,elev)                           | 2               | none           |
| 3pm Dew point temperature | 183                 | 3 independent (long,lat,elev)                           | 2               | none           |
| Radiation with rainfall   | 45                  | 3 independent (long,lat,elev)<br>1 dependent (rainfall) | 2               | none           |

**Table 2. ANUSPLIN variables created for the long-term plant productivity modelling.**

| ANUSPLIN Climate Variable | Data points (knots) | Fitted variables              | Order of spline | Transformation |
|---------------------------|---------------------|-------------------------------|-----------------|----------------|
| Average temperature       | 1126                | 3 independent (long,lat,elev) | 2               | -              |
| Frost days (number)       | 700                 | 3 independent (long,lat,elev) | 2               | Square root    |

#### 2.2.4.1 Average Temperature

ANUSPLIN (SPLINA) was fitted to data created from averaging the BoM monthly minimum and maximum temperature data and then calculating long-term averages. Both a spline and a partial spline function were fitted to these long-term average data (1,126 stations). A second order spline function was fitted to this data using three independent variables (longitude, latitude and elevation). The iterative process meant that five surfaces were produced and the final surface coefficient file was used in the productivity modelling. Errors associated with these surfaces were approximately  $\pm 0.5^{\circ}\text{C}$ . Figure 5 shows the fitted monthly long-term continental average temperature ( $T_{avg}$ ).

A second order partial spline function fitted to two independent variables (longitude and latitude) and one independent variable (elevation) was used to check lapse rate values. The lapse rates for average temperature for the Australian continent ranged from  $4.9^{\circ}\text{C}$  in June to  $6.0^{\circ}\text{C}$  in December.

#### 2.2.4.2 Number of Frost Days

SPLINA was used to fit the average of monthly frost days from data provided by the BoM. A second order spline function using three independent variables (longitude, latitude and elevation) was used to create the frost surfaces. Values on the output grids which were greater than the number of days in the month were reduced to the number of days in the month and any values less than 0 were converted to 0. Errors associated with the surfaces ranged from  $\pm 0.2$  days in summer to  $\pm 2$  days in winter. Figure 5 shows the fitted continental long-term average number of frost days per month ( $F$ ).

#### 2.2.4.3 Radiation

Long-term radiation surfaces (with rainfall as a covariate,  $R_p$ ) from the ANUCLIM package were used for both the long-term and monthly productivity indices.

For the slope and aspect corrected solar radiation the ratio of diffuse ( $R_F$ ) to global radiation ( $R_G$ ) was derived for each station from the original BoM data. For each of the 32 BoM stations which had monthly measurements of direct and global radiation a long-term average was determined from the monthly




Figure 5. Long-term monthly average temperature.






Figure 6. Long-term average number of frost days per month.



data to calculate the ratio of diffuse to global radiation. A 2-dimensional spline (using longitude and latitude) was fitted to the data to give a spatial coverage of the ratio expressed as a percentage for each month. This ratio surface was then applied to the ANUCLIM long-term radiation ( $R_P$ ) surface to produce coverages of direct and diffuse radiation. By definition the effects of slope and aspect apply to direct solar radiation only.

$$R_D = R_P - (R_P * R_F / R_G) \quad (3)$$

The radiation calculator is a computer program written by ScienceSpeak in C<sup>+</sup> and is based on a spreadsheet by J.B. Moncrieff of Edinburgh University. The radiation calculator estimates the correction to direct radiation from location (latitude) and slope and aspect (calculated from the 9 second DEM). The calculator only uses the zenith-angle correction factor from the Moncrieff spreadsheet as transmissivity effects are assumed to be mainly incorporated in the measured flat-earth measurements. Output is a grid of percentage change to flat earth which is used to adjust the direct radiation grid. This adjusted direct radiation ( $R_A$ ) grid is then added to the diffuse radiation grid to give the corrected global radiation ( $R_C$ ) grids.

$$R_C = R_A + R_F \quad (4)$$

Where  $R_C$  is the corrected global radiation,  $R_A$  is the slope and aspect adjusted direct radiation and  $R_F$  is the diffuse radiation. Grids of net radiation were derived where:

$$\begin{aligned} R_N &= (R_C * 0.8) - 4 & \text{if } R_C > 5 \\ R_N &= 0 & \text{if } R_C \leq 5 \end{aligned} \quad (5)$$

#### 2.2.4.4 Vapour Pressure Deficit (VPD)

The air absorbs water vapour and as water vapour is a gas it exerts a pressure in addition to that of the air. The higher the air temperature the more water vapour can be absorbed. Saturation vapour pressure ( $e^\circ$ ) specifies the maximum amount of water vapour which air, at a given temperature, can hold. The saturation vapour pressure increases with increasing

air temperature. The air inside a green leaf, near the surfaces of the cells where water is evaporating, is very nearly always saturated.

Vapour pressure deficit (VPD) is the difference between the actual vapour pressure and the saturation vapour pressure. For water vapour, VPD is normally in the range 0.1 kPa (very humid) to 3 kPa (very dry air), or 1 to 30 mbar. A low VPD means a high air humidity, and vice-versa. VPD indicates the 'drying effect' of the air, the higher the VPD the stronger the drying effect, so the stronger the driving force on transpiration.

The ANUCLIM program has long-term average coverages for 9am and 3pm dry bulb temperature (the ambient air temperature) and the 9am and 3pm dew point temperature (the temperature to which air must be cooled if the moisture present is to reach saturation). VPD was calculated by averaging the 9am and 3pm difference between the saturation vapour pressure for the dry-bulb temperature and the saturation vapour pressure for the dew point temperature.

$$VPD = ((e^\circ(T_{dry9}) - e^\circ(T_{dew9})) + (e^\circ(T_{dry3}) - e^\circ(T_{dew3}))) / 2 \quad (6)$$

### 2.2.5 Monthly Average Climate Variables

The production of monthly and annual productivity indices for Australia from 1970 to 2002 required monthly climate surfaces from 1968 to 2002. Climate surfaces for all variables were created using the ANUSPLIN software package and data from the BoM (see Table 3). Long-term radiation data was used as data was not available for monthly radiation for the early part of the time period and data numbers were considered insufficient for the latter part of the time period.

The methods described in the previous section were used to produce a gridded set of maps for monthly and annual means for temperature (minimum, maximum and average), rainfall and number of frost days, from January 1968 to December 2002 (Table 3).

### 2.2.5.1 Average Temperature


Monthly average temperature ( $T_{avg}$ ) data was calculated by averaging the BoM monthly minimum and maximum temperature data at each station. Any month which had less than 15 days recorded measurements was removed from the data set and all years which did not have 12 months of data were

also removed. The average number of station numbers used for all years was 489.5 ranging from 346 stations in 1986 to 741 stations in 2002.

SPLINA was then fitted to the average temperature data set. Both a spline and a partial spline function was fitted to the data. A second order spline function using three independent variables

**Table 3. Monthly climate variables.**

| ANUSPLIN Climate Variable | Data points (knots) | Fitted variables              | Order of spline | Transformation |
|---------------------------|---------------------|-------------------------------|-----------------|----------------|
| Average temperature       | 346-741             | 3 independent (long,lat,elev) | 2               | none           |
| Maximum temperature       | 346-741             | 3 independent (long,lat,elev) | 2               | none           |
| Minimum temperature       | 363-733             | 3 independent (long,lat,elev) | 2               | none           |
| Rainfall                  | 4719-7080 (2000)    | 3 independent (long,lat,elev) | 2               | Square root    |
| Frost days                | 350-739             | 3 independent (long,lat,elev) | 2               | none           |



**Figure 7. Annual average temperature derived from monthly data grids for 1968-2002 and examples of low (1976), average (1982) and high (1996) temperature years.**

(longitude, latitude and elevation) was used to create the average temperature surfaces. The average error associated with the grids was approximately  $\pm 0.5^{\circ}\text{C}$ .

The continental annual average temperature ranged from  $20.75^{\circ}\text{C}$  in 1976 to  $22.37^{\circ}\text{C}$  in 1998 (see graph, Figure 7). The mean continental average temperature for the period 1968 to 2002 was  $21.64^{\circ}\text{C}$ . The graph in Figure 7 shows a rising trend of  $0.45^{\circ}\text{C}$  but most of the rise in temperature is due to the relatively cooler years of 1974 to 1978 with only a  $0.2^{\circ}\text{C}$  rising trend evident in the last twenty years.

## 2.2.5.2 Maximum Temperature

Monthly maximum temperature ( $T_{max}$ ) was fitted from data from the BoM monthly maximum temperature data set. Any month which had less than 15 days recorded measurements was removed from the data set and all years which did not have 12 months of data were also removed. The average number of station numbers used for all years was 489.5, ranging from 346 stations in 1986 to 741 stations in 2002.

SPLINA was then fitted to the average temperature data set. Both a spline and a partial spline function was fitted to the data. A second order spline function using three independent variables (longitude, latitude and elevation) was used to create the average temperature surfaces. The average error associated with the data grids ranged from 0.2 to  $0.75^{\circ}\text{C}$ .




Figure 8. Australian average maximum temperature ( $T_{max}$ ) and the 12 month running mean.

Figure 8 shows the variation in the Australian average maximum temperature and the 12 month running mean shows an overall increase ( $\sim 0.4^{\circ}\text{C}$ ) over the period January 1968 to December 2002. Continental monthly maximum temperatures had a mean value of  $28.53^{\circ}\text{C}$  and ranged from a low of

$18.9^{\circ}\text{C}$  in July 1968 to a high of  $36.7^{\circ}\text{C}$  in December 1972. Figure 9 shows example maps for the high (July 2002 and December 1972) and low (July 1968 and December 1975) continental average values.




Figure 9. Maximum temperature grids for high and low continental average values for July and December.

### 2.2.5.3 Minimum Temperature

Monthly maximum temperature ( $T_{min}$ ) was fitted from data from the BoM monthly minimum temperature data set. Any month which had less than 15 days recorded measurements was removed from the data set and all years which did not have 12 months of data were also removed. The average number of station numbers used for all years was 516, ranging from 363 stations in 1986 to 733 stations in 2002.

Both spline and partial spline functions were fitted to the BoM data. A second order spline function using three independent variables (longitude, latitude and elevation) was used to create the minimum temperature surfaces. The average error associated with the maps ranged from 0.6 to 1.0°C.

Figure 10 shows the variation in the Australian average minimum temperature and the 12 month running mean shows an overall increase ( $\sim 0.5^{\circ}\text{C}$ ) over the period January 1968 to December 2002. Figure 11 shows the continental monthly minimum temperatures have a mean value of  $14.8^{\circ}\text{C}$  and ranged from a low of  $5.5^{\circ}\text{C}$  in July 1982 to a high of  $22.3^{\circ}\text{C}$  in February 1983. Figure 11 shows example maps for the high (February 1983 and July 1973) and low (February 1972 and July 1982) continental average values.





Figure 10. Australian average minimum temperature ( $T_{min}$ ) and the 12 month running mean.



**Figure 11. Minimum temperature grids for high and low continental average values for July and February.**

#### 2.2.5.4 Rainfall

Rainfall ( $P$ ) was fitted to the data from the BoM monthly rainfall data set. Any month in which had less than 20 days recorded measurements was removed from the data set and all years which did not have 12 months of data were also removed from the data set. The average number of station numbers used for all years was 5,972 ranging from 4,719 stations in 1994 to 7,080 stations in 1970.

A second order spline function using three independent variables (longitude, latitude and elevation) and a square root transformation was used to create the rainfall surfaces. The average

grid values ranged from 15.5 mm in November 1982 to 219 mm in January 1974. The error associated with the grids ranged from 14% to 30%, with the higher errors resulting from the lower than average value grids.

Figure 12 shows the Australian continental average monthly and annual rainfall. The mean value for the continental monthly average is 40 mm and ranges from a low of 7 mm in October 2002 to a high of 235 mm in January 1974. The mean annual rainfall was 485 mm and ranged from 327 mm in 2002 to 770 mm in 1974. Figure 13 shows annual rainfall maps for 1970 to 2002.




Figure 12. Australian continental average monthly rainfall.






Figure 13. Annual rainfall 1970–2002.



2.2.5.5 Frost Days

The number of frost days per month (*F*) was fitted to the data from the BoM monthly data set. Any month which had less than 20 days recorded measurements was removed from the data set and all years which did not have 12 months of data were also removed. The average number of station numbers used for all years was 548, ranging from 350 stations in 2002 to 739 stations in 1973.

A second order spline function using three independent variables (longitude, latitude and elevation) was used to create the frost surfaces. Values on the output grids which were greater than the number of days in the month were reduced to the number of days in the month and any values less than 0 were converted to 0. The continental average annual number of frost days ranged from 21.1 days in 1982 to 6.1 days in 1991(Figures 14 and 15) with a sharp decline in the latter half of the period. The error associated with the data grids ranged from 0.01 days in the summer months to 3 days in the winter months, with the higher errors resulting from the higher than average value grids.





Figure 14. Australian continental average annual number of frost days.



**Figure 15. Example grids of low (1991, average = 6.1 days), medium (1987, average = 14.0 days) and high (1982, average = 21.1 days) number of days frost.**

#### 2.2.5.6 Radiation

As the BoM monthly data would not support the derivation of monthly surfaces, long-term averages were used. For monthly productivity it was decided that 'flat earth' estimates of global radiation would be used as slope and aspect correction at the 1 km scale was thought to be inappropriate. The radiation with rainfall surfaces from ANUCLIM were used. Grids of net radiation were derived where:

$$R_N = (R_p * 0.8) - 4 \quad \text{if } R_p > 5$$

$$R_N = 0 \quad \text{if } R_p \leq 5 \quad (7)$$

### 2.3 NORMALISED DIFFERENCE VEGETATION INDEX (NDVI)

The NDVI provides a crude estimate of vegetation health and a means of monitoring changes in vegetation over time and is one of the most common vegetation indices derived from remotely sensed data. NDVI is related to vegetation as healthy vegetation reflects very well in the near infrared part of the spectrum. Green leaves have a reflectance of 20 percent or less in the 0.5 to 0.7 micron range (green to red,  $\lambda_R$ ) and about 60 percent in the 0.7 to 1.3 micron range (near infra-red,  $\lambda_I$ ). The value is then normalised to the range  $-1 \leq \text{NDVI} \leq 1$  to

partially account for differences in illumination and surface slope but is undefined when  $\lambda_R$  and  $\lambda_I$  are zero. The NDVI is computed by the product of the ratio of the two electromagnetic wavelengths.

$$\text{NDVI} = (\lambda_I - \lambda_R) / (\lambda_I + \lambda_R) \quad (8)$$

The typical range is between about -0.1 ( $\lambda_I$  less than  $\lambda_R$  for a not very green area) to 0.6 (for a very green area). NDVI values vary with absorption of red light by plant chlorophyll and the reflection of infrared radiation by water-filled leaf cells. Because photosynthesis occurs in the green parts of plant material the NDVI is normally used to estimate green vegetation.

The Advanced Very High Resolution Radiometer (AVHRR) sensor collects imagery at a 1 km resolution that is then used to generate a NDVI product for Australia. The Environmental Resources Information Network (ERIN) of the Australian Government Department of Environment and Heritage .01 degree NDVI archive contains data from the National Oceanic and Atmospheric Administration (NOAA) Satellite 9, 11, 14 and 16. The NOAA 14 data is calibrated using the equations in 'Revised post-launch calibration of channels 1 and 2 of the AVHRR on board the NOAA-14 spacecraft' by Rao and Chen on the NOAA web site at

<http://orbit-net.nesdis.noaa.gov/>. Data from NOAAs 9, 11 and 16 are cross-calculated to NOAA 14 by assuming that the reflectance of the 10,000 pixels with the lowest standard deviation in the NOAA 14 dataset from each of 5 of Australian Deserts are stable.

The data for April to September 1994 obtained from NOAA 11 was very poor due to very low sun angles, thus large angular corrections. For this period, NOAA 9 NDVI images were calibrated and then composited into the NOAA 11 image data.

The images were cloud screened by masking out pixels that did not appear to be biologically consistent with the pixels in the images taken immediately before and after. The first stage of this process was performed by a filter that masks out pixels that show a large decrease followed by a large increase. This procedure was also designed to remove 'dropout' values and abnormally high values probably associated with low sun angles. This was followed by a manual (subjective) stage where suspect areas were compared on screen with adjacent images. The cloud masks were then adjusted as required. It should be noted that the screening procedure contained a degree of subjectivity and was not exhaustive and the cloud masks were expected to change slightly with further work (N. Fitzgerald pers. comm.).

NDVI values were available for the ten year period, 1991 to 2000. Long-term averages were derived from the data and used in both the long-term and monthly productivity indices to provide a comprehensive cloud free data set. Monthly and annual data were not usable because of the extent of cloud cover obscuring data signals over the shorter time spans.

The 28 day average NDVI (in BIL file format) ( $NDVI_b$ ) files supplied by ERIN had values which ranged from 1 to 256. NDVI values less than 50 were considered to be cloud and given a no data value of -99. For each month NDVI was then rescaled:

$$\begin{aligned} NDVI &= (NDVI_b - 50) / 200 & \text{if } NDVI_b > 50 \\ NDVI &= -99 & \text{if } NDVI_b \leq 50 \end{aligned} \quad (9)$$

The satellite-derived NDVI data represents the photosynthetic capacity of all vegetation within a 1 km pixel for a given month, correlated with the fraction of photosynthetically active radiation (PAR) absorbed ( $fPAR$ ). Sellers (1985, 1987) derived an important relationship between leaf area index (LAI), absorbed photosynthetically active radiation (APAR) and NDVI. This research found that under specified canopy properties APAR was linearly related to NDVI and curvilinear related to LAI. LAI is usually positively correlated to an increase in the difference between  $\lambda_l$  and  $\lambda_R$  radiation. This relationship has been shown to hold generally over a number of different biomes. LAI can be calculated from:

$$LAI = \ln ( 1 - (NDVI * 1.0611) + 0.3431 ) / - 0.5 \quad (10)$$

### 3. CALCULATION OF PLANT PRODUCTIVITY

Monthly values of environmental constraints on productivity indices were expressed by modifiers calculated from the temperature, vapour pressure deficit (*VPD*) of the atmosphere, soil water deficit and number of frost days.

#### 3.1 TEMPERATURE MODIFIER

The growth of any plant species is limited by temperatures outside the optimum range for that species. A general assumption is made that, in any particular region, the plants are well-adapted to the existing temperature range.  $T_{low}$  was set to  $\frac{1}{2}$  the minimum temperature of the coldest month if the minimum temperature of the coldest month was greater than or equal to 0. If the minimum temperature of the coldest month was less than 0,  $T_{low}$  was set to  $\frac{1}{2}$  the minimum temperature of the coldest month plus the minimum temperature of the coldest month.  $T_{high}$  was set to 5°C above the maximum temperature of the hottest month of the

year and  $T_{opt}$  as equal to the average of  $T_{low}$  and  $T_{high}$ .  $T_{avg}$  was the average temperature described in Sections 2.2.4.1 and 2.2.5.1. The equation describing the effects of temperature is:

$$T_{mod} = \frac{T_{avg} - T_{low}}{T_{opt} - T_{low}} \left( \frac{T_{high} - T_{avg}}{T_{high} - T_{opt}} \right) \quad (11)$$

$T_{low}$  represents the monthly average temperature below which plant growth stops;  $T_{high}$  is the monthly average temperature above which plant growth stops and  $T_{opt}$  is the optimum temperature for growth. Minimum and maximum temperature data were derived from the ESOCLIM and ANUSPLIN packages and  $T_{avg}$ ,  $T_{high}$ ,  $T_{low}$  and  $T_{opt}$  were derived from these values. The temperature modifier ( $T_{mod}$ ) was 1 when  $T_{avg} = T_{opt}$ . Equation 10 gives a hyperbolic response curve, with  $T_{mod} = 0$  when  $T_{avg} = T_{low}$  or  $T_{high}$ . Consequently  $T_{mod}$  generally had relatively small effects on the calculation of plant productivity. Figure 16 shows the spatial temperature modifier equations for January and July long-term averages.




Figure 16. Temperature modifier spatial equations (from Equation 11) for January and July long-term averages.

### 3.2 FROST MODIFIER

A frost modifier ( $F_{mod}$ ) was applied, using the simple assumption that frost temporarily inactivates the photosynthetic mechanism in foliage, so there is no growth on a frost day. The modifier is simply one minus of number of frost days per month ( $F$ ) divided by the number of days in the month (Figure 17).

$$F_{mod} = (1 - F / \text{Days in month}) \quad (12)$$

### 3.3 VAPOUR PRESSURE DEFICIT (VPD) MODIFIER

The  $VPD$  modifier is important because plants respond to the  $VPD$  rather than relative humidity and the  $VPD$  provides the driving force for evaporation of water from the leaves. There is a gradient of water vapour from the leaves to the air because the air in the leaf is saturated whereas the air outside is drier. The larger the  $VPD$  the steeper the gradient. The  $VPD$  acting on stomatal, and hence canopy, conductance. The equation used is:

$$VPD_{mod} = \text{Exp}(-0.05 * VPD) \quad (13)$$

Figure 18 shows the  $VPD$  modifier spatial equations for the January and July long-term averages. This modifier essentially acts as a control on the rate of water loss; it is conditional upon soil water balance (see over page).





Figure 17. The frost modifier equation for the July long-term average.



**Figure 18. VPD modifier spatial equations for January and July long-term averages.**

### 3.4 SOIL WATER CONTENT ( $S_W$ )

Soil water content ( $S_W$ ) is derived from water balance calculations, which take into account the maximum soil water holding capacity ( $S_C$ ) in the root zone of plants. Plant water use, transpiration ( $Trans$ ), is calculated from the equation for equilibrium evaporation ( $E_{eq}$ , see Landsberg and Gower 1997; p 79), modified by feed-back from current soil water content, and a conventional water balance equation:

$$E_{eq} = ((0.67 * R_N * (1 - 0.05)) / 2.47) * \text{days in month} \quad (14)$$

Where  $E_{eq}$  is the equilibrium evaporation and  $R_N$  is the net radiation (see Equation 7)

Monthly transpiration ( $Trans$ ) was calculated as:

$$Trans(j) = E_{eq}(j) * S_{Wmod}(j-1) \quad (15)$$

Where  $Trans(j)$  is the transpiration for the month ( $j$ ),  $E_{eq}(j)$  is the equilibrium evaporation for the month and  $S_{Wmod}(j-1)$  is the soil water modifier (see Equation 20) for the previous month (Figure 20).

Rainfall interception ( $I$ ) is derived from the leaf area index (LAI) and is defined as:

$$I = 0.05 + (0.02 * LAI) \quad (16)$$

Australian continental rainfall interception ( $I$ ) values were derived for long-term averages only as monthly data was not available for the period 1970 to 2002. Figure 21 shows the long-term average  $I$  for January.






Figure 19. Spatial equation for equilibrium evaporation, January 1995.




Figure 20. Spatial equation for transpiration, January 1995.




Figure 21. Spatial equation for interception, January long term average.

The conventional water balance ( $W$ ) equation is derived from the rainfall ( $P$ ), the rainfall interception ( $I$ ) and transpiration ( $Trans$ ) and is given by:

$$W = (P * (1 - I)) - Trans \quad (17)$$

The soil water content ( $S_W$ ) is derived by adding the water balance (change in soil water content over the month) ( $W$ ) to the previous months soil water content (Equation 18, Figure 23). The soil water content is reduced to the soil water holding capacity if it is greater than soil holding water capacity. Negative soil water content values are set to zero.

$$S_W(j) = S_W(j-1) + W \quad (18)$$




Figure 22. Spatial equation for water balance, January 1995.




Figure 23. Spatial equation for soil water content, January 1995.



Initial (first month)  $S_W$  was taken as 0.75 of the soil water holding capacity ( $S_C$ ) and carries over from one time step to the next. The soil moisture calculation was sequenced so that initial  $S_W$  was set by running the water balance model for two years prior to initiating the productivity model. For the January 1970 to December 2002 monthly analysis, the  $S_W$  calculation sequence was initialised using the 1968 and 1969 climate data.


The soil water modifier ( $S_{Wmod}$ ) (Equation 20) was calculated from the moisture ratio ( $M$ ), which is the  $S_W$  divided by  $S_C$ . The equation describes the variable effect of  $M$  across the range from wet soil ( $M \approx 1$ ) to dry soil ( $M \approx 0$ ).

$$M = S_W / S_C \quad (19)$$

The soil water and  $VPD$  modifiers are not multiplicative; the lowest one applies. The argument is that if plant growth (conversion of radiant energy into biomass) is limited more by  $VPD$  than soil water (i.e. if  $VPD_{mod} < S_{Wmod}$ ) then soil water is not a limiting factor, even if soil water content is relatively low. The converse applies; i.e. if  $S_{Wmod} < VPD_{mod}$ , soil water is the limiting factor.

$$S_{Wmod} = 1 / (1 + ((1 - M) / 0.6)^7) \text{ if } 1 / (1 + ((1 - M) / 0.6)^7) < VPD_{mod}$$

$$S_{Wmod} = VPD_{mod} \text{ if } 1 / (1 + ((1 - M) / 0.6)^7) > VPD_{mod} \quad (20)$$



**Figure 24. Spatial equation for moisture ratio, January 1995.**




Figure 25. Spatial equation for soil water modifier, January 1995.

### 3.5 ABSORBED PHOTOSYNTHETICALLY ACTIVE RADIATION (APAR)


In the original version of the 3-PG model (following Landsberg and Waring 1997), APAR is multiplied by a factor that converts it to a biomass accumulation equivalent. This, in effect, amalgamates two physiological processes: the conversion of absorbed  $\text{CO}_2$  into initial carbon products (gross primary production) and the loss of a proportion of those products by respiration to give net primary productivity. The value of the conversion factor ( $\epsilon$ ,  $\text{gm C MJ}^{-1}$  APAR) used was obtained from the literature (Potter *et. al.* 1993, Ruimey *et. al.* 1994, Landsberg *et. al.* 1997). There is significant variation in  $\epsilon$  values, but no clear pattern in relation to plant type, so a 'best estimate' value of  $1.25 \text{ gm C MJ}^{-1}$  APAR was used. In the productivity model the net primary productivity equivalent was taken as an index of 'plant productivity' and was

not used to infer an absolute mass increase. Used in this way the conversion factor was not critical as it had only a linear (direct multiplicative) effect on the index.

Absorbed photosynthetically active radiation (APAR) was estimated from global solar radiation, interpolated from station data provided by the BoM, and from a linear relation with the satellite-derived NDVI which represents the photosynthetic capacity of all vegetation within each 1 km pixel and is correlated with the fraction of PAR absorbed ( $f\text{PAR}$ ).

APAR is calculated as half the amount of short-wave (global) incoming radiation ( $R_p$ ) absorbed by plant canopies:

$$\text{APAR} = R_p * 0.5 * (1 - \text{Exp}(-0.5 * \text{LAI})) \times \text{days in month} \quad (21)$$




**Figure 26. Long-term average APAR for January and July.**

Where LAI is the Leaf Area Index and the coefficient 0.5 is a general value for the extinction coefficient. In a slope and aspect corrected form of the productivity model radiation with rainfall ( $R_P$ ) is replaced by slope and aspect corrected radiation ( $R_C$ ). APAR was calculated for long-term means as the available radiation and LAI data were long-term averages.

### 3.6 PLANT PRODUCTIVITY INDEX

Productivity is reduced by modifiers reflecting non-optimal nutrition, soil water status, temperature and atmospheric vapour pressure deficits. Modifiers are dimensionless factors with values between 0 (complete restriction of growth) and 1 (no limitation). Modifiers used in this way are discussed by Landsberg (1986), McMurtrie *et. al.* (1994) and Landsberg and Waring (1997).



**Figure 27. Spatial equations for productivity indices, January and July long-term averages.**

Monthly values of plant productivity index (Figure 27) are expressed by APAR multiplied by the modifiers calculated from soil water balance, temperature, number of frost days and soil nutrient values.


$$\text{Productivity Index} = \text{APAR} * T_{mod} * S_N * S_{Wmod} * 0.01 * F_{mod} \quad (22)$$

Australia's annual plant productivity is the sum of the 12 monthly values for the year of interest

$$\text{Productivity Index}_{\text{ann}} = \sum_{j=1}^{j=12} \text{Productivity Index}_j \quad (23)$$

where  $j$  is the month. Figure 28 shows the long-term annual average productivity index for Australia and Figure 29 shows the annual productivity indices for 1970 to 2002.

A comparison of the annual productivity index maps in Figure 29 and the annual rainfall maps (Figure 13) shows the dominance of rainfall in the estimation of plant productivity.



**Figure 28. Derived long-term average plant productivity index for Australia.**






Figure 29. Derived annual continental productivity indices 1970-2002.

#### 4. SELECT REFERENCES

- AUSLIG 1994. *GEODATA TOPO-250K Data User Guide, Version 1 Data, Edition 2*. Australian Surveying & Land Information Group, Commonwealth Department of Administrative Services, Canberra, Australia.
- Bates, D. & Wahba, G. 1982. *Computational methods for generalised cross-validation with large data sets*. In Baker, C. and Miller, G. (eds), *Treatment of Integral Equations by Numerical Methods*, Academic Press, London.
- Eldén L. 1984. *A note on the computation of the generalized cross-validation function for ill-conditioned least squares problems*. BIT 24:467-472.
- Hillel, D. 1980. *Applications of Soil Physics*. Academic Press, New York.
- Hutchinson, M.F. 2001. *ANUSPLIN User Guide Version 4.2*. Centre for Resource and Environmental Studies, Australian National University, Canberra.
- Hutchinson, M.F. 2000. *ANUDEM Software*. Centre for Resource and Environmental Studies, Australian National University, Canberra. <http://cres.anu.edu.au/outputs/software.html>
- Hutchinson, M.F. 1998a. *Interpolation of rainfall data with thin plate smoothing splines: I. Two dimensional smoothing of data with short range correlation*. Journal of Geographic Information and Decision Analysis 2(2):153-167. [http://www.geodec.org/gida\\_4.htm](http://www.geodec.org/gida_4.htm)
- Hutchinson, M.F. 1998b. *Interpolation of rainfall data with thin plate smoothing splines: II. Analysis of topographic dependence*. Journal of Geographic Information and Decision Analysis 2(2):168-185. [http://www.geodec.org/gida\\_4.htm](http://www.geodec.org/gida_4.htm)
- Hutchinson, M.F. 1995. *Interpolating mean rainfall using thin plate smoothing splines*. International Journal of GIS 9:385-403.
- Hutchinson, M.F. 1993. *On thin plate splines and kriging*. In Tarter, M.E. & Lock, M.D. (eds), *Computing Science and Statistics*, Vol. 25. Interface Foundation of North America, University of California, Berkeley, pp. 55-62.
- Hutchinson, M.F. 1991a. *The application of thin plate smoothing splines to continent-wide data assimilation*. In Jasper, J.D. (ed), *BMRC Research Report No.27, Data Assimilation Systems*, Bureau of Meteorology, Melbourne, pp. 104-113.
- Hutchinson, M.F. 1991b. *Climatic analyses in data sparse regions*. In Muchow, R.C. & Bellamy, J.A. (eds), *Climatic Risk in Crop Production*, CAB International, Wallingford, pp. 55-71.
- Hutchinson, M.F. 1989. *A new objective method for spatial interpolation of meteorological variables from irregular networks applied to the estimation of monthly mean solar radiation, temperature, precipitation and windrun*. CSIRO Division of Water Resources Research Technical Memorandum 89/5: 95-104.
- Hutchinson, M.F. 1984. *A summary of some surface fitting and contouring programs for noisy data*. CSIRO Division of Land Use Research and Mathematics and Statistics, Consulting Report No. ACT 84/6, CSIRO, Canberra.
- Hutchinson, M.F. & Bischof, R.J. 1983. *A new method for estimating the spatial distribution of mean seasonal and annual rainfall applied to the Hunter Valley, New South Wales*. Australian Meteorological Magazine 31: 179-184.
- Hutchinson, M.F., Booth, T.H., McMahon, J.P. & Nix, H.A. 1984. *Estimating monthly mean values of daily total solar radiation for Australia*. Solar Energy 32: 277-290.
- Hutchinson, M.F. & Gessler, P.E. 1994. *Splines - more than just a smooth interpolator*. Geoderma 62: 45-67.

- Hutchinson, M.F., Houlder, D.J., Nix, H.A. & McMahon, J.P. 1999. *ANUCLIM User Guide, Version 5.0*. Centre for Resource and Environmental Studies, Australian National University, Canberra.
- Hutchinson, M.F., Nix, H.A. & McMahon, J.P. 1992. *Climate constraints on cropping systems*. In Pearson, C.J. (ed), *Field Crop Ecosystems of the World*, Elsevier, pp 37-58.
- Hutchinson, M.F., Nix, H.A., McMahon, J.P. & Ord, K.D. 1996. *The development of a topographic and climate database for Africa*. In *Proceedings of the Third International Conference/Workshop on Integrating GIS and Environmental Modelling*, NCGIA, Santa Barbara, California. Also on NCGIA Web Page.  
[http://www.ncgia.ucsb.edu/conf/SANTA\\_FE\\_C\\_D-ROM/main.html](http://www.ncgia.ucsb.edu/conf/SANTA_FE_C_D-ROM/main.html)
- Hutchinson, M.F., Stein, J.L. & Stein, J.A. 2000. *Derivation of nested catchments and sub-catchments for the Australian continent*. Centre for Resource and Environmental Studies, Australian National University, Canberra.  
<http://cres.anu.edu.au/outputs/programs.html>
- Hutchinson, M.F., Stein, J.A. & Stein, J.L. 2001. *Upgrade of the 9 Second Digital Elevation Model for Australia*. Centre for Resource and Environmental Studies, Australian National University, Canberra.  
<http://cres.anu.edu.au/outputs/programs.html>
- Kesteven, J.L. 1998. *A Spatial and Temporal Analysis of Australian Climate Fields*, PhD Thesis, ANU, November 1998.
- Kesteven, J.L. & Hutchinson, M.F. 1996. *Spatial modelling of climatic variables on a continental scale*. In *Proceedings of the Third International Conference/Workshop on Integrating GIS and Environmental Modelling*, NCGIA, Santa Barbara, California. Also on NCGIA Web Page.  
[http://www.ncgia.ucsb.edu/conf/SANTA\\_FE\\_C\\_D-ROM/main.html](http://www.ncgia.ucsb.edu/conf/SANTA_FE_C_D-ROM/main.html)
- Landsberg, J. J. 1986. *Physiological Ecology of Forest Production*. Academic Press, Sydney, 198 pp.
- Landsberg, J.J., & Gower, S.T. 1997. *Application of Physiological Ecology to Forest Management*. Academic Press, San Diego, California. 354p.
- Landsberg, J.J. & Waring, R.H. 1997. *A generalised model of forest productivity using simplified concepts of radiation-use efficiency, carbon balance and partitioning*. *Forest Ecology and Management* 95:209–228.
- Landsberg, J.J., Prince, S.D., Jarvis, P.G., McMurtrie, R.E., Luxmoore, R. & Medlyn, B.E. 1997. *Energy conversion and use in forests: An analysis of forest production in terms of radiation utilisation efficiency*. In *The Use of Remote Sensing in the Modeling of Forest Productivity at Scales from the Stand to the Globe*. (eds) Gholz, H.L., Nakane, K., and Shimoda, H. Kluwer Academic Publishers: Dordrecht.
- Laslett, G.M. 1994. *Kriging and splines: An empirical comparison of their predictive performance in some applications*. *Journal of the American Statistical Association* 89:391-409.
- Laslett, G.M., McBratney, A.B., Pahl, P.J. & Hutchinson, M.F. 1987. *Comparison of several spatial prediction methods for soil pH*. *Journal of Soil Science* 38, 325-341.
- McKenney, D.W., Hutchinson, M.F., Kesteven, J.L. & Venier, L.A. 2001. *Canada's plant hardiness zones revisited using modern climate interpolation techniques*. *Canadian Journal of Plant Sciences* 81: 129-143.
- McKenzie, N. J. & Hook, J. 1992. *Interpretations of the Atlas of Australian Soils*. Consulting Report to the Environmental Resources Information Network (ERIN). CSIRO Division of Soils Technical Report 94/1992.



- McMurtrie, R.E., Gholz, H.L., Linder, S. & Gower, S.T. 1994. *Climatic factors controlling the productivity of pine stands: a model-based analysis*. Ecological Bulletins, 43: 173-188.
- McKenzie, N.J., Jacquier, D.W., Ashton, L.J. & Cresswell, H.P. 2000. *Estimation of Soil Properties Using the Atlas of Australian Soils*, CSIRO Land and Water Technical Report 11/00.
- Northcote, K.H. with Beckmann, G.G., Bettenay, E., Churchward, H.M., Van Dijk, D.C., Dimmock, G.M., Hubble, G.D., Isbell, R.F., McArthur, W.M., Murtha, G.G., Nicolls, K.D., Paton, T.R., Thompson, C.H., Webb, A.A. & Wright, M.J. 1960-1968. *Atlas of Australian Soils*, Sheets 1 to 10. With explanatory data (CSIRO Aust. and Melbourne University Press: Melbourne).
- Northcote, K.H., Hubble, G.D., Isbell, R.F., Thompson, C. H. & Bettenay, E. 1975. *A Description of Australian Soils*. (CSIRO: Melbourne).
- Potter, C.S., Randerson, J.T., Field, C.B., Matson, P.A., Vitousek, P.M., Mooney, H.A. & Klooster, S.A. 1993. *Terrestrial ecosystem production: A process model based on global satellite and surface data*. Global Biogeochemical Cycles 7:811-841.
- Price, D.T., McKenney, D.W., Nalder, I.A., Hutchinson, M.F. & Kesteven, J.L. 2000. *A comparison of two statistical methods for spatial interpolation of Canadian monthly mean climate data*. Agricultural and Forest Meteorology 101: 81-94.
- Ruimey, A., Saugier, B. & Dedieu, G. 1994. *Methodology for the estimation of terrestrial net primary production from remotely sensed data*. J. Geophys. Res. 99:5263-5283.
- Sellers, P.J. 1985. *Canopy Reflectance, Photosynthesis and Transpiration*. International Journal of Remote Sensing, 68:1335-1372.
- Sellers, P.J. 1987. *Canopy Reflectance, Photosynthesis, and Transpiration 2. The role of Biophysics in the Linearity of Their Interdependence*. Remote Sensing of the Environment, 21:143-183.
- Stace, H.C.T., Hubble, G.D., Brewer, R., Northcote, K.H., Sleeman, J.R., Mulcahy, M.J. & Hallsworth, E.G. 1968. *A Handbook of Australian Soils*. Rellim Tech. Publ.: Glenside, S. A.
- Wahba G. 1979. *How to smooth curves and surfaces with splines and cross-validation*. Tech. Report 555. University of Wisconsin-Madison, Statistics Department.
- Wahba G. 1990. *Spline Models for Observational Data*. CBMS-NSF Regional Conference Series in Mathematics 59. SIAM, Philadelphia. Society for Industrial and Applied Mathematics, 169pp.

## **APPENDIX 1**

### **Preliminary Investigations: Spatial Estimation of Plant Productivity and Classification of Non-Commercial Vegetation Types**

**URS Australia**

**Joe Landsberg**

**Jenny Kesteven**

**July 2001**



# TABLE OF CONTENTS

|   | Page No. |
|---|----------|
| A1. Overview  | 43       |
| A2. Collation and Analysis of Growth Data – URS Australia                     | 44       |
| A2.1 Assembly of Datasets   | 44       |
| A2.2 Coding   | 45       |
| A2.3 Selection of Vegetation Types  | 45       |
| A2.4 Preparation of Growth Tables   | 45       |
| A2.5 Findings   | 45       |
| A2.5.1 Review of Publications   | 45       |
| A2.5.2 Comparability Issues and Adjustment Factors                            | 47       |
| A2.5.3 Forest Growth Data   | 47       |
| A2.6 Discussion   | 48       |
| A2.7 References   | 48       |
| A3. Spatial Estimation of Plant Productivity – Joe Landsberg & Jenny Kesteven | 50       |
| A3.1 Introduction   | 50       |
| A3.2 Background   | 50       |
| A3.3 Planning and Procedure   | 50       |
| A3.4 The Plant Productivity Model   | 52       |
| A3.5 Analysis of Plant Productivity by IBRA Regions                           | 52       |
| A3.6 Plant Productivity Values by IBRA Regions                                | 53       |
| A3.7 Spatial Variation in Plant Productivity within Regions                   | 55       |
| A3.8 Analysis of Plant Productivity by Carnahan Vegetation Classes            | 62       |
| A3.9 Conclusions  | 63       |
| A3.10 Follow Up Research  | 64       |
| A3.11 Relevant Citations  | 64       |
| A4. Forest Growth Data Tables – URS Australia                                 | 65       |

# LIST OF TABLES

|   | Page No. |
|---|----------|
| Table 1a. Growth data for sites in South Australia, Hassall & Associates (1999a).   | 66       |
| Table 1b. Growth data for sites in Victoria, Hassall & Associates (1998).           | 74       |
| Table 1c. Growth data for sites in Western Australia, Hassall & Associates (1999b). | 81       |
| Table 2a. Growth data for sites in South Australia, Wong <i>et. al.</i> (2000).     | 88       |
| Table 2b. Growth data for sites in Victoria, Wong <i>et. al.</i> (2000).            | 92       |

# LIST OF FIGURES

|   | Page No. |
|---|----------|
| <b>Figure 1.</b> Derived long-term average plant productivity index for Australia.  | 52       |
| <b>Figure 2.</b> Annual average Plant Productivity by IBRA regions.   | 53       |
| <b>Figure 3.</b> Monthly plant productivity index in four IBRA regions: South East Highlands (SEH); Jarrah Forest (JF); Darling River Plains (DRP); and Central Kimberly (CK).  | 54       |
| <b>Figure 4.</b> Monthly water balance in the four IBRA regions for which plant productivity data are shown in Figure 3. Negative values indicate the dryness of the area but note that in the calculation of water balances for the model soil water content cannot go below zero.   | 55       |
| <b>Figure 5a.</b> Change in the standard error of the mean productivity value with various numbers of sample points, up to 100, in the Wet Topics region. There was no indication that a minimum value had been reached. <b>b.</b> Spatial distribution of plant productivity across the Wet Tropics region. The data exhibit significant variation.  | 56       |
| <b>Figure 6a.</b> Change in the standard error of the mean productivity value with various numbers of sample points, up to 100, in the Tanami Desert region. There was no indication that a minimum value had been reached. <b>b.</b> Spatial distribution of mean annual plant productivity across the Tanami Desert. The data exhibit significant variation, with a marked peak in the low productivity range.  | 57       |
| <b>Figure 7a.</b> Change in the standard error of the mean productivity value with various numbers of sample points, up to 300, in the Jarrah Forest region. The curve indicates that 2-300 sample points would be required to give an accurate value of the mean. <b>b.</b> Spatial distribution of mean annual plant productivity across the Jarrah Forest region. The curve is skewed towards lower values.  | 58       |
| <b>Figure 8a.</b> Change in the standard error of the mean productivity value with various numbers of sample points, up to 300, in the South Eastern Highlands. The curve indicates that 2-300 sample points would be required to give an accurate value of the mean. <b>b.</b> Spatial distribution of mean annual plant productivity across the South Eastern Highlands region. The distribution is more uniform than in most other regions.            | 59       |
| <b>Figure 9a.</b> Change in the standard error of the mean productivity value with various numbers of sample points, up to 300, in the Darling River Plains. The curve indicates that at least 300 sample points would be required to give an accurate value of the mean. <b>b.</b> Spatial distribution of mean annual plant productivity across the Darling River Plains. The distribution is reasonably uniform, slightly skewed to the higher values. | 60       |
| <b>Figure 10a.</b> Change in the standard error of the mean productivity value with varying numbers of sample points, up to 300, in the Central Kimberley region. The curve has not reached stable values; at least 300 sample points would be required to give an accurate value of the mean. <b>b.</b> Spatial distribution of mean annual plant productivity across the Central Kimberley region. The distribution is approximately normal.            | 61       |
| <b>Figure 11.</b> The distribution of open woodland shaded by the plant productivity index value.   | 62       |
| <b>Figure 12.</b> Range of plant productivity values for the forest classes of present Carnahan vegetation classes.   | 63       |



## A1. OVERVIEW

The work reported here represents preliminary investigations toward developing the capability of the National Carbon Accounting System (NCAS) to estimate carbon sequestration in ‘non-commercial’ vegetation types, largely environmental plantings and native regrowth. This preliminary work provided guidance on the preferred framework and evolution of methods presented in the main body of this report.

Most of the collection of forest growth data and modelling activity has historically been focussed on commercial plantings which have higher productivity. Non-commercial plantings are important because of their associated benefits e.g. windbreaks, salinity control, erosion control, habitat and biodiversity, visual amenity. With the drive to more sustainable land management, there is a need for more of such plantings integrated with whole farm and whole region planning. Likewise, native regrowth following disturbance, e.g. fire and land clearing, is a common feature of the Australian landscape, and is a desirable form of revegetation. Carbon sequestration can likewise facilitate sound land management.

A relatively small, yet important, part of Australia’s National Greenhouse Gas Inventory is the carbon stored in environmental plantings and native regrowth; however there are few data available on which to develop Growth Tables, whereby empirical estimates may be made for specific projects or for regions. The NCAS is working to fill this gap, and this project is a means to that end. The NCAS has chosen the second of two broad options to do so:

1. Sample stands of environmental plantings and native regrowth systematically across Australia, along with associated environmental data, and develop the relationships needed to develop Growth Tables; or

2. Stratify the Australian landmass on the basis of plant productivity and use this to undertake a more-efficient stratified sampling and modelling of the growth potential within these strata.

This project comprises two steps:

1. Collate and analyse available data for the main types of environmental plantings and regrowth vegetation relevant to forming Growth Tables (as reported in Section A2 and A4); and, concurrently
2. Develop the possible methodologies for productivity stratification and vegetation classification; and apply the stratification methodology to produce the documented GIS layer over Australia (as reported in Section A3).

The original intent of this work was to develop a classification method for non-commercial plantings around potential carbon accumulation and to develop Growth Tables for regions of varying productivity throughout Australia. The aim was to classify vegetation into a manageable number of types of similar carbon accumulation patterns. This would provide default estimates of the carbon accumulation for a particular vegetation class in a particular region/location.

This was planned as a desktop study with 3 steps:

1. Draft classification of environmental plantings and native regrowth.
  - a) Define classes (classification)
  - b) Define productivity regions (stratification);
2. Consult with experts to refine the classification; and
3. Refine the classification.

The initial intention was to produce a series of 'look-up' Growth Tables for each of the types of non-commercial plantings, so that they could be applied in the relevant regions as defined in the first step of this project.

However, it was soon recognised that there were few data available concerning carbon accumulation patterns i.e. 'plant growth rates' of these vegetation types. Thus the classification would be extremely coarse and, on its own, would add little value. This conclusion led to the development of methods explored in Richards (2002) and described in detail in Richards (2001).

In light of this, an alternative approach of developing Growth Tables for typical vegetation types was pursued and is reported in the following Section A2 and data is tabulated in Section A4.

To guide data collection a list of about 40 vegetation types was compiled, but feedback was that this needed to be greatly expanded by incorporating height classes as well. Height class is generally used as a site index, that is, as a measure of site productivity. There are few experts in this field, and those who provided information or advice confirmed the paucity of this data and that it would be difficult to provide the necessary Growth Tables for each vegetation type.

Feedback received confirmed that the best way to proceed was to collate as much of the available growth information for native regrowth and environmental plantings that would allow potential development of growth models and likely carbon accumulation for particular regions. This sparse data could then be used as a basis for interpolation, guided by relative site productivity assessments, to fill gaps. The paucity of purely 'non-commercial' data led to the inclusion of data from some commercial plantings data.

## A2. COLLATION AND ANALYSIS OF GROWTH DATA

URS Australia

### A2.1 ASSEMBLY OF DATASETS

Datasets were sourced widely and because the data were sparse, all possibilities were considered: Fortech (now URS Forestry) files, CSIRO, Greening Australia, State agencies, Bureau of Rural Sciences (BRS), and relevant Universities.

Data sources included:

- The Australian Government Bushcare/Natural Heritage Trust (NHT) funded work undertaken by Hassalls 'Measurements of Plantations of Trees in [region] to Improve the Estimates of Carbon Sequestration in Low Rainfall Areas', for the following regions: South Australia, Victoria and WA;
- The publication commissioned by the AGO (NCAS Technical Report No. 25) by Grierson, Williams and Adams (2000) which listed sources of unpublished data; and
- The ABARE Trees on Farms survey, for leads as to what might be available. (Wilson *et al.* 1995)

A number of other published and unpublished reports and databases were reviewed. In most cases, these resources did not incorporate measured volumetric growth data and/or were not made available to URS Australia.

The surprising finding was just how little *relevant* data were available, even from native forestry agencies. This was after extensive consultations. The paucity of data led to the inclusion of some commercial plantings data. Because of this, careful consideration of the comparability and compatibility of data was undertaken before calculating Growth Tables.

It was clear that a large volume of related data exists, but there would be a need to establish its utility for this particular purpose. It is likely that further data will be identified and its utility should also be examined. Synthesis of such data has considerable cost benefit advantages over field sampling.

## A2.2 CODING

Information was gathered under data fields that would be useful for developing Growth Tables and interpreting the variation, including:

- Site information (ID #, name/location/coordinates, year established, area planted, species list);
- Biomass estimate (value, age of trees at time of measurement, type of estimate);
- Mean Annual Increment (MAI);
- Planting density (trees per hectare);
- Wood density (kg/m<sup>3</sup>);
- Site factors, e.g. rainfall, soil type/quality, topography, access to water, salinity;
- Husbandry, e.g. tubestock/direct seed, weed and pest control, fire, irrigation; and
- Source of information.

## A2.3 SELECTION OF VEGETATION TYPES

The selection of vegetation types for analysis of growth parameters was based on two considerations:

- The datasets that were made available – the best represented types can support the most robust Growth Tables and model calibrations; and
- Having a wide range of vegetation types, to give most scope for estimating the performance of other vegetation types by ‘interpolation’.

## A2.4 PREPARATION OF GROWTH TABLES

The revised plan was that the data for each species would be summarised statistically to understand the causes of variation. These have been presented in Section A4, Tables 1 and 2.

Having compiled the growth tables and examined the data, it became apparent that the paucity of data and the different methodologies used to arrive at this data, meant that comparisons of growth and the estimation of growth tables was inappropriate. Adjustments and assumptions could be made to lessen the impacts of these incompatibilities. Undertaking these adjustments was outside the scope of this project.

## A2.5 FINDINGS

### A2.5.1 Review of Publications

#### **Grierson *et. al.* (2000) Review of Unpublished Biomass-Related Information: Western Australia, South Australia, New South Wales and Queensland**

This presents data for mature vegetation or at least snap shots in time, e.g. Table 2. There are references to publications and datasets that may be able to supply growth rate data. The publication represents the first stage (identification of potentially usable datasets) of a two stage process. The second stage will provide comprehensive assembling and synthesis of this data.

Particular issues that were noted within Grierson *et. al.* (2000) include:

- Mallee-type species will likely need different estimates for single and multi-stemmed species;
- p. 7-8: Forestry SA, see Kiddle *et. al.* (1987), Boardman *et. al.* (1992) for growth curves of native species growth in plantations;
- p. 9: see reviews by Eamus *et. al.* (2000), Snowden *et. al.* (2000), Keith *et. al.* (2000), Burrows *et. al.* (2001) for mulga and other species;

- Root:shoot ratios - most decline with age, a few increase and a few have no clear trend (Table 4, p. 12). There is a wide range of mean values, even within the same genus. There is clear evidence of an environmental gradient and further work on the dominant factors of change, e.g. soil depth texture etc., nutrient and water availability that may permit modelled estimates;
- p. 21: see Todd (2000) for jarrah regeneration;
- p. 23: WA oil mallee growth data. Not really relevant as these are dense plantings. Might illuminate the shape of the growth curve;
- p. 24: see chenopod growth (Koonamore data, see Noble (1977) for recovery after over-grazing), also Sinclair (Botany Department, Adelaide University);
- p. 26: cypress pine growth rates quoted;
- MAI quoted for tropical dry forest and woodland;
- p. 29: wide variation in (fine) root biomass, similar to variation in leaf biomass; and
- p. 30: mountain ash data – summary values, see Grierson *et. al.* (1992).

**Hassall & Associates (1998, 1999a 1999b)**  
***Measurements of Plantations of Trees in [region] to Improve the Estimates of Carbon Sequestration in Low Rainfall Areas***

This set of three reports funded by Bushcare and the NHT aims to improve the knowledge on carbon sequestration estimates from tree planting in three States, Victoria, South Australia and Western Australia. Approximately a dozen sites were sampled in each state, incorporating a large range of climatic and physical variation and species mix.

Very diverse growth rates were observed throughout the regions studied. The diversity of the species mix, site characteristics and differing husbandry made comparisons between different

sites and regions difficult and few conclusions could be reached regarding the effect of different environmental conditions on biomass accumulation.

The approach used a constant wood density value for all tree ages in their estimation of volume, plus their use of total stem volume including bark as the base for estimating volume, make comparisons difficult with other major sources of data. Also, the surveying of multiple species plots necessitates the use of mean estimates for many of the numerical results, leading to further incompatibilities with other datasets.

Summaries of growth data for each site in the three States are provided in Section A4, Tables 1a-1c.

**Eamus *et. al.* (2000) *Review of Allometric Relationships for Estimating Woody Biomass: Queensland, the Northern Territory and Western Australia***

This report entailed the collection of primary volumetric data from a number of sites in Queensland and the Northern Territory. The report itself concentrates on constructing allometric equations linking this data with growth parameters such as height (DBH) and growth rate. The fundamental datasets from this work were not available for this study.

**Wong *et. al.* (2000) *Forecasting Growth of Key Agroforestry Species in South Eastern Australia***

This report, prepared for the Agroforestry Joint Venture Program, aimed to improve the reliability of forecasting tree growth and yield of key agroforestry, i.e. commercial species on cleared land across South Australia and Victoria. As part of this, primary data including volumetric information, was collected from long-established trials for six major *Eucalypt* species.

Despite focussing on commercial plantings, the Wong report was included in this study given the paucity of purely non-commercial data.

Summaries of growth data for each site in each State are provided in Section A4, Tables 2a-2b.

### Miscellaneous hardcopy and online databases and reports

A number of other published and unpublished reports and databases were reviewed. In most cases, these resources did not incorporate measured volumetric growth data and/or were not made available for this study. These included:

- *TREEDAT* – Stores the results of field trials of species, provenances and other sub-species taxa so they can be selectively retrieved to assist people to choose plants for nominated end uses and sites.
- *MPTDAT* – Provides information on the growth of Australian trees in various locations and assists staff of the Australian Tree Seed Centre.
- *Subtropical Tree Site Management (STSM)* – Provides a recording, analysis and delivery structure for data on tree growth, management techniques, and species selection in the subtropical region of Australia.
- *Optimising the Growth of Trees Planted on Farms* – This report summarises farm tree and shrub plantings on the Northwest slopes and Plains and Northern Tablelands of New South Wales.

The records available for *E. camaldulensis* in Hassall & Associates (1998, 1999a and 1999b) were one-off snap-shots only, hence no growth estimates for the one plot were able to be estimated. Further records for *E. camaldulensis* and other species are thought to be available in other inaccessible (and/or costly to obtain) databases and in the records of individuals and other operators in the industry.

#### A2.5.2 Comparability Issues and Adjustment Factors

---

For the data that was collected, there were several comparability and compatibility issues which prevented the generation and comparison of biomass growth curves. These are discussed below.

**Wood Density.** Biomass growth data can be estimated in Wong *et. al.* (2000) by applying estimates of wood density at various ages to the supplied volumetric growth data. This differs from Hassall & Associates (1998, 1999a and 1999b) where constant wood density values are assumed in its biomass projections. URS Australia initially attempted to estimate biomass values for Wong using wood density estimates provided from URS Forestry consultant staff, but subsequently abandoned this given the constant density assumptions used in Hassall & Associates and other biasing factors described below.

**Basis for Estimating Volume.** Hassall & Associates (1998, 1999a and 1999b) used total stem volume overbark as the basis for estimating volume. Wong's South Australian data was for total stem volume under bark, and the Victorian data was stem volume to 2 cm under bark. Even with similar wood density assumptions, the difference in base volume estimates prevents direct comparability of any resultant biomass estimates.

**Unclear Volume Parameters.** The height at which diameter measurements are made for estimating volume directly influences all subsequent calculations. The same applies for the shape of the tree bole. Hassall & Associates (1998, 1999a and 1999b) assumes a conical shape, while the Wong reporting is unclear about this assumption. Wong points out that the methodologies are not consistent by saying "*The growth data has been collected by numerous forestry organisations over the past 10 to 15 years. Tree measurements followed the standard measurational procedures for each State organisation...*"

The inconsistency in the current data collections highlights the merit of the NCAS in developing standard protocols for new data collection, making the protocols widely available, and requiring them to be used for all new data collection.

#### A2.5.3 Forest Growth Data

---

All data compiled in this study is summarised in the Growth Tables presented in Section A4.



## A2.6 DISCUSSION

The above is a modest harvest for a reasonable search effort. It is however a fair reflection of how sparse forest growth data are, and how difficult they are to access. There is doubtless more data around, effectively 'lost' in old files, and it would take more resources than were available (including financial purchase) to access them.

Adding to the problem of accessing data, is the current lack of suitable data. There is little data that includes biomass estimates. The focus on height parameters for the majority of vegetation and revegetation databases, rather than biomass, is one of the factors that limited the development of the Growth Tables. The incompatibilities of the data that do exist and are available is another major factor, as discussed above.

It would be possible to estimate biomass accumulation using allometric equations of height growth data, but this was outside the scope of the study and is in itself potentially subject to considerable error – see Eamus *et. al.* (2000).

In summary, the growth data contained in Section A4, Tables 1 and 2 provides a good representation of biomass accumulation levels for some major native species in differing husbandry, environmental and climatic conditions across southern Australia. It could be argued that with appropriate adjustments and suitably broad assumptions, growth tables could be estimated and compared over different datasets, as has been discussed in Section 3.2, albeit with certain caveats for compatibility and comparability.

## A2.7 REFERENCES

- Boardman, R., Thomas, M., Wood, R.R. & Ferguson, T. 1992. *Selected Plantations of Native Trees Growing in the Semi-Arid Zone of South Australia and Their Capacity to Sequester Atmospheric Carbon Dioxide*. Forest Research Branch, Central and Northern Region, Forestry SA, Adelaide.
- Burrows, W.H., Hoffmann, M.B., Compton, J.F., & Back, P.V. 2001. *Allometric Relationships and Community Biomass Stocks in White Cypress Pine (Callitris glaucophylla) and Associated Eucalypts of the Carnarvon Area - South Central Queensland (with additional data for Scrub Leopardwood - Flindersia dissosperma)*. National Carbon Accounting System Technical Report No. 33 Australian Greenhouse Office, Canberra, 16pp.
- Eamus, D., McGuinness, K. & Burrows, W. 2000. *Review of Allometric Relationships for Estimating Woody Biomass: Queensland, the Northern Territory and Western Australia*. National Carbon Accounting System Technical Report No. 5a, Australian Greenhouse Office, Canberra, pp56.
- Grierson, P.F., Adams, M.A. & Attiwill, P.M. 1992. *Estimates of Carbon Storage in the Above-ground Biomass of Victoria's Forests*. Aust. J. Botany: 631-640.
- Grierson, P., Williams, K. & Adams, M. 2000. *Review of Unpublished Biomass-Related Information: Western Australia, South Australia, New South Wales and Queensland*. National Carbon Accounting System Technical Report No. 25, Australian Greenhouse Office, Canberra, pp86.
- Hassall & Associates 1998. *Carbon sequestration in low rainfall areas: the measurement of plantations of trees in Victoria*. Environment Australia, Canberra, pp47.
- Hassall & Associates 1999a. *The measurement of plantations of trees in South Australia to improve the estimates of carbon sequestration in low rainfall areas*. Environment Australia, Canberra, pp96.

- Hassall & Associates 1999b. *The measurement of plantations of trees in Western Australia to improve the estimates of carbon sequestration in low rainfall areas*. Environment Australia, Canberra, p56.
- Keith, H., Barrett, D. & Keenan, R. 2000. *Review of Allometric Relationships for Estimating Woody Biomass: New South Wales, the Australian Capital Territory, Victoria, Tasmania and South Australia*. National Carbon Accounting System Technical Report No. 5b, Australian Greenhouse Office, Canberra, pp111.
- Kiddle, G. Boardman, R. & Van Der Sommen, F. 1987. *A Study of Growth and Characteristics of Woodlot and Amenity Tree Planting in Semi-Arid Rural South Australia*. Final Report, Forestry SA, Adelaide.
- Noble, I.R. 1977. *Long-term biomass dynamics in an arid chenopod shrub community at Koonamore, South Australia*. Australian Journal of Botany. 25:639-653.
- Richards, G.P. 2001. *The FullCAM Carbon Accounting Model: Development, Calibration and Implementation for the National Carbon Accounting System*. National Carbon Accounting System Technical Report No. 28, Australian Greenhouse Office, Canberra, pp49.
- Richards, G.P. 2002. *Biomass Estimation: Approaches for Assessment of Stocks and Stock Change*. National Carbon Accounting System Technical Report No. 27, Australian Greenhouse Office, Canberra, pp140.
- Snowdon, P., Eamus, D., Gibbons, P., Khanna, P., Keith, H., Raison, J. & Kirschbaum, M. 2000. *Synthesis of Allometrics, Review of Root Biomass and Design of Future Woody Biomass Sampling Strategies*. National Carbon Accounting System Technical Report No. 17, Australian Greenhouse Office, Canberra, pp114.
- Todd, M.C. 2000. *The Role of Nutrient Cycling in the Sustainability of Young Plant Communities on Mined Sites*. Ph. D. Thesis, University of Western Australia, Perth.
- Wilson, S., Whitham, J., Bhati, U.N. & Tran, Y. 1995. *Trees on Farms. Survey of trees on Australian Farms: 1993-94*. ABARE Research Report 95.7, ABARE, Canberra.
- Wong, J., Baker, T., Duncan, M., McGuire, D. & Bulman, P. 2000. *Forecasting growth of key agroforestry species in south-eastern Australia*. Rural Industries Research and Development Corporation, Canberra.

## A3. SPATIAL ESTIMATION OF PLANT PRODUCTIVITY

Joe Landsberg & Jenny Kesteven

### A3.1 INTRODUCTION

Preliminary plant productivity indices were calculated for the 80 IBRA regions of Australia (following Thackway and Cresswell, 1995) using a variant of the 3-PG (spatial) model based on the relationship between the photosynthetically active radiation absorbed by plant canopies (APAR) and the biomass productivity of those canopies. The model used a monthly time step. The factor converting APAR to biomass was reduced from the selected optimum value by modifiers dependent on soil fertility, atmospheric vapour pressure deficits, soil water content and temperature. The ESOCLIM package was used to generate climate surfaces for the country. Soil fertility and water holding capacity values were obtained from the Digital Soil Atlas of Australia. Leaf Area Index, essential for the calculation of APAR, was estimated from 10-year mean values of Normalised Difference Vegetation Indices, for 1 km pixels, for the whole country. Incoming short-wave radiation – and hence APAR – was corrected for slope and aspect using a Digital Elevation Map (DEM), but with 1 km pixels (which made no significant difference to the results). All analyses were carried out on estimates based on the assumption of flat terrain.

One hundred point plant productivity values, evenly spaced, were calculated for every IBRA region. Statistical analysis indicated that, as expected, there were significant differences in productivity between regions and, more importantly, that there was significant within-region variation. IBRA regions were not homogenous for biomass production. Part of the reason for this lay in the fact that most of them crossed climatic zones, and contained a variety of soils.

The result of this preliminary exercise was a map (and digital values) of plant productivity index, by months, for Australia. This provided general estimates of plant productivity in any region, and hence a basis for estimates of carbon sequestration and a framework for the analysis of carbon accumulation by various vegetation types. The products of the project could be tested empirically by statistical methods applied to all the measured biomass data available, or by modelling productivity in detail, for selected locations for which the appropriate soil and climatic data were provided as inputs.

### A3.2 BACKGROUND

URS Australia were contracted by the Australian Greenhouse Office (AGO) to produce a map of Australia showing the regional distribution of estimated plant productivity potential, in terms of carbon sequestration.

The task was to develop an information system which could be used to obtain reasonable estimates of probable carbon accumulation rates in any region of the country. These regional estimates would also provide the framework for more detailed research and analysis of carbon flows and carbon storage capacities.

It was recognised that a simple modelling approach would have to be adopted and Dr. Joe Landsberg was sub-contracted to do the work.

### A3.3 PLANNING AND PROCEDURE

Several meetings were held to determine the scope of the project, and the best approach to take. It was agreed that the project would be based around a simplified version of the 3-PG model (Landsberg and Waring, 1997), which has been extensively tested as a tool for the spatial estimation of plant productivity (see Coops *et. al.*, 2001). This is described in the following section.

It was agreed that the project would be based around an analysis of plant productivity potential by IBRA regions<sup>1</sup>. These provided defined areas,

classified in terms of clearly stated characteristics, that are well recognised and quite widely used. It was assumed that the IBRA regions were based on data likely to correlate with plant productivity.

The work was to be GIS-based, using a raster system, with sampling on the basis of pixels, or sample points, rather than polygons whose definitions involved assumptions that may not be justified. Initial discussions proposed the use of ~250 m pixels, since this was the scale of the Digital Elevation Map (DEM) available and it was proposed that the short-wave radiation incident on land surfaces should be corrected for slope and aspect. However, this involved an enormous amount of computing time so the calculations were initially done using 1 km pixels, compatible with the scale at which estimates of Leaf Area Index (LAI), derived from satellite data, were available for the country as a whole. Estimates of plant productivity using radiation not corrected for slope and aspect were referred to as the 'flat earth' estimates. Water bodies were screened out of the Normalised Difference Vegetation Index (NDVI) data.

The basic procedure was to sample 100 pixels per IBRA region, for each of which plant productivity would be calculated. This allowed statistical evaluation of the variation in plant productivity across a region – information that cannot be obtained from analysis of polygons. To determine whether 100 samples was enough, 300 samples from each of 4 regions were to be analysed to determine the number needed to reduce the standard error of the mean to the lowest possible (stable) level. Note that taking 100 samples per region, across regions that vary in size from 2,400 to 42,400 km<sup>2</sup>, means

that the sampling density will vary enormously. This was considered reasonable on the grounds that the larger regions were, virtually by definition, more homogeneous<sup>2</sup>.

It was agreed that the AGO would make available all the data bases it had, or had access to, and that it would be actively involved in the project through its Scientific Adviser, Dr Jenny Kesteven. That is, the project would be, in effect, collaborative with the AGO. In fact the AGO did most of the computing work involving data bases and GIS analysis, Joe Landsberg produced the productivity model and they worked together on de-bugging and analyses. The data bases required, and their sources, are described below.

The project took a great deal longer to complete than was initially planned. This was largely a result of delays in obtaining data; Australia-wide digital soil maps, including values for fertility and water holding capacity in each pixel, had to be obtained from ERIN (Environmental Resources Information Network, The Australian Government Department of Environment and Heritage) and interpretation provided by Dr Neil McKenzie (CSIRO Division of Land and Water; see McKenzie *et. al.* 2000). Climatic data for each month were produced using the ANUCLIM package; the number of frost days had to be calculated for each month for the whole country. The greatest delays were caused by the requirement for estimates of Leaf Area Index (LAI) for the whole country. These were initially provided by Dr Nicholas Coops (CSIRO Division of Forestry and Forest Products), using NDVI<sup>3</sup> data for 1995. However, the data had gaps and it was eventually decided that using data from a single year, even one

---

1 The IBRA (Interim Biogeographic Regionalisation for Australia) regions were developed to provide a framework for establishing priorities for delivering the National Reserves System Cooperative Program (NRSCP) (Thackway and Creswell, 1995). They were derived by compiling the best available data and information about each State and Territory including specialist field knowledge, published resource and environmental reports and are based on climate, lithology/geology, landform, vegetation, flora and fauna and land use. There are 80 IBRA regions within Australia.

2 In fact there was no reason to assume that plant productivity would be homogenous across IBRA regions (Richard Thackway; pers. comm.)

3 Normalised Digital Vegetation Index, obtained from satellite data. This is a ratio of the reflectances in the infra-red and near infrared wavebands. It actually provides a good measure of the amount of chlorophyll in the vegetation, but this generally correlates well with LAI. Empirical equations are used to calculate LAI from NDVI.




that conformed, in many respects, to average climatic conditions, was a dangerous procedure. There is always significant seasonal variation in weather conditions across Australia, and a year that is 'average' in one region may be far from average in others. LAI was therefore estimated from 10-year means of NDVI (from Landsat) for the whole country, provided by ERIN. Joe Landsberg produced the code (in Visual Basic) for a simplified, spatial version of the 3-PG model. This was implemented in the ARCINFO system by the AGO.

#### A3.4 THE PLANT PRODUCTIVITY MODEL

The long-term average plant productivity maps used in this preliminary study (Figure 1) were generated using the methods described in the main body of this report (Section 3.6, see page 33).

#### A3.5 ANALYSIS OF PLANT PRODUCTIVITY BY IBRA REGIONS

There were no significant differences between the 'flat earth' and slope and azimuth-corrected estimates of APAR when the analysis was based on 1 km pixels. All the analyses presented in this section are based on the 'flat earth' analysis.



**Figure 1. Derived long-term average plant productivity index for Australia.**



Two approaches were used to characterise plant productivity in the IBRA regions. As noted in Section A3.1, 100 points uniformly spaced, were sampled in each region. This provided the basis for statistical analysis of the variation in primary productivity across these areas, allowing assessment of whether they were homogeneous for this property. The characteristics used to calculate these point estimates of plant productivity were those of the pixels on which each sampling point fell. Subsequently the model was applied to the whole surface of Australia to give a complete map of plant productivity (Figure 1).

**A3.6 PLANT PRODUCTIVITY VALUES BY IBRA REGIONS**

Average annual plant productivity values for each of the 80 IBRA regions are given in Figure 2. As we would expect, the highest productivity is in the Wet Tropics (13.4), followed by the New South Wales North Coast (11.1), while the lowest productivity areas are the dry desert areas (Little Sandy and Simpson-Strezlecki deserts, each with 1.1).

Standard analysis of variance shows that, between IBRA regions, many of the productivity values were significantly different, an unsurprising result of little interest in itself. For comparisons between particular values the critical value for F-tests with 95% confidence (that differences are real) was 1.28.





Figure 2. Annual average Plant Productivity by IBRA regions.

The widest range of calculated productivity values was in the Wet Tropics (between 5.8 and 22.5; Standard Deviation 4.22), followed by the Central Mackay Coast (between 6.2 and 18.2; Standard Deviation 2.98). However, taking Standard Deviation as a ratio of the mean regional value, the largest variation was in the Victorian Volcanic Plain (mean productivity index = 5.9, ratio = 56 %), followed by the Tanami Desert (mean productivity index = 2.3, ratio = 47 %) and the Little Sandy Desert (mean productivity index = 1.1, ratio = 44 %).


Figure 3 shows the variation in average monthly plant productivity across four IBRA regions with markedly different characteristics and more than three-fold difference in mean annual productivity<sup>4</sup>.

The regions were (mean productivity index in parenthesis): Jarrah Forest (4.7); South East Highlands (7.5), Darling River Plains (5.3); and Central Kimberly (2.1). Figure 4 shows the monthly water balance for the same regions. Clearly, water balance, while it explains some of the monthly variation, does not account for all of it, particularly in the case of the South East Highlands, where maximum productivity was in the spring even though the water balance for plant growth in that period was decreasing. Productivity in early summer was reduced by the lack of water in the soil. Similarly, in the Jarrah Forest region, maximum productivity was in the late winter/spring period, based on water stored from the winter rainfall. Here too, productivity declines in the summer.



**Figure 3. Monthly plant productivity index in four IBRA regions: South East Highlands (SEH); Jarrah Forest (JF); Darling River Plains (DRP); and Central Kimberly (CK).**

<sup>4</sup> Maps of monthly plant productivity, water balance and the effects of temperature (temperature modifier) can be produced.





**Figure 4. Monthly water balance in the four IBRA regions for which plant productivity data are shown in Figure 3. Negative values indicate the dryness of the area but note that in the calculation of water balances for the model soil water content cannot go below zero.**

### A3.7 SPATIAL VARIATION IN PLANT PRODUCTIVITY WITHIN REGIONS

To assess the spatial variation of plant productivity across IBRA regions, we plotted the standard errors (SEs) of the mean values obtained with varying numbers of samples for the two regions (Wet Tropics and the Tanami) with the widest range in point values. In both cases (Figs. 5a, 6a) the SE decreased with increasing numbers of sample points, but showed little sign of reaching a steady value. The spatial distribution of plant productivity values in the two regions was quite different. In the Wet Tropics (Figure 5b) there was a fairly uniform distribution of values, with a peak towards the higher productivity end of the spectrum, while in the Tanami (Figure 6b) the distribution was skewed strongly left (towards the low productivity end of

the spectrum). There was little apparent relationship between the spatial distribution of plant productivity and the SE of the sample mean.

To evaluate the number of samples needed to minimise the SE of the mean, we took 300 sample points from each of the four regions (Jarrah Forest, South East Highlands, Darling River Plains, Central Kimberly) analysed earlier. The results are shown in Figures 7a, 8a, 9a and 10a, where the SEs are plotted against number of samples. The spatial distributions of plant productivity are presented in Figures 7b, 8b, 9b and 10b. The data indicate that, in most cases, at least 200 sample points would need to be analysed to minimise the SE of the mean value. Again, the spatial distribution of plant productivity values provides little guidance to the number of samples needed.



**Figure 5a. (top)** Change in the standard error of the mean productivity value with various numbers of sample points, up to 100, in the Wet Topics region. There was no indication that a minimum value had been reached. **b. (bottom)** Spatial distribution of plant productivity across the Wet Tropics region. The data exhibit significant variation.









Figure 6a. (top) Change in the standard error of the mean productivity value with various numbers of sample points, up to 100, in the Tanami Desert region. There was no indication that a minimum value had been reached. b. (bottom) Spatial distribution of mean annual plant productivity across the Tanami Desert. The data exhibit significant variation, with a marked peak in the low productivity range.









**Figure 7a. (top)** Change in the standard error of the mean productivity value with various numbers of sample points, up to 300, in the Jarrah Forest region. The curve indicates that 2-300 sample points would be required to give an accurate value of the mean. **b. (bottom)** Spatial distribution of mean annual plant productivity across the Jarrah Forest region. The curve is skewed towards lower values.



**Figure 8a. (top) Change in the standard error of the mean productivity value with various numbers of sample points, up to 300, in the South Eastern Highlands. The curve indicates that 2-300 sample points would be required to give an accurate value of the mean. b. (bottom) Spatial distribution of mean annual plant productivity across the South Eastern Highlands region. The distribution is more uniform than in most other regions.**



**Figure 9a. (top)** Change in the standard error of the mean productivity value with various numbers of sample points, up to 300, in the Darling River Plains. The curve indicates that at least 300 sample points would be required to give an accurate value of the mean. **b. (bottom)** Spatial distribution of mean annual plant productivity across the Darling River Plains. The distribution is reasonably uniform, slightly skewed to the higher values.



**Figure 10a. (top)** Change in the standard error of the mean productivity value with varying numbers of sample points, up to 300, in the Central Kimberley region. The curve has not reached stable values; at least 300 sample points would be required to give an accurate value of the mean. **b. (bottom)** Spatial distribution of mean annual plant productivity across the Central Kimberley region. The distribution is approximately normal.

### A3.8 ANALYSIS OF PLANT PRODUCTIVITY BY CARNAHAN VEGETATION CLASSES

An analysis of productivity by Carnahan vegetation (present) classes was done by overlaying the Carnahan vegetation on the productivity surface. While the distribution of productivity values for most of the Carnahan vegetation classes shows that

there was a preferred value or niche, the range for each vegetation class could be large. Figure 11 shows the distribution of open woodland as shaded by the plant productivity index value. Due to the wide distribution of the open woodland vegetation class it is not surprising that the productivity index values have a range from 1 to 25.




Figure 11. The distribution of open woodland shaded by the plant productivity index value.



Figure 12 clearly shows the wide range of plant productivity values for each vegetation class (e.g. open woodland ranges from a productivity index of 1 to 20, open forest ranges from 2 to 20). It also clearly shows that the overlap between the vegetation classes is large and would not allow for discrimination of classes by plant productivity. This will have implications for any stratification based on vegetation classes.

### A3.9 CONCLUSIONS

The mean productivity values obtained for IBRA regions appear to be ‘reasonable’, insofar as they correspond well to vegetation and climatic regions. The method of calculation used here is based on the premise that there is a linear relationship between the PAR absorbed by vegetation and production. This is not linear over short periods (e.g. hours-days), and the use of the simple Beer’s Law equation to calculate energy absorption by plant canopies could lead to significant errors over such periods. However when we consider time periods as long as a month, the assumption that biomass production is linearly related to biomass production is justified (Landsberg *et. al.*, 1997) and Beer’s Law provides good estimates of the energy absorbed.




Figure 12. Range of plant productivity values for the forest classes of present Carnahan vegetation classes.

The full version of 3-PG (spatial) model has been shown to provide good estimates of the productivity of forest types ranging from large, long-lived conifers (Coops *et. al.* 2001) to native forests in New South Wales (Tickle *et. al.* 2001) and fast-growing eucalypt plantations in Brazil (A.C. de Almeida and J.J.Landsberg; unpublished data). The model has also been tested against biomass measurements made in New Zealand re-growth scrub (White *et. al.*, 2000; Neal Scott, Landcare NZ; pers. comm.) and was found to perform well. These examples show that, given appropriate inputs, the basic model provides good estimates of productivity for specific locations, allowing confidence in the model as an estimator of wide-scale productivity values.

The spatial variation in plant productivity across the IBRA regions indicates that, whatever their value in terms of ecological regions and guides for the establishment of reserves, these regions are not homogeneous in terms of productivity. It is clear that the IBRA regions not only contain a variety of soil types – which would have been impossible to avoid – but, in most cases, they lay across the boundaries of several quite different climatic regions. This would explain some of the variance that needs to be able to be taken into account when making calculations for specific locations.

### A3.10 FOLLOW UP RESEARCH

Ultimately, a variant of the 3-PG model (as used here in these preliminary investigations, following Landsberg *et. al.* 1997) was used as the productivity model, in addition to the long-term average plant production index and top soil moisture deficits for the NCAS (as fully described in the main body of this report). Annual plant productivity indices were also subsequently used in development of the full carbon accounting model for the NCAS (see Richards 2001).

### A3.11 RELEVANT CITATIONS

- Coops, N.C., Waring, R.H., Brown, S.R. & Running, S.W. 2001. *Comparisons of Predictions of Net Primary Production and Seasonal Patterns in Water Use Derived with Two Forest Growth Models in Southwestern Oregon*. Ecological Modelling. 142:61-81.
- Landsberg, J.J. 1986. *Coupling of carbon, water and nutrient interactions in woody plant soil systems*. Tree Physiology 2.
- Landsberg, J.J. & Gower, S.T. 1997. *Applications of Physiological Ecology to Forest Management*. Academic Press: San Diego Press. 354pp.
- Landsberg, J.J. & Waring, R.H. 1997. *A generalised model of forest productivity using simplified concepts of radiation-use efficiency, carbon balance and partitioning*. Forest Ecology and Management 95:209–228.
- Landsberg, J.J., Prince, S.D., Jarvis, P.G., McMurtrie, R.E., Luxmoore, R. & Medlyn, B.E. 1997. *Energy conversion and use in forests: An analysis of forest production in terms of radiation utilisation efficiency*. In The Use of Remote Sensing in the Modeling of Forest Productivity at Scales from the Stand to the Globe. Eds: Gholz, H.L., Nakane, K., & Shimoda, H. Kluwer Academic Publishers: Dordrecht.
- McKenzie, N.J., Jacquier, D.W., Ashton, L.J. & Cresswell, H.P. 2000. *Estimation of Soil Properties Using the Atlas of Australian Soils*. CSIRO Land and Water Technical Report 11/00, February 2000.
- McMurtrie, R.E., Gholz, H.L., Linder, S. & Gower, S.T. 1994. *Climatic factors controlling the productivity of pine stands: A model-based analysis*. Ecological Bulletins 43:173–188.
- Potter, C.S., Randerson, J.T., Field, C.B., Matson, P.A., Vitousek, P.M., Mooney, H.A. & Klooster, S.A. 1993. *Terrestrial ecosystem production: A process model based on global satellite and surface data*. Global Biogeochemical Cycles 7:811–841.

Richards, G.P. 2001. *The FullCAM Carbon Accounting Model: Development, Calibration and Implementation for the National Carbon Accounting System*. National Carbon Accounting System Technical Report No. 28, Australian Greenhouse Office, Canberra, pp49.

Ruimey, A., Saugier, B. & Dedieu, G. 1994. *Methodology for the estimation of terrestrial net primary production from remotely sensed data*. J. Geophys. Res. 99:5263–5283.

Thackway, R. & Cresswell, I.D. (Eds). 1995. *An Interim Biogeographic Regionalisation for Australia: A Framework for Establishing the National System of Reserves*, Version 4.0. Australian Nature Conservation Agency, Canberra.  
[http://www.environment.gov.au/bg/nrs/ibraimcr/ibra\\_95/](http://www.environment.gov.au/bg/nrs/ibraimcr/ibra_95/)

Tickle, P.K., Coops, N.C. & Hafner, S.D. 2001. *Assessing Forest Productivity at Local Scales Across a Native Eucalypt Forest Using a Process Model, 3PG-SPATIAL*. Forest Ecology and Management. 152:275-291.

White, J.D., Coops, N.C. & Scott, N.A. 2000 *Predicting broad-scale forest and scrub biomass for New Zealand: Investigating the application of a physiologically-based model*. Ecological Modeling 131: 175–190.

## A4. FOREST GROWTH DATA

### URS Australia

Compiled forest growth data tables are presented overleaf.

A4 - Table 1a. Growth Data for Sites in South Australia, Hassall &amp; Associates (1999a)

| Site Information (South Australia, Hassall & Associates, 1999a) |   |      |              |   |  |     |                |                                   |                              |   |  |
|---|---|------|--------------|---|--|-----|----------------|-----------------------------------|------------------------------|---|--|
| ID  | Name, Location, Coordinates   | Est. | Area Planted | Species List & Purpose  | Biomass<br>t/ha                          | Age | MAI<br>t/ha/yr | Planting<br>Density<br>(trees/ha) | Volume<br>m <sup>3</sup> /ha | Site Factors<br>e.g. rainfall, soil type/quality,<br>topography, access to water, salinity  | Establishment & Stand Management<br>e.g. site preparation and prior landuse, tubestock/<br>direct seed, weeds/disease/pests, fire, irrigation  |
| 1   | Hynam Park,<br>Lower South East<br><br>Lat/long: 365713, 1405244      | 1990 | -            | <i>E. camaldulensis</i> ,<br><i>E. leucoxylon</i> ,<br><i>E. occidentalis</i><br><br>Shelterbelt                                | 88.03 <sup>1</sup><br>66.02 <sup>2</sup> | 8   | 11.00          | -                                 | N/A                          | Rainfall: Av 581 mm<br><br>Soil: Sandy loam over clay, pH 5.5<br>at surface to 7.0 at 50 cm<br><br>Topography: Flat plain<br><br>Water: Rainfed, adjacent to dairy farm<br>which may or may not be irrigated<br>(unspecified). Depth to<br>groundwater unknown. | Site preparation unknown. Previously used for<br>dairy grazing.<br><br>Direct seeding.<br><br>Selective thinnings for firewood.<br><br>Some stunted growth due to crowding out.<br>No other problems associated with<br>weeds/disease/pests, drought or fire.<br><br>No history of irrigation on the trees or in the<br>adjacent dairy pasture is specified. |
| 2   | Struan Fairlamb,<br>Lower South East<br><br>Lat/long: 360622,1404534  | 1993 | 0.7 ha       | <i>E. ovata</i> ,<br><i>E. occidentalis</i> ,<br><i>Allocasuarina<br/>verticillate</i><br><br>Purpose not<br>specified          | 4.76 <sup>1</sup><br>3.57 <sup>2</sup>   | 5   | 0.95           | 1,089                             | N/A                          | Rainfall: Av 558 mm<br><br>Soil: Black rendzina soils with an<br>impervious limestone layer at 30 cm,<br>pH 7.0 at surface to 8.5 at 30 cm<br><br>Topography: Flat plain<br><br>Water: Rainfed, depth to<br>groundwater unknown.                                | Site preparation unknown. Previously used for<br>sheep grazing.<br><br>Tubestock.<br><br>No thinnings.<br><br>Some incidence of pests (spitfire caterpillars)<br>and weeds. No treatment. No other problems.<br><br>Rainfed only.  |
| 3   | Struan Roadside,<br>Lower South East<br><br>Lat/long: 370620, 1404729 | 1990 | 0.9 ha       | <i>E. viminalis</i><br><i>E. leucoxylon</i> ,<br><i>Acacia<br/>pycnantha</i><br><br>Purpose not<br>specified                    | 96.19 <sup>1</sup><br>72.14 <sup>2</sup> | 8   | 12.02          | -                                 | N/A                          | Rainfall: Av 558 mm<br><br>Soil: Sandy clay loam over clay Terra<br>Rossa, pH 6.0 at surface to 7.0 at 30 cm<br><br>Topography: Not specified<br><br>Water: Rainfed, depth to<br>groundwater unknown.   | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>Thinnings history not specified.<br><br>Some incidence of insect attack and weeds,<br>but no damage. Some stunted growth due to<br>crowding out. No other problems.<br><br>Rainfed only.   |
| 4   | Mark Morris,<br>Kangaroo Island<br><br>Lat/long: 354911, 1374211      | 1993 | 2.5 ha       | <i>E. cneorifolia</i> ,<br><i>E. diversifolia</i> ,<br><i>E. cosmophylla</i><br><br>Groundwater<br>recharge/salinity<br>control | 18.29 <sup>1</sup><br>13.7 <sup>2</sup>  | 5   | 3.66           | -                                 | N/A                          | Rainfall: Av 532 mm<br><br>Soil: Clay loam over clay, pH 6.0 at<br>surface to 7.0 at 50 cm<br><br>Topography: Not specified<br><br>Water: Rainfed, depth to<br>groundwater unknown.   | Site preparation unknown. Previous use<br>mixed farming.<br><br>Direct seeding.<br><br>Thinnings history not specified.<br><br>No problems associated with weeds/disease/pests,<br>drought or fire.<br><br>Rainfed only.   |

| Site Information (South Australia, Hassall & Associates, 1999a) |   |      |              |  |   |     |             |                             |                           | Establishment & Stand Management  |   |
|---|---|------|--------------|--|---|-----|-------------|-----------------------------|---------------------------|---|---|
| ID  | Name, Location, Coordinates   | Est. | Area Planted | Species List & Purpose   | Biomass t/ha                            | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. site preparation and prior landuse, tubestock/direct seed, weeds/disease/pests, fire, irrigation   |
| 5   | Snellings Beach (tubestock)<br>Kangaroo Island<br>Lat/long: 354016, 1370360   | 1992 | 0.3 ha       | <i>E. creorifolia</i> ,<br><i>E. diversifolia</i> ,<br><i>Melaleuca lanceolata</i><br><br>Soil stabilisation, shelterbelt, amenity | 7.90 <sup>1</sup><br>5.93 <sup>2</sup>  | 6   | 1.32        | -                           | N/A                       | Rainfall: Av 655 mm<br><br>Soil: Clay loam over clay, pH 6.0 at surface to 6.5 at 50 cm<br><br>Topography: Not specified<br><br>Water: Rainfed only (following irrigation for first 12 months), depth to groundwater unknown. | Site preparation unknown.<br>Previously used for sheep grazing.<br><br>Tubestock.<br><br>Thinnings history not specified.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed, plus drip fed for initial 12 months.       |
| 6   | Snellings Beach (direct seed)<br>Kangaroo Island<br>Lat/long: 354015, 1370354 | 1992 | 0.3 ha       | <i>Allocasuarina verticillata</i><br><br>Shelterbelt, amenity  | 5.81 <sup>1</sup><br>4.36 <sup>2</sup>  | 6   | 0.97        | -                           | N/A                       | Rainfall: Av 655 mm<br><br>Soil: Clay loam over clay, pH 6.0 at surface to 6.5 at 50 cm<br><br>Topography: Rolling coastal hills<br><br>Water: Rainfed, depth to groundwater unknown<br><br>Aspect: S.                        | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>Thinnings history not specified.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |
| 7   | Magpie Creek, Mid North<br>Lat/Long: 334340, 1382353                          | 1992 | 1.5 ha       | <i>E. camaldulensis</i> ,<br><i>E. leucocydon</i> ,<br><i>E. odorata</i><br><br>Soil stabilisation                                 | 10.06 <sup>1</sup><br>7.55 <sup>2</sup> | 6   | 2.01        | 2,333                       | N/A                       | Rainfall: Av 408 mm<br><br>Soil: Clay loam over deep limestone, pH 9.0 at surface to 50 cm<br><br>Topography: Gently sloping hills<br><br>Water: Rainfed, depth to groundwater approx. 7m<br><br>Aspect: Unknown.             | Site preparation unknown. Previous use unknown.<br><br>Tubestock.<br><br>Thinnings history not specified. Replanting in 1996 to fill originally missed areas.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.   |
| 8   | Blyth-Brinkworth Corridor, Mid North<br>Lat/long: 334352, 1382514             | 1996 | 1.2 ha       | <i>E. odorata</i> ,<br><i>E. porosa</i> ,<br><i>E. calamifolia</i><br><br>Visual amenity   | 0.14 <sup>1</sup><br>0.11 <sup>2</sup>  | 2   | 0.07        | 916                         | N/A                       | Rainfall: Av 408 mm<br><br>Soil: Clay loam, pH 8.5 at surface to 50 cm<br><br>Topography: Flat plain<br><br>Water: Rainfed, depth to groundwater approx. 15m.   | Site preparation unknown. Previously used for mixed farming.<br><br>Direct seeding.<br><br>Thinnings history not specified.<br><br>Some problems with weeds. Potential impact on growth rates of trees. No treatment. No other problems.<br><br>Rainfed only. |



A4 - Table 1a. Growth Data for Sites in South Australia, Hassall &amp; Associates (1999a) continued.

| Site Information (South Australia, Hassall & Associates, 1999a) |   |      |              |   | Biomass                                | MAI | Planting           | Volume             | Site Factors  | Establishment & Stand Management  |  |
|---|---|------|--------------|---|--|-----|--------------------|--------------------|---|---|--|
| ID  | Name, Location, Coordinates   | Est. | Area Planted | Species List & Purpose  | t/ha                                   | Age | Density (trees/ha) | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity | e.g. site preparation and prior landuse, tubestock/direct seed, weeds/disease/pests, fire, irrigation   |  |
| 9   | Blyth, Mid North<br><br>Lat/long: 334957, 1382859                         | 1996 | 2.5 ha       | <i>E. odorata</i> ,<br><i>Allocasuarina verticillata</i> ,<br><i>Acacia pycnantha</i><br><br>Visual amenity           | 0.18 <sup>1</sup><br>0.14 <sup>2</sup> | 2   | 0.09               | 833                | N/A   | Rainfall: Av 427 mm<br><br>Soil: Clay loam, pH approx. 8.0<br><br>Topography: Flat plain<br><br>Water: Rainfed, depth to groundwater 10-15m.  | Site preparation unknown. Previously used as a railway corridor.<br><br>Direct seeding.<br><br>No Thinnings.<br><br>Significant weed infestation requiring treatment. No other problems.<br><br>Rainfed only.  |
| 10  | Mourilya (direct seed)<br>Eyre Peninsula<br><br>Lat/long: 343612, 1354336 | 1992 | 1 ha         | <i>E. cladocalyx</i> ,<br><i>Acacia pycnantha</i><br><i>Allocasuarina verticillata</i> ,<br><br>Purpose not specified | 6.34 <sup>1</sup><br>4.75 <sup>2</sup> | 6   | 1.06               | -                  | N/A   | Rainfall: Av 495 mm<br><br>Soil: Sandy loam, pH approx. 6.5<br><br>Topography: Gently sloping low hills<br><br>Water: Rainfed, depth to groundwater 10-15m<br><br>Aspect: S.                                  | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.         |
| 11  | Mourilya (tubestock)<br>Eyre Peninsula<br><br>Lat/long: 343453, 1354336   | 1994 | 2 ha         | <i>E. cladocalyx</i> ,<br><i>E. lansdowneana</i> var. <i>albopurpurea</i><br><i>E. leucoxydon</i><br><br>Shelterbelt  | 7.53 <sup>1</sup><br>5.64 <sup>2</sup> | 4   | 1.88               | -                  | N/A   | Rainfall: Av 424 mm<br><br>Soil: Sandy loam over clay, pH unknown<br><br>Topography: Gently sloping low hills<br><br>Water: Rainfed, depth to groundwater 5m<br><br>Aspect: S.                                | Site preparation unknown. Previously used for mixed farming.<br><br>Tubestock.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only. |
| 12  | Warrow, Eyre Peninsula<br><br>Lat/long: 341855, 1352840                   | 1992 | 2.5 ha       | <i>E. incrassata</i> ,<br><i>E. porosa</i> ,<br><i>Allocasuarina verticillata</i><br><br>Habitat                      | 5.35 <sup>1</sup><br>4.01 <sup>2</sup> | 6   | 0.89               | -                  | N/A   | Rainfall: Av 420 mm<br><br>Soil: Sand over limestone, pH 7.5 at surface to 8.5-9.0 at 30-40 cm<br><br>Topography: Flat plain - gentle slope<br><br>Water: Rainfed, depth to groundwater 3m<br><br>Aspect: SW. | Site preparation unknown. Previously used for grazing.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |

| Site Information (South Australia, Hassall & Associates, 1999a) |  |      |              |   |  |     |             |                             |                           | Establishment & Stand Management   |   |
|---|--|------|--------------|---|--|-----|-------------|-----------------------------|---------------------------|--|---|
| ID  | Name, Location, Coordinates  | Est. | Area Planted | Species List & Purpose  | Biomass t/ha                               | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity   | e.g. site preparation and prior landuse, tubestock/direct seed, weeds/disease/pests, fire, irrigation   |
| 13  | Jackson (direct seed)<br>Upper South East<br><br>Lat/long: 362251, 1401934 | 1990 | -            | <i>E. gracilis</i> ,<br><i>E. leucoxydon</i> ,<br><i>E. leptophylla</i><br><br>Purpose not specified  | 133.37 <sup>1</sup><br>100.03 <sup>2</sup> | 8   | 16.67       | -                           | N/A                       | Rainfall: Av 560 mm<br><br>Soil: Sand over shallow clay, pH 6.0 at surface to 7.0 at 50 cm<br><br>Topography: Rolling plains<br><br>Water: Rainfed, depth to groundwater unknown.                            | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |
| 14  | Jackson (tubestock)<br>Upper South East<br><br>Lat/long: 362248, 1401934   | 1988 | 0.6 ha       | <i>E. gracilis</i> ,<br><i>E. fasciculosa</i> ,<br><i>E. camaldulensis</i><br><br>Purpose not specified   | 15.21 <sup>1</sup><br>11.40 <sup>2</sup>   | 10  | 1.52        | -                           | N/A                       | Rainfall: Av 560 mm<br><br>Soil: Sand, low salinity, pH 6.0-6.5 at surface to 7.0-8.5 at 50 cm<br><br>Topography: Rolling plains<br><br>Water: Rainfed, depth to groundwater unknown.                        | Site preparation unknown. Previous use unknown.<br><br>Tubestock.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.   |
| 15  | Meningie-Coonalpyn Road, Upper South East<br><br>Lat/long: 354132, 1392646 | 1993 | -            | <i>E. leucoxydon</i> ,<br><i>Allocasuarina verticillata</i> ,<br><i>Allocasuarina muelleriana</i><br><br>Roadside shelterbelt, revegetation trial | 10.77 <sup>1</sup><br>8.08 <sup>2</sup>    | 5   | 2.15        | -                           | N/A                       | Rainfall: Av 468 mm<br><br>Soil: Non-wetting sand, low salinity, pH 8.0 at surface to 9.0 at 50 cm<br><br>Topography: Flat plain<br><br>Water: Rainfed, depth to groundwater unknown.                        | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |
| 16  | Del Fabbro, Upper South East<br><br>Lat/long: 363921, 1401708              | 194  | 10 ha        | <i>Melaleuca brevifolia</i><br><i>Melaleuca halmaturorum</i><br><i>E. leucoxydon</i><br><br>Groundwater recharge/salinity control                 | 6.89 <sup>1</sup><br>5.16 <sup>2</sup>     | 4   | 1.72        | -                           | N/A                       | Rainfall: Av 521 mm<br><br>Soil: Poor sands over clay, evidence of salinity, pH 6.0<br><br>Topography: Rolling plain<br><br>Water: Rainfed, plus shallow depth to watertable, particularly in winter months. | Site fenced off and sprayed for weeds prior to sowing. Previously used as sheep grazing.<br><br>Direct seeding.<br><br>No thinnings.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed, plus access to shallow groundwater. |

A4 - Table 1a. Growth Data for Sites in South Australia, Hassall &amp; Associates (1999a) continued.

| Site Information (South Australia, Hassall & Associates, 1999a) |                                       |      |              | Biomass   | MAI | Planting           | Volume             | Site Factors   | Establishment & Stand Management  |
|---|---------------------------------------|------|--------------|---|-----|--------------------|--------------------|--|---|
| ID  | Name, Location, Coordinates           | Est. | Area Planted | Species List & Purpose  | Age | Density (trees/ha) | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. site preparation and prior landuse, tubestock/direct seed, weeds/disease/pests, fire, irrigation   |
| 17  | Langhorne Creek Cemetery, Murraylands | 1992 | -            | <i>E. leucoxydon</i> ,<br><i>Acacia pycnantha</i> ,<br><i>Acacia brachybotrya</i>                                       | 6   | -                  | N/A                | Rainfall: Av 392 mm<br><br>Soil: Clay with increasing levels of limestone at 50 cm, pH 6.0-6.5 at surface to 7.5-8.5 at 50 cm<br><br>Topography: Low gently sloping hillside | Site preparation unknown. Previously used for dairy farming and cropping.<br><br>Direct seeding.<br><br>Thinnings of lower limbs and inferior saplings. |
|   | Lat/long: 351749, 1390312             |      |              | Seed collection   |     |                    |                    | Water: Rainfed, depth to watertable approx. 3m.  | No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed, plus possible access to groundwater.                                  |
|   |                                       |      |              |   |     |                    |                    |  |   |
| 18  | Hartley Dump, Murraylands             | 1992 | -            | <i>E. leucoxydon</i> ,<br><i>E. porosa</i> ,<br><i>Acacia calamifolia</i>   | 6   | -                  | N/A                | Rainfall: Av 392 mm<br><br>Soil: Clay with varying levels of limestone, pH 8.0 at surface to 9.0 at 30 cm<br><br>Topography: Gently rolling hills                            | Site preparation unknown. Adjoining rubbish dump/quarry.<br><br>Direct seeding.<br><br>Thinnings history unknown.                                       |
|   | Lat/long: 351111, 1390205             |      |              | Site rehabilitation   |     |                    |                    | Water: Rainfed, depth to watertable unknown.   | No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |
|   |                                       |      |              |   |     |                    |                    |  |   |
| 19  | Swanport, Murraylands                 | 1991 | -            | <i>E. leucoxydon</i> spp.<br><i>leucoxydon</i> ,<br><i>E. leucoxydon</i><br>spp. <i>stephenae</i> ,<br><i>E. porosa</i> | 7   | -                  | N/A                | Rainfall: Av 346 mm<br><br>Soil: Sand, low salinity, pH 7.0 at surface to 8.0-8.5 at 50 cm<br><br>Topography: Gently undulating  | Site preparation unknown. Previously used for agricultural purposes.<br><br>Tubestock.<br><br>Thinnings history unknown.                                |
|   | Lat/long: 350855, 1391842             |      |              | Purpose not specified   |     |                    |                    | Water: Rainfed, depth to watertable 7-10m<br><br>Aspect: W.  | Some problems associated with weeds and rabbits which may be affecting growth. No other problems.<br><br>Rainfed only.                                  |
|   |                                       |      |              |   |     |                    |                    |  |   |
| 20  | Hugh Longbottom, Yorke Peninsula      | 1992 | 0.9 ha       | <i>E. porosa</i> ,<br><i>Acacia pycnantha</i> ,<br><i>Acacia notabilis</i>  | 6   | -                  | N/A                | Rainfall: Av 504 mm<br><br>Soil: Clay-loam, pH unknown<br><br>Topography: Gently rolling plain   | Site preparation unknown. Previously used for cropping.<br><br>Direct seeding.<br><br>Thinnings history unknown.  |
|   | Lat/long: 342137, 1373235             |      |              | Shelterbelt   |     |                    |                    | Water: Rainfed, depth to watertable unknown.   | No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |
|   |                                       |      |              |   |     |                    |                    |  |   |

| Site Information (South Australia, Hassall & Associates, 1999a) |   |      |              |  |  |     |             |                             |                           | Establishment & Stand Management   |   |
|---|---|------|--------------|--|--|-----|-------------|-----------------------------|---------------------------|--|---|
| ID  | Name, Location, Coordinates   | Est. | Area Planted | Species List & Purpose   | Biomass t/ha                             | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity   | e.g. site preparation and prior landuse, tubestock/direct seed, weeds/disease/pests, fire, irrigation   |
| 21  | Maitland-Minlaton Road, Yorke Peninsula<br>Lat/long: 342637, 1373920      | 1994 | 0.6 ha       | <i>E. porosa</i> ,<br><i>Acacia spinescens</i> ,<br><i>Dodenaëa viscosa</i><br><br>Roadside revegetation trial | 2.10 <sup>1</sup><br>1.57 <sup>2</sup>   | 4   | 0.52        | -                           | N/A                       | Rainfall: Av 504 mm<br><br>Soil: Clay-loam, pH unknown<br><br>Topography: Gently rolling plain<br><br>Water: Rainfed, depth to watertable unknown.   | Site preparation unknown. Previous use not specified.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |
| 22  | Klopp, Yorke Peninsula<br>Lat/long: 342855, 1374520                       | 1990 | 0.27 ha      | <i>E. porosa</i> ,<br><i>E. socialis</i> ,<br><i>E. calycogona</i><br><br>Roadside planting                    | 22.56 <sup>1</sup><br>16.92 <sup>2</sup> | 8   | 2.82        | -                           | N/A                       | Rainfall: Av 414 mm<br><br>Soil: Sandy loam, pH 8.5 at surface to 9.0 at 50 cm<br><br>Topography: Rolling plains<br><br>Water: Rainfed, depth to watertable unknown but believed to seasonally be within root zone<br><br>Aspect: N. | Site preparation unknown. Previous use not specified.<br><br>Tubestock.<br><br>Thinnings history unknown.<br><br>Some weeds present but not affecting growth. No other problems associated with disease, pests, drought or fire.<br><br>Rainfed but with potential access to groundwater. |
| 23  | Mount Barker Tannery, Mount Lofty Ranges<br>Lat/long: 350332, 1385151     | 1992 | 12 ha        | <i>E. viminialis</i> ,<br><i>E. leucoxylon</i> ,<br><i>E. fasciculosa</i><br><br>Site rehabilitation, amenity  | 30.30 <sup>1</sup><br>22.72 <sup>2</sup> | 6   | 5.05        | -                           | N/A                       | Rainfall: Av 769 mm<br><br>Soil: Sandy loam over clay, pH 8.5 to 9.0<br><br>Topography: Undulating hills<br><br>Water: Rainfed, depth to watertable unknown.   | Fresh soils spread over contaminated site. Previously used as a tannery and later on as pasture for grazing.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.                       |
| 24  | Flaxley Holistic Trading, Mount Lofty Ranges<br>Lat/long: 350820, 1384950 | 1991 | 5 ha         | <i>E. fasciculosa</i> ,<br><i>E. obliqua</i> ,<br><i>Acacia melanoxylon</i><br><br>Revegetation                | 97.55 <sup>1</sup><br>73.17 <sup>2</sup> | 7   | 13.94       | -                           | N/A                       | Rainfall: Av 740 mm<br><br>Soil: Sandy loam, pH 6.0 to 5.5<br><br>Topography: Unknown<br><br>Water: Rainfed, depth to watertable unknown.  | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |

**A4 - Table 1a. Growth Data for Sites in South Australia, Hassall & Associates (1999a) continued.**

| Site Information (South Australia, Hassall & Associates, 1999a) |  |      |              |   | Biomass                                  | MAI | Planting           | Volume             | Site Factors   | Establishment & Stand Management   |
|---|--|------|--------------|---|--|-----|--------------------|--------------------|--|--|
| ID  | Name, Location, Coordinates  | Est. | Area Planted | Species List & Purpose  | t/ha                                     | Age | Density (trees/ha) | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. site preparation and prior landuse, tubestock/direct seed, weeds/disease/pests, fire, irrigation  |
| 25  | Flaxley Lodge (tubestock)<br>Mount Lofty Ranges<br><br>Lat/long: 350812, 1384950 | 1990 | 0.3 ha       | <i>E. globulus</i><br><br>Shelterbelt, farm forestry  | 46.37 <sup>1</sup><br>34.78 <sup>2</sup> | 8   | -                  | N/A                | Rainfall: Av 740 mm<br><br>Soil: Sandy loam, pH 5.5 to 6.0<br><br>Topography: Flat<br><br>Water: Rainfed, depth to watertable unknown.   | Site preparation unknown. Previously used for sheep grazing<br><br>Tubestock.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.        |
| 26  | Dingo Plains (direct seed)<br>Mallee<br><br>Lat/long: 352042, 1402337            | 1991 | 0.9 ha       | <i>Acacia argyrophylla</i><br><i>Acacia calamiifolia</i> ,<br><i>Acacia microcarpa</i><br><br>Purpose not specified | 3.49 <sup>1</sup><br>2.61 <sup>2</sup>   | 7   | -                  | N/A                | Rainfall: Av 368 mm<br><br>Soil: Clay loam with significant calcium carbonate deposits, pH 7.5 at surface to 8.0 at 30 cm<br><br>Topography: Rolling plain<br><br>Water: Rainfed, depth to watertable unknown. | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.               |
| 27  | Lameroo (tubestock)<br>Mallee<br><br>Lat/long: 352012, 1402539                   | 1990 | 1.1 ha       | <i>E. leucosylon</i><br><i>E. occidentalis</i><br><br>Farm forestry trial   | 14.35 <sup>1</sup><br>10.76 <sup>2</sup> | 8   | -                  | N/A                | Rainfall: Av 368 mm<br><br>Soil: Sand over clay loam, pH 7.0 at surface to 8.0 at 30 cm<br><br>Topography: Rolling plain<br><br>Water: Rainfed, depth to watertable unknown.                                   | Site preparation unknown. Previous use unknown.<br><br>Tubestock.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.                    |
| 28  | Wilkawatt (direct seed)<br>Mallee<br><br>Lat/long: 352541, 1402451               | 1992 | 0.6 ha       | <i>Dodonaea viscosa</i> ,<br><i>E. porosa</i> ,<br><i>Melaleuca acuminata</i><br><br>Roadside shelterbelt           | 24.99 <sup>1</sup><br>18.74 <sup>2</sup> | 6   | -                  | N/A                | Rainfall: Av 368 mm<br><br>Soil: Red brown loam over brown clay, pH 7.0 at surface to 8.5 at 40 cm<br><br>Topography: Flat plain<br><br>Water: Rainfed, depth to watertable unknown.                           | Site preparation unknown. Previously used for mixed cropping.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only. |



| Site Information (South Australia, Hassall & Associates, 1999a) |  |      |              |  |  |     |             |                             |                           | Establishment & Stand Management  |  |
|---|--|------|--------------|--|--|-----|-------------|-----------------------------|---------------------------|---|--|
| ID  | Name, Location, Coordinates                                | Est. | Area Planted | Species List & Purpose   | Biomass t/ha                             | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. site preparation and prior landuse, tubestock/direct seed, weeds/disease/pests, fire, irrigation  |
| 29  | Kalisch, Riverland<br>Lat/long: 341245, 1400423            | 1995 | 4 ha         | <i>Acacia oswaldii</i> ,<br><i>Acacia ligulata</i> ,<br><i>Acacia cyanophylla</i><br><br>Purpose not specified                 | 0.00 <sup>1</sup><br>0.00 <sup>2</sup>   | 3   | 0.00        | -                           | N/A                       | Rainfall: Av 266 mm<br><br>Soil: Sand, pH 6.5 at surface to 7.5 at 50 cm<br><br>Topography: Flat plain<br><br>Water: Rainfed, depth to watertable unknown.  | Site preparation unknown. Previously used for grazing.<br><br>Direct seeding.<br><br>No thinnings.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.   |
| 30  | Karelia, Riverland<br>Lat/long: 1401751, 341156            | 1994 | 0.7 ha       | <i>Acacia ligulata</i> ,<br><i>Acacia oswaldii</i> ,<br><i>Acacia sp.</i> (juvenile)<br><br>Dryland revegetation demonstration | 0.07 <sup>1</sup><br>0.05 <sup>2</sup>   | 4   | 0.02        | -                           | N/A                       | Rainfall: Av 241 mm<br><br>Soil: Sandy loam over limestone with significant calcium carbonate throughout, pH 9.0<br><br>Topography: Rolling plain<br><br>Water: Rainfed, depth to watertable unknown.                                 | Site preparation unknown. Previous use unknown.<br><br>Direct seeding.<br><br>No thinnings.<br><br>No problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.  |
| 31  | Orlando-Wyndham, Riverland<br>Lat/long: 340849, 1395434    | 1994 | 0.9 ha       | <i>Casuarina obesa</i> ,<br><i>E. camaldulensis</i> ,<br><i>E. largiflorens</i><br><br>Groundwater recharge control            | 31.97 <sup>1</sup><br>23.98 <sup>2</sup> | 4   | 7.99        | -                           | N/A                       | Rainfall: Av 235 mm<br><br>Soil: Sandy loam, pH 7.0-7.5<br><br>Topography: River floodplain<br><br>Water: Rainfed plus access irrigation from neighboring vineyards, depth to watertable unknown but expected to be within root zone. | Site preparation unknown. Previous use unknown.<br><br>Tubestock.<br><br>Thinnings history unknown.<br><br>Some weeds present but no threat. No other problems associated with disease, pests, drought or fire.<br><br>Rainfed plus irrigated. |
| 32  | Little Desert Lodge, Victoria<br>Lat/long: 362725, 1413937 |      |              | <i>Allocasuarina verticillata</i> ,<br><i>E. fasciculosa</i> ,<br><i>E. leucoxydon</i><br><br>Soil stabilisation, habitat      | 3.00 <sup>1</sup><br>2.25 <sup>2</sup>   | 6   | 0.5         | -                           | N/A                       | Rainfall: Av 418 mm<br><br>Soil: Sand, pH 7.0 at surface to 7.5 at 50 cm<br><br>Topography: Low undulating plain<br><br>Water: Rainfed, depth to watertable unknown.  | Site preparation unknown. Previously used for cattle grazing.<br><br>Direct seeding.<br><br>Thinnings history unknown.<br><br>No problems associated with weeds, disease, pests, drought or fire.<br><br>Rainfed only.                         |

<sup>1</sup> measured using bole and branches  
<sup>2</sup> measured using bole only

A4 - Table 1b. Growth Data for Sites in Victoria, Hassall &amp; Associates (1998)

| Site Information (Victoria, Hassall & Associates, 1998) |  |      |                  |  |                                     |     |                |                                   |                              |  |   |
|---|--|------|------------------|--|-------------------------------------|-----|----------------|-----------------------------------|------------------------------|--|---|
| ID  | Name, Location, Coordinates  | Est. | Area Planted     | Species List & Purpose   | Biomass<br>t/ha                     | Age | MAI<br>t/ha/yr | Planting<br>Density<br>(trees/ha) | Volume<br>m <sup>3</sup> /ha | Site Factors<br>e.g. rainfall, soil type/quality,<br>topography, access to water, salinity   | Husbandry<br>e.g. tubestock/direct seed, weed & pest control,<br>fire, irrigation   |
| 1   | Wail,<br>near Horsham, Vic<br><br>Vic Roads map ref: 26 C10            | 1973 | Approx<br>130 ha | Sugar Gum<br><i>(E. cladocalyx)</i><br><br>Wood products                                       | 81 <sup>1</sup><br>61 <sup>2</sup>  | 23  | 3.5            | N/A                               | N/A                          | Rainfall: Av 420 mm<br><br>Soil: Clay loam overlying impeding layer<br>of clay, pH 6.5-7.5<br><br>Topography: Flat<br><br>Water: No runon/runoff, depth to<br>groundwater unknown. | Seedlings planted in block formation, thinnings<br>history unknown.<br><br>Land previously was native bush, cleared,<br>cultivated and ripped before planting.<br><br>Weeds/disease/pests have not been a problem.<br><br>Unburnt.<br><br>Rainfed only, some periods of drought.                              |
| 2   | Wail,<br>near Horsham, Vic.<br><br>Vic Roads map ref: 26 C10           | 1975 | Approx<br>130 ha | See above  | 121 <sup>1</sup><br>90 <sup>2</sup> | 21  | 5.8            | N/A                               | N/A                          | See above  | See above   |
| 3   | Barrett Reserve,<br>near Horsham, Vic.<br><br>Vic Roads map ref: 26 F8 | 1978 | Approx<br>32 ha  | Shining Gum<br><i>(E. sieberi)</i><br>Sugar Gum<br><i>(E. cladocalyx)</i><br><br>Wood Products | 77 <sup>1</sup><br>58 <sup>2</sup>  | 17  | 4.5            | N/A                               | N/A                          | Rainfall: Av 454 mm<br><br>Soil: Black with high clay content,<br>pH 7.0-7.5<br><br>Topography: Flat<br><br>Water: No runon/runoff, depth to<br>groundwater unknown.               | Seedlings planted in block formation, selective<br>thinnings in parts of the plantation.<br><br>Land preparation unknown, previously used<br>for grazing.<br><br>Weeds/disease/pests have not been a problem.<br><br>Unburnt.<br><br>Rainfed only, some periods of drought which<br>may have affected growth. |
| 4   | Barrett Reserve,<br>near Horsham, Vic.<br><br>Vic Roads map ref: 26 F8 | 1986 | Approx<br>32 ha  | See above  | 16 <sup>1</sup><br>12 <sup>2</sup>  | 9   | 1.8            | N/A                               | N/A                          | See above  | See above   |
| 5   | Barrett Reserve,<br>near Horsham, Vic.<br><br>Vic Roads map ref: 26 F8 | 1973 | Approx<br>32 ha  | See above  | 72 <sup>1</sup><br>54 <sup>2</sup>  | 22  | 3.3            | N/A                               | N/A                          | See above  | See above   |
| 6   | Barrett Reserve,<br>near Horsham, Vic.<br><br>Vic Roads map ref: 26 F8 | 1971 | Approx<br>32 ha  | See above  | 78 <sup>1</sup><br>59 <sup>2</sup>  | 24  | 3.3            | N/A                               | N/A                          | See above  | See above   |

| Site Information (Victoria, Hassall & Associates, 1998) |   |      |               |  |                                      |     |             |                             |                           | Husbandry   |   |
|---|---|------|---------------|--|--------------------------------------|-----|-------------|-----------------------------|---------------------------|---|---|
| ID  | Name, Location, Coordinates                                     | Est. | Area Planted  | Species List & Purpose   | Biomass t/ha                         | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 7   | Barrett Reserve, near Horsham, Vic.<br>Vic Roads map ref: 26 F8 | 1975 | Approx 32 ha  | Sugar Gum ( <i>E. cladocalyx</i> )<br>Shining Gum ( <i>E. sieberi</i> )<br>Wood Products | 64 <sup>1</sup><br>48 <sup>2</sup>   | 20  | 3.2         | N/A                         | N/A                       | See above   | See above   |
| 8   | You Yangs, near Geelong, Vic.<br>Vic Roads map ref: 77 H9       | 1966 | Approx 450 ha | Sugar Gum ( <i>E. cladocalyx</i> )<br>Wood Products                                      | 145 <sup>1</sup><br>108 <sup>2</sup> | 30  | 4.8         | N/A                         | N/A                       | Rainfall: Av 491 mm<br>Soil: Granitic<br>Topography: Flat<br>Water: No runoff/runoff, depth to groundwater unknown.   | Trees probably planted as seedlings, block formations. No thinnings.<br>Land preparation unknown.<br>No significant problems with weeds, disease or pests.<br>Fire has occurred but does not appear to have affected the trees significantly.<br>Rainfed only, some periods of drought which may have affected the trees.                       |
| 9   | You Yangs, near Geelong, Vic.<br>Vic Roads map ref: 77 H9       | 1965 | Approx 450 ha | See above  | 175 <sup>1</sup><br>131 <sup>2</sup> | 31  | 5.6         | N/A                         | N/A                       | See above   | See above   |
| 10  | You Yangs, near Geelong, Vic.<br>Vic Roads map ref: 77 H9       | 1958 | Approx 450 ha | Brown Mallett ( <i>E. astringens</i> )<br>Wood Products                                  | 175 <sup>1</sup><br>131 <sup>2</sup> | 39  | 4.5         | N/A                         | N/A                       | See above   | See above   |
| 11  | You Yangs, near Geelong, Vic.<br>Vic Roads map ref: 77 H9       |      | Approx 450 ha | Brown Mallett ( <i>E. astringens</i> )<br>Wood Products                                  | 115 <sup>1</sup><br>86 <sup>2</sup>  | 38  | 3.0         | N/A                         | N/A                       | See above   | See above   |
| 12  | 'Titanga', near Lismore, Vic.                                   | 1994 | Approx 8 ha   | Southern Blue Gum ( <i>E. globulus</i> ssp. <i>globulus</i> )<br>Wood Products           | 7.1 <sup>1</sup><br>5.3 <sup>2</sup> | 2   | 3.6         | N/A                         | N/A                       | Rainfall: Av 630 mm<br>Soil: Granitic sand and red lateritic gravel, pH unknown<br>Topography: Gentle slope from elevation 30m to 200m, NW aspect<br>Water: Surface runoff downslope, depth to groundwater unknown. | Seedlings used in rows. No thinnings.<br>Land ripped, formed and sprayed along rows before planting, previously used for grazing.<br>Weeds a problem during establishment, more recently attacks from insects. No treatment undertaken. No other problems.<br>Rainfed only, some periods of drought that do not appear to have affected growth. |

**A4 - Table 1b. Growth Data for Sites in Victoria, Hassall & Associates (1998) continued.**

| Site Information (Victoria, Hassall & Associates, 1998) |  |      |               |   |                                      |             |                             |                           |  |   |
|---|--|------|---------------|---|--------------------------------------|-------------|-----------------------------|---------------------------|--|---|
| ID  | Name, Location, Coordinates              | Est. | Area Planted  | Species List & Purpose                    | Biomass t/ha                         | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity                                   | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 13  | 'Lingalonga', Mtiamo, NW of Bendigo, Vic | 1984 | Approx 40 ha  | River Red Gum ( <i>E. camaldulensis</i> ) | 66 <sup>1</sup><br>50 <sup>2</sup>   | 12          | 5.5                         | N/A                       | Rainfall: Av 399 mm<br><br>Soil: Grey cracking clay, pH neutral  | Grown from seed collected on property in to 2 rows along roads and property boundaries. No thinnings, but some trees blown over or dead.  |
|   | Vic Roads map ref: 30 B4                 |      |               | Amenity                                   |                                      |             |                             |                           | Topography: Flat<br><br>Water: Some surface runon during wet months; depth to groundwater 1m.                          | Limited preparation prior to planting, deep ripped but no weed control, previously used for irrigated dairy pastures.<br><br>Initial weed problems, recently rabbits/hares in plague numbers. No treatments undertaken.<br><br>Unburnt.<br><br>Rainfed mainly, but close to irrigated pastures. |
| 14  | Cohuna, N of Bendigo, Vic                | 1979 | Approx 2.5 ha | River Red Gum ( <i>E. camaldulensis</i> ) | 286 <sup>1</sup><br>214 <sup>2</sup> | 17          | 16.8                        | N/A                       | Rainfall: Av 450 mm<br><br>Soil: Clay loams, pH 6.0-6.5, saline in areas   | Seedlings planted in block formations. No thinnings since.<br><br>Land deep ripped and formed prior to planting, previously used for irrigated annual and perennial pastures.   |
|   | Vic Roads map ref: 21 E6                 |      |               | Shelterbelts, pest control, amenity       |                                      |             |                             |                           | Topography: Flat<br><br>Water: Some surface runon from irrigation; depth to groundwater > 1m.                          | Some disease and pests present during establishment, but no treatments used.<br><br>Unburnt.<br><br>Rainfed, plus some areas flooded occasionally from adjacent irrigated pastures.   |
| 15  | Stanhope, W of Shepparton, Vic           | 1955 | N/A           | River Red Gum ( <i>E. camaldulensis</i> ) | 902 <sup>1</sup><br>677 <sup>2</sup> | 37          | 24.4                        | N/A                       | Rainfall: Av 500 mm<br><br>Soil: Lemnos loam, pH 4.5, no salinity, high fertility from a long history of dairy farming | Seedlings planted mainly as single rows along fence lines. No thinnings since.<br><br>Land not prepared before planting.  |
|   | Vic Roads map ref: 31 H9                 |      |               | Shelterbelts                              |                                      |             |                             |                           | Topography: Flat<br><br>Water: Regularly irrigated plus rainfed, depth to groundwater 1-2m.                            | Hares a problem during establishment. A once-off outbreak of Sawflies affected young trees and growth, treated with 'Curacine'.<br><br>Unburnt.<br><br>Regularly irrigated (approx once per fortnight) via adjacent irrigated dairy pastures.   |

| Site Information (Victoria, Hassall & Associates, 1998) |                                      |      |              |   |  |     |                |                                   |                              |   |   |
|---|--------------------------------------|------|--------------|---|--|-----|----------------|-----------------------------------|------------------------------|---|---|
| ID  | Name, Location, Coordinates          | Est. | Area Planted | Species List & Purpose  | Biomass<br>t/ha                          | Age | MAI<br>t/ha/yr | Planting<br>Density<br>(trees/ha) | Volume<br>m <sup>3</sup> /ha | Site Factors<br>e.g. rainfall, soil type/quality,<br>topography, access to water, salinity  | Husbandry<br>e.g. tubestock/direct seed, weed & pest control,<br>fire, irrigation   |
| 16  | Stanhope,<br>W of Shepparton, Vic    | 1958 | N/A          | River Red Gum<br>( <i>E. camaldulensis</i> )  | 2,889 <sup>1</sup><br>2,167 <sup>2</sup> | 42  | 16.8           | N/A                               | N/A                          | See above   | See above   |
|   | Vic Roads map ref: 31 H9             |      |              | Shelterbelts  |  |     |                |                                   |                              |   |   |
| 17  | Stanhope,<br>W of Shepparton, Vic    | 1960 | N/A          | River Red Gum<br>( <i>E. camaldulensis</i> )  | 4131<br>3102                             | 39  | 10.6           | N/A                               | N/A                          | See above   | See above   |
|   | Vic Roads map ref: 31 H9             |      |              | Shelterbelts  |  |     |                |                                   |                              |   |   |
| 18  | Stanhope,<br>W of Shepparton, Vic    | 1965 | N/A          | River Red Gum<br>( <i>E. camaldulensis</i> )  | 1,620 <sup>1</sup><br>1,215 <sup>2</sup> | 32  | 50.6           | N/A                               | N/A                          | See above   | See above   |
|   | Vic Roads map ref: 31 H9             |      |              | Shelterbelts  |  |     |                |                                   |                              |   |   |
| 19  | Katamatite,<br>NE of Shepparton, Vic | 1965 | N/A          | River Red Gum<br>( <i>E. camaldulensis</i> )  | 1,271 <sup>1</sup><br>953 <sup>2</sup>   | 21  | 60.5           | N/A                               | N/A                          | Rainfall: Av 500 mm<br><br>Soil: Moira loam, pH 5.5, no salinity,<br>high fertility from a history of<br>dairy farming<br><br>Topography: Flat<br><br>Water: Regularly irrigated plus rainfed,<br>depth to groundwater varies between 3m<br>and 9m depending on season. | Seedlings planted as a single row along property<br>boundary. No thinnings since.<br><br>Land ripped and formed before planting.<br><br>Hares a problem during establishment. No other<br>problems since.<br><br>Regularly irrigated (approx once per fortnight) via<br>adjacent water channel for dairy pasture. |
|   | Vic Roads map ref: 33 C3             |      |              | Shelterbelts,<br>amenity  |  |     |                |                                   |                              |   |   |
| 20  | Leichardt,<br>near Bendigo, Vic.     | 1987 | 10 ha        | River Red Gum<br>( <i>E. camaldulensis</i> )  | 5.1 <sup>1</sup><br>3.8 <sup>2</sup>     | 9   | 0.6            | N/A                               | N/A                          | Rainfall: Av 373 mm<br><br>Soil: Red loam, slightly acidic, rocky<br>in parts<br><br>Topography: Gentle undulating<br><br>Water: Rainfed only, depth to<br>groundwater varies between 1m and 2m.  | Seedlings planted in widely spaced shelterbelt<br>formations. No thinnings.<br><br>Land previously used for grazing, ripped prior<br>to planting.<br><br>Rabbits initially a problem but less so now due<br>to ongoing control. No other problems.<br><br>Rainfed only.   |
|   | Vic Roads map ref: 44 B5             |      |              | Drooping Sheoak<br>( <i>Allocasuarina<br/>verticillata</i> )<br><br>Various Mallee<br>species<br><br>Erosion control,<br>waterable/ dryland<br>salinity control |  |     |                |                                   |                              |   |   |



**A4 - Table 1b. Growth Data for Sites in Victoria, Hassall & Associates (1998) continued.**

| Site Information (Victoria, Hassall & Associates, 1998) |   | Biomass |              | MAI  | Planting | Volume             | Site Factors  | Hubandry   |
|---|---|---------|--------------|--|----------|--------------------|---|--|
| ID  | Name, Location, Coordinates   | Est.    | Area Planted | Species List & Purpose   | Age      | Density (trees/ha) | e.g. rainfall, soil type/quality, topography, access to water, salinity   | e.g. tubestock/direct seed, weed & pest control, fire, irrigation  |
| 21  | Bear's Lagoon, NE of Bendigo, Vic.<br><br>Vic Roads map ref: 29 H7                  | 1972    | N/A          | River Red Gum<br>( <i>E. camaldulensis</i> )<br><br>Black Box<br>( <i>E. largiflorens</i> )<br><br>Manna Gum<br>( <i>E. viminalis</i> )<br>Shelterbelts              | 24       | N/A                | MAI: 7.9<br><br>Rainfall: Av 450 mm<br><br>Soil: Heavy clay loam, pH 5.5<br><br>Topography: Flat<br><br>Water: Rainfed with occasional flooding, depth to groundwater 7m.                       | Seedlings planted in a single row along fence lines. No thinnings since.<br><br>Land preparation unknown. Previously used for mixed farming.<br><br>No known problems from insects, disease, pests, drought or fire.<br><br>Rainfed, with occasional flooding.   |
| 22  | Burke's Flat, W of Bendigo, Vic.<br><br>Vic Roads map ref: 43 A4                    | 1983    | 5-10 ha      | Red Ironbark<br>( <i>Eucalyptus sideroxylon</i> )<br><br>Yellow Gum<br>( <i>Eucalyptus leucoxylon</i> )<br><br>Erosion control, watertable/ dryland salinity control | 13       | 600-300            | MAI: 5.6<br><br>Rainfall: Av 458 mm<br><br>Soil: Heavily leached, gradational, pH 4.0-4.5<br><br>Topography: Gently undulating<br><br>Water: Rainfed only, depth to groundwater <10m.           | Trees planted as tubestock. No thinnings but some natural deaths and regeneration.<br><br>Land ripped prior to planting. Previously used for grazing.<br><br>Some insect damage from 1989-2001 during the winter months. No other problems from insects, disease, pests, drought or fire.<br><br>Rainfed only. |
| 23  | Colbinabbin Range (Toolleen), W of Shepparton, Vic.<br><br>Vic Roads map ref: 45 D5 | 1982    | 5-10 ha      | Grey Box<br>( <i>Eucalyptus microcarpa</i> )<br><br>Drooping Sheoak<br>( <i>Allocasuarina verticillata</i> )<br><br>Watertable/ dryland salinity control, amenity    | 14       | N/A                | MAI: 1.8<br><br>Rainfall: Av 491 mm<br><br>Soil: Gradational red brown, mild acidity, not saline<br><br>Topography: 10% slope on hilltop<br><br>Water: Rainfed only, depth to groundwater <10m. | Seedlings planted in rows on hilltop. No thinnings, but significant natural attrition.<br><br>Land ripped prior to planting. Previously used for grazing.<br><br>No known problems from insects, disease, pests or fire. Severe drought in 1982/83 killed approx 55% of plot.<br><br>Rainfed only.             |

| Site Information (Victoria, Hassall & Associates, 1998) |   |      |              |  |   |     |                |                                   |                              |  |   |
|---|---|------|--------------|--|---|-----|----------------|-----------------------------------|------------------------------|--|---|
| ID  | Name, Location, Coordinates                 | Est. | Area Planted | Species List & Purpose                               | Biomass<br>t/ha   | Age | MAI<br>t/ha/yr | Planting<br>Density<br>(trees/ha) | Volume<br>m <sup>3</sup> /ha | Site Factors<br>e.g. rainfall, soil type/quality,<br>topography, access to water, salinity   | Husbandry<br>e.g. tubestock/direct seed, weed & pest control,<br>fire, irrigation   |
| 24  | Pyalong,<br>SW of Shepparton, Vic.          | 1991 | 0.25 ha      | Red Stringybark<br>( <i>E. macrorhyncha</i> )        | 6 <sup>1</sup>  | 5   | 1.2            | N/A                               | N/A                          | Rainfall: Av 631 mm  | Trees planted via direct seeding. Some removal of unwanted species since planting, but otherwise no thinnings.  |
|   | Vic Roads map ref: 60 F3                    |      |              | Black Wattle<br>( <i>Acacia mearnsii</i> )           | 4.5 <sup>2</sup>  |     |                |                                   |                              | Soil: Ironstone of sedimentary origin, pH approx 4.8<br><br>Topography: 20% hillside<br><br>Water: Rainfed only, depth to groundwater unknown. | Some ripping to prepare the land prior to seeding, but not over the entire plot. Previously used for grazing.<br><br>No problems from insects, disease, pests, drought or fire.<br><br>Rainfed only.  |
|   |   |      |              |  | Watertable/ dryland salinity control, experimental  |     |                |                                   |                              |  |   |
| 25  | Puckapunyal Range,<br>SW of Shepparton, Vic | 1980 | 40 ha        | Red Ironbark<br>( <i>Eucalyptus sideroxydon</i> )    | 659 <sup>1</sup>  | 15  | 43.9           | N/A                               | N/A                          | Rainfall: Av 598 mm  | Trees planted as seedlings on contour lines around hilltop. No thinnings.   |
|   | Vic Roads map ref: 61 B1                    |      |              | Yellow Gum<br>( <i>E.leucoxydon</i> )                | 494 <sup>2</sup>  |     |                |                                   |                              | Soil: Heavy clay (buckshot gravel), pH 5.5-6.0<br><br>Topography: Gentle slope or flat   | Before planting, the site was contour ripped and sprayed. Previously used for grazing and army training.  |
|   |   |      |              |  | Grey Box<br>( <i>E. microcarpa</i> )<br><br>Watertable/dryland salinity control, ecology & biodiversity |     |                |                                   |                              | Water: Rainfed plus groundwater discharge in parts.  | Attacks by insects have been assumed although not recorded as the cause of foliage decline over the years. No treatment undertaken. No other problems with disease, fire or drought.<br><br>Rainfed, although proximity to groundwater discharge is having an impact on growth rates. |
| 26  | Toolleen,<br>E of Bendigo, Vic.             | 1987 | N/A          | River Red Gum<br>( <i>Eucalyptus camaldulensis</i> ) | 104 <sup>1</sup>  | 9   | 11.6           | N/A                               | N/A                          | Rainfall: Av 450 mm  | Seedlings planted as a double row windbreak along fence lines. No thinnings.  |
|   | Vic Roads map ref: 45 D5                    |      |              | Yate<br>( <i>E. accidentalis</i> )                   | 78 <sup>2</sup>   |     |                |                                   |                              | Soil: Red volcanic loam, with heavy clay lower horizon, pH approx 5.9<br><br>Topography: Unknown   | Site was deep ripped (up to 40 cm) and sprayed. Previously used for mixed farming.  |
|   |   |      |              |  | Watertable/ dryland salinity control, shelterbelts  |     |                |                                   |                              | Water: Rainfed only, depth to groundwater varies around 1m depending on season.  | Outbreak of spittfires treated with Malathion. Some trees have died and been replanted because of drought in 1982/83. No other problems from insects, disease, pests or fire.<br><br>Rainfed, with occasional flooding. Flooding frequency unknown.                                   |

3 measured using bole and branches  
4 measured using bole only

**A4 - Table 1c. Growth Data for Sites in Western Australia, Hassall & Associates (1999b)**

| Site Information (Western Australia, Hassall & Associates, 1999b) |                             |      |              | Biomass  | MAI                                  | Planting | Volume             | Site Factors       | Husbandry   |  |   |
|---|-----------------------------|------|--------------|--|--------------------------------------|----------|--------------------|--------------------|---|--|---|
| ID  | Name, Location, Coordinates | Est. | Area Planted | Species List & Purpose   | t/ha                                 | Age      | Density (trees/ha) | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity | e.g. tubestock/direct seed, weed & pest control, fire, irrigation  |   |
| 1   | Wongon Hills                | 1965 | 3.5 ha       | Dwarf Sugar Gum ( <i>E.cladocalyx</i> var. <i>nana</i> )<br><br>Amenity  | 76 <sup>1</sup><br>57 <sup>2</sup>   | 32.5     | 2.4                | 150                | N/A   | Rainfall: Av 392 mm<br><br>Soil: Yellow/brown sand over deep loamy sand, pH approx 4.8-5.2 (CaCl2)<br><br>Topography: Gently undulating<br><br>Water: Rainfed only, depth to groundwater >1m<br><br>Aspect: NE.                          | Site preparation unknown. Previously cropping/grazing.<br><br>Tubestock planted in rows<br><br>No thinnings and no replanting.<br><br>Some species have suffered from competition from more vigorous species. No other problems associated with weeds/disease/pests, drought or fire.<br><br>Rainfed only.    |
| 2   | Wongon Hills                | 1965 | 3.5 ha       | Sugar Gum ( <i>E. cladocalyx</i> )<br><br>Amenity  | 160 <sup>1</sup><br>120 <sup>2</sup> | 32.5     | 4.9                | 110                | N/A   | See above  | See above   |
| 3   | Wongon Hills                | 1951 | 8 ha         | Salt River Gum ( <i>E. sargentii</i> )<br><br>Erosion control, watertable/dryland salinity control   | 73 <sup>1</sup><br>55 <sup>2</sup>   | 46.5     | 1.6                | 200                | N/A   | Rainfall: Av 392 mm<br><br>Soil: Black clayey sand over brown sandy loam, pH approx 5.0 (CaCl2)<br><br>Topography: Flat<br><br>Water: Rainfed plus possible access to groundwater, depth to groundwater approx 20 cm<br><br>Aspect: N/A. | Site preparation unknown. Prior land use unknown.<br><br>Tubestock planted in rows along contours.<br><br>No thinnings, but site replanted to replace deaths and increase density.<br><br>Incidence of weeds/disease/pests, drought or fire unknown.<br><br>Rainfed plus access to shallow groundwater.       |
| 4   | Ejandling West              | 1984 | 20 ha        | Dwarf Sugar Gum ( <i>E. cladocalyx</i> var. <i>nana</i> )<br><br>Coastal Mort ( <i>E. platypus</i> var. <i>heterophylla</i> )<br><br>Erosion control, watertable/dryland salinity control, windbreak | 68 <sup>1</sup><br>51 <sup>2</sup>   | 13.5     |                    | 285                | N/A   | Rainfall: Av 311 mm<br><br>Soil: Sand over yellow brown clayey sand, non-saline, pH approx 4.5<br><br>Topography: Gentle slope<br><br>Water: Rainfed only, depth to groundwater approx 10-12m<br><br>Aspect: W.                          | Site ripped prior to planting. Previously used for cropping/grazing.<br><br>Tubestock planted in rows across contours.<br><br>No thinnings. Some initial loss of trees due to drought and wind damage, but none since.<br><br>Incidence of weeds/disease/pests, drought or fire unknown.<br><br>Rainfed only. |

A4 - Table 1c. Growth Data for Sites in Western Australia, Hassall &amp; Associates (1999b) continued.

| Site Information (Western Australia, Hassall & Associates, 1999b)                         |                             |      |              |  |                                      |             |                |                             |                           |
|---|-----------------------------|------|--------------|--|--------------------------------------|-------------|----------------|-----------------------------|---------------------------|
| ID  | Name, Location, Coordinates | Est. | Area Planted | Species List & Purpose   | Biomass t/ha                         | Age t/ha/yr | MAI (trees/ha) | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha |
| Site Factors  |                             |      |              |  |                                      |             |                |                             |                           |
| e.g. rainfall, soil type/quality, topography, access to water, salinity                   |                             |      |              |  |                                      |             |                |                             |                           |
| Husbandry   |                             |      |              |  |                                      |             |                |                             |                           |
| e.g. tubestock/direct seed, weed & pest control, fire, irrigation                         |                             |      |              |  |                                      |             |                |                             |                           |
| 5   | Ejanding                    | 1985 | 2 ha         | Salt River Gum ( <i>E. sargentii</i> )<br><br>Erosion control, waterable/dry/and salinity control    | 10 <sup>1</sup><br>7 <sup>2</sup>    | 12.5        |                | 500                         | N/A                       |
| Rainfall: Av 311 mm   |                             |      |              |  |                                      |             |                |                             |                           |
| Soil: yellow brown sand over light grey sandy clay loam, moderate salinity, pH approx 5.0 |                             |      |              |  |                                      |             |                |                             |                           |
| Topography: unknown   |                             |      |              |  |                                      |             |                |                             |                           |
| Water: Rainfed plus possible access to groundwater, depth to groundwater approx 20 cm     |                             |      |              |  |                                      |             |                |                             |                           |
| Aspect: N/A.  |                             |      |              |  |                                      |             |                |                             |                           |
| 6   | Trayning                    | 1987 | 1 ha         | River Red Gum ( <i>E. camaldulensis</i> )<br><br>Erosion control, waterable/dry/and salinity control | 45 <sup>1</sup><br>34 <sup>2</sup>   | 10.5        |                | 450-500                     | N/A                       |
| Rainfall: Av 323 mm   |                             |      |              |  |                                      |             |                |                             |                           |
| Soil: Yellow brown sand over bright yellow brown sandy loam, pH 6.1                       |                             |      |              |  |                                      |             |                |                             |                           |
| Topography: Midslope, very gentle slope   |                             |      |              |  |                                      |             |                |                             |                           |
| Water: Rainfed only, depth to groundwater approx 30 cm                                    |                             |      |              |  |                                      |             |                |                             |                           |
| Aspect: NNE.  |                             |      |              |  |                                      |             |                |                             |                           |
| Site ripped and sprayed prior to planting. Previously used for cropping.                  |                             |      |              |  |                                      |             |                |                             |                           |
| Seedlings planted in contoured rows using a tree planter.                                 |                             |      |              |  |                                      |             |                |                             |                           |
| No thinnings. Some recent deaths due to drought. No replanting.                           |                             |      |              |  |                                      |             |                |                             |                           |
| No damage from weeds/disease/pests or fire.   |                             |      |              |  |                                      |             |                |                             |                           |
| Rainfed only.   |                             |      |              |  |                                      |             |                |                             |                           |
| 7   | Trayning                    | 1987 | 1 ha         | See above  | 7.3 <sup>1</sup><br>5.4 <sup>2</sup> | 10.5        |                | 450-500                     | N/A                       |
| See above except:   |                             |      |              |  |                                      |             |                |                             |                           |
| Water: Rainfed only, depth to groundwater >1.5m.  |                             |      |              |  |                                      |             |                |                             |                           |
| 8   | Trayning                    | 1987 | 1 ha         | Salt River Gum ( <i>E. sargentii</i> )<br><br>Erosion control, waterable/dry/and salinity control    | 28 <sup>1</sup><br>21 <sup>2</sup>   | 10.5        |                | 450-500                     | N/A                       |
| See above except:   |                             |      |              |  |                                      |             |                |                             |                           |
| pH 5.1  |                             |      |              |  |                                      |             |                |                             |                           |
| Aspect: SSW.  |                             |      |              |  |                                      |             |                |                             |                           |
| See above except:   |                             |      |              |  |                                      |             |                |                             |                           |
| Some deaths due to borers in the drier areas.   |                             |      |              |  |                                      |             |                |                             |                           |
| 9   | Trayning                    | 1987 | 1 ha         | See above  | 3.9 <sup>1</sup><br>3.0 <sup>2</sup> | 10.5        |                | 450-500                     | N/A                       |
| See above   |                             |      |              |  |                                      |             |                |                             |                           |



| Site Information (Western Australia, Hassall & Associates, 1999b) |                             |      |              |   |                                    |      |             |                             |                           | Site Factors   | Husbandry   |
|---|-----------------------------|------|--------------|---|------------------------------------|------|-------------|-----------------------------|---------------------------|--|---|
| ID  | Name, Location, Coordinates | Est. | Area Planted | Species List & Purpose  | Biomass t/ha                       | Age  | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 10  | Bencubbin                   | 1982 | N/A          | River Red Gum<br>( <i>E. camaldulensis</i> )<br><br>Windbreak   | 16 <sup>1</sup><br>12 <sup>2</sup> | 15.5 |             | 500                         | N/A                       | Rainfall: Av 322 mm<br><br>Soil: Deep yellow woodjill sands, pH 4.5 (H <sub>2</sub> O)<br><br>Topography: High up on a gently sloping hillside<br><br>Water: Rainfed only, depth to groundwater unknown but deep<br><br>Aspect: W.                               | Site ripped and sprayed prior to planting. Previously used for cropping/grazing.<br><br>Seedlings planted in rows. Thinnings history unknown.<br><br>Initial deaths due to rabbits. Treated by baiting and protective netting. Seedlings replanted. Some recent deaths due to drought and associated borer activity. No treatment.<br><br>Rainfed only. |
| 11  | Beacon                      | 1986 | 0.25 ha      | River Red Gum<br>( <i>E. camaldulensis</i> )<br><br>Shade and shelter   | 11 <sup>1</sup><br>8 <sup>2</sup>  | 11.5 |             | 200                         | N/A                       | Rainfall: Av 314 mm<br><br>Soil: Reddish brown sandy loam, low salinity, pH 7.6-7.8 (H <sub>2</sub> O)<br><br>Topography: Flat<br><br>Water: Rainfed plus access to groundwater. Depth to groundwater approx 1.5m, salinity 6000 mS/m<br><br>Aspect: N/A.        | Site preparation history unknown. Previously used for cropping/grazing.<br><br>Seedlings planted in rows by hand. Thinnings history unknown.<br><br>The only site difficulty appears to be reasonably shallow and saline groundwater.<br><br>Rainfed plus access to groundwater.  |
| 12  | Murkinbudin                 | 1986 | 1.5 ha       | York Gum (I)<br><br>Strickland's Gum<br>( <i>E. stricklandii</i> )<br><br>Sheath's Gum<br>( <i>E. sheathiana</i> )<br><br>Erosion control, waterable/dryland salinity control | 18 <sup>1</sup><br>13 <sup>2</sup> | 11.5 |             | 250                         | N/A                       | Rainfall: Av 285 mm<br><br>Soil: Brown sandy loam, over bedrock, salt scalds in parts, moderately acid.<br><br>Topography: Very gentle midslope<br><br>Water: Rainfed plus potential access to groundwater seep. Depth to groundwater unknown.<br><br>Aspect: S. | Site ripped but not sprayed. Previously used for cropping/grazing.<br><br>Seedlings planted in 5 contoured rows adjacent salt scald using a tree planter. Thinnings history unknown.<br><br>No reported problems with pests/weeds/disease, fire or drought.<br><br>Rainfed, plus potential access to groundwater  |

A4 - Table 1c. Growth Data for Sites in Western Australia, Hassall &amp; Associates (1999b) continued.

| Site Information (Western Australia, Hassall & Associates, 1999b) |                             |      |                               |  | Biomass                            | MAI  | Planting | Volume                         | Site Factors       | Husbandry  |  |
|---|-----------------------------|------|-------------------------------|--|------------------------------------|------|----------|--------------------------------|--------------------|--|--|
| ID  | Name, Location, Coordinates | Est. | Area Planted                  | Species List & Purpose   | t/ha                               | Age  | t/ha/yr  | Density (trees/ha)             | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. tubestock/direct seed, weed & pest control, fire, irrigation  |
| 13  | Belka East                  | 1988 | 2 ha                          | River Red Gum ( <i>E. camaldulensis</i> )<br><br>Red Ironbark ( <i>E. sideroxylon</i> )<br><br>Erosion control, watertable/dryland salinity control, amenity | 56 <sup>1</sup><br>42 <sup>2</sup> | 9.5  |          | N/A                            | N/A                | Rainfall: Av 332 mm<br><br>Soil: Sand over bright yellowish brown sandy loam, pH 4.2<br><br>Topography: Midslope<br><br>Water: Rainfed plus potential access to groundwater seep, depth to groundwater >5m<br><br>Aspect: NE.            | Site ripped and sprayed the year before. Previously used for cropping/grazing.<br><br>Seedlings planted in contoured rows perpendicular from seep using a tree planter. No thinnings.<br><br>No reported problems with pests/weeds/disease, fire or drought.<br><br>Rainfed, plus potential access to groundwater. |
| 14  | Belka East                  | 1988 | 2 ha                          | See above, except only<br><br>Red Ironbark ( <i>E. sideroxylon</i> )   | 30 <sup>1</sup><br>22 <sup>2</sup> | 9.5  |          | N/A                            | N/A                | See above except:<br><br>No access to groundwater seep, greater depth.   | See above except:<br><br>Lower growth rates associated with droughting.  |
| 15  | Bakers Hill                 | 1978 | N/A (larger than those below) | River Red Gum ( <i>E. camaldulensis</i> )<br><br>Erosion control, watertable/dryland salinity control, experiment  | 38 <sup>1</sup><br>282             | 19.5 |          | Originally 550, now much lower | N/A                | Rainfall: Av 593 mm<br><br>Soil: Brownish black gravelly sand over gravelly yellow orange sand, pH unknown.<br><br>Topography: Very gently sloping low hillside.<br><br>Water: Rainfed, depth to groundwater >2m<br><br>Aspect: Unknown. | Site preparation unknown. Previously used for grazing.<br><br>Trees planted in rows. Thinnings occurred several years ago.<br><br>History of problems with pests/weeds/disease, fire or drought unknown.<br><br>Rainfed only.  |
| 16  | Bakers Hill                 | 1977 | 1 ha                          | See above  | 46 <sup>1</sup><br>35 <sup>2</sup> | 20.5 |          | 650                            | N/A                | See above except:<br><br>Soil: Dark brown gravelly loamy sand over a gravelly clayey sand.   | See above, except:<br><br>No thinnings.  |
| 17  | Bakers Hill                 | 1984 | 1.5 ha                        | See above  | 30 <sup>1</sup><br>22 <sup>2</sup> | 13.5 |          | 260                            | N/A                | See above except:<br><br>Soil: Dark brown loamy sand over yellowish brown light clay with mottles. Adjacent saline areas.<br><br>Water: Rainfed, depth to groundwater <2m.   | See above  |

| Site Information (Western Australia, Hassall & Associates, 1999b) |                             |      |              |   |                                      |      |             |                             |                           | Husbandry  |  |
|---|-----------------------------|------|--------------|---|--------------------------------------|------|-------------|-----------------------------|---------------------------|--|--|
| ID  | Name, Location, Coordinates | Est. | Area Planted | Species List & Purpose  | Biomass t/ha                         | Age  | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity   | e.g. tubestock/direct seed, weed & pest control, fire, irrigation  |
| 18  | North Bannister             | 1976 | 0.75 ha      | River Red Gum ( <i>E. camaldulensis</i> )<br><br>Erosion control, watertable/dryland salinity control, experiment | 197 <sup>1</sup><br>148 <sup>2</sup> | 21.5 |             | 815                         | N/A                       | Rainfall: Av 760 mm<br><br>Soil: Duplex with sandy gravel over sandy clay, impenetrable layers in parts. pH unknown.<br><br>Topography: Gentle midslope.<br><br>Water: Rainfed. Depth to groundwater >2.8m<br><br>Aspect: W. | Site preparation unknown. Previously used for grazing.<br><br>Trees planted in randomised blocks. No thinnings.<br><br>No reported problems with pests/weeds/disease, fire or drought.<br><br>Rainfed, plus potential access to groundwater. |
| 19  | North Bannister             | 1976 | 0.75 ha      | Blue Gum ( <i>E. globulus</i> )<br><br>Erosion control, watertable/dryland salinity control                       | 531 <sup>1</sup><br>398 <sup>2</sup> | 21.5 |             | 815                         | N/A                       | See above  | See above  |
| 20  | North Bannister             | 1976 | 1.5 ha       | River Red Gum ( <i>E. camaldulensis</i> )<br><br>Erosion control, watertable/dryland salinity control             | 146 <sup>1</sup><br>110 <sup>2</sup> | 21.5 |             | 815                         | N/A                       | See above  | See above except:<br><br>Survival rate poor (66%), apparently due to impenetrable layer in soil horizon and droughting.  |
| 21  | Dryandra                    | 1976 | 1.5 ha       | River Red Gum ( <i>E. camaldulensis</i> )<br><br>Erosion control, watertable/dryland salinity control, experiment | 43 <sup>1</sup><br>32 <sup>2</sup>   | 21.5 |             | 815                         | N/A                       | Rainfall: Av 530 mm<br><br>Soil: Sandy loam with gravel over light to dense brown clay, pH unknown.<br><br>Topography: Gentle slope.<br><br>Water: Rainfed. Depth to groundwater >1.6m.<br><br>Aspect: N.                    | Site preparation unknown. Previously used for grazing.<br><br>Trees planted in randomised blocks. No thinnings.<br><br>No reported problems with pests/weeds/disease, fire or drought.<br><br>Rainfed, plus potential access to groundwater. |
| 22  | Dryandra                    | 1976 | 1.5 ha       | See above   | 80 <sup>1</sup><br>60 <sup>2</sup>   | 21.5 |             | 815                         | N/A                       | See above  | See above  |

A4 - Table 1c. Growth Data for Sites in Western Australia, Hassall &amp; Associates (1999b) continued.

| Site Information (Western Australia, Hassall & Associates, 1999b) |                             |      |              |  |                                     |     |             |                             |                           |  |  |
|---|-----------------------------|------|--------------|--|-------------------------------------|-----|-------------|-----------------------------|---------------------------|--|--|
| ID  | Name, Location, Coordinates | Est. | Area Planted | Species List & Purpose   | Biomass t/ha                        | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity   | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation  |
| 23  | Bingham River               | 1976 | N/A          | Blue Gum ( <i>E. globulus</i> )<br><br>Erosion control, watertable/dryland salinity control                          | 83 <sup>1</sup><br>62 <sup>2</sup>  | 20  |             | N/A                         | N/A                       | Rainfall: Av 713 mm<br><br>Soil: Sand over sandy clay, pH unknown.<br><br>Topography: Gentle slope.<br><br>Water: Rainfed. Depth to groundwater unknown.   | Site preparation unknown. Previously used for mixed farming.<br><br>Establishment and management regime unknown.   |
| 24  | Bingham River               | 1978 | N/A          | See above  | 69 <sup>1</sup><br>52 <sup>2</sup>  | 18  |             | N/A                         | N/A                       | See above  | See above  |
| 25  | Bowelling                   | 1981 | N/A          | River Red Gum ( <i>E. camaldulensis</i> )<br><br>Erosion control, watertable/dryland salinity control, wood products | 113 <sup>1</sup><br>85 <sup>2</sup> | 14  |             | 600                         | N/A                       | Rainfall: Av 692 mm<br><br>Soil: Heavy clay, moderately saline (EC 140m S/m), pH unknown.<br><br>Topography: Flat.<br><br>Water: Rainfed, plus possible access to groundwater. Depth to groundwater <1.5m. Water table salinity 2000 mS/m. | Site preparation unknown. Previously used for grazing.<br><br>Trees planted in mounded rows. Thinnings history unknown.<br><br>No reported problems with pests/weeds/disease, fire or drought.<br><br>Rainfed, plus potential access to groundwater. |
| 26  | Cordinger                   | 1984 | N/A          | Blue Gum ( <i>E. globulus</i> )<br><br>Erosion control, watertable/dryland salinity control, wood products           | 121 <sup>1</sup><br>91 <sup>2</sup> | 12  |             | 375                         | N/A                       | Rainfall: Av 621 mm<br><br>Soil: N/A<br><br>Topography: Very gentle slope to flat.<br><br>Water: Rainfed. Depth to groundwater unknown.  | Site preparation unknown. Previously used for grazing.<br><br>Trees planted in rows. Thinnings history unknown.<br><br>No reported problems with pests/weeds/disease, fire or drought.<br><br>Rainfed.   |
| 27  | Cordinger                   | 1984 | N/A          | River Red Gum ( <i>E. camaldulensis</i> )  | 36 <sup>1</sup><br>27 <sup>2</sup>  | 12  |             | 325                         | N/A                       | See above  | See above except:<br><br>Trees planted on a salt scald and barley grass site.  |

| Site Information (Western Australia, Hassall & Associates, 1999b) |                             |      |              |  | Biomass                              | MAI  | Planting | Volume             | Site Factors       | Husbandry  |   |
|---|-----------------------------|------|--------------|--|--------------------------------------|------|----------|--------------------|--------------------|--|---|
| ID  | Name, Location, Coordinates | Est. | Area Planted | Species List & Purpose   | t/ha                                 | Age  | t/ha/yr  | Density (trees/ha) | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity  | e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 28  | Dinninup                    | 1983 | < 1 ha       | Blue Gum<br>( <i>E. globulus</i> )<br><br>Erosion control, waterable/dryland salinity control, wind break        | 83 <sup>1</sup><br>62 <sup>2</sup>   | 14.5 |          | 1,100              | N/A                | Rainfall: Av 601 mm<br><br>Soil: Black loam over sandy clay loam, pH unknown.<br><br>Topography: Gentle midslope.<br><br>Water: Rainfed. Depth to groundwater >2m although it is wet in winter.<br><br>Aspect: E.                      | Site ripped and sprayed but not mounded prior to planting. Previously used for grazing.<br><br>Seedlings hand planted in irregularly spaced rows. Thinnings history unknown.<br><br>No reported problems with weeds/disease or fire. Some initial deaths from kangaroos and some recent extensive deaths due to droughting.<br><br>Rainfed. |
| 29  | Dinninup                    | 1984 | < 1 ha       | River Red Gum<br>( <i>E. camaldulensis</i> )   | 77 <sup>1</sup><br>58 <sup>2</sup>   | 13.5 |          | 1,100              | N/A                | See above  | See above except:<br><br>No deaths associated with droughting.  |
| 30  | Dinninup                    | 1983 | < 1 ha       | Blue Gum<br>( <i>E. globulus</i> )   | 146 <sup>1</sup><br>109 <sup>2</sup> | 14.5 |          | 900                | N/A                | See above except:<br><br>Soil: Grey sand over yellow brown loamy sand.   | See above   |
| 31  | Mayanup                     | 1987 | 5 ha         | River Red Gum<br>( <i>E. camaldulensis</i> )<br><br>Erosion control, waterable/dryland salinity control, amenity | 76 <sup>1</sup><br>57 <sup>2</sup>   | 8.5  |          | 500                | N/A                | Rainfall: Av 668 mm<br><br>Soil: Shallow sand over clay, moderately saline (EC 50-140 mS/m), pH unknown.<br><br>Topography: Gentle slope to flat.<br><br>Water: Rainfed. Shallow, depth unknown, saline (2,200 mS/m)<br><br>Aspect: N. | Site ripped and mounded prior to planting. Previously use unknown.<br><br>Thinnings history unknown.<br><br>No reported problems with pests/weeds/disease, drought or fire.<br><br>Rainfed, plus access to groundwater.   |

<sup>5</sup> measured using bole and branches  
<sup>6</sup> measured using bole only

**A4 - Table 2a. Growth Data for Sites in South Australia, Wong et al. (2000)**

| Site information (South Australia, Wong et al., 2000) |   | Biomass |              | MAI                                       |      | Planting |         | Volume             |                    | Site Factors  |  | Husbandry   |  |
|---|---|---------|--------------|---|------|----------|---------|--------------------|--------------------|---|--|---|--|
| ID  | Name, Location, Coordinates                           | Est.    | Area Planted | Species List & Purpose                    | t/ha | Age      | t/ha/yr | Density (trees/ha) | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity                                   |  | e.g. tubestock/direct seed, weed & pest control, fire, irrigation |  |
| 1   | RT123C, Mt. Lofty Ranges<br><br>Lat/long: 3453, 13849 | 1988    | N/A          | <i>E. globulus</i><br>Southern Blue Gum   |      | 3        | N/A     | N/A                | 4                  | Rainfall: Av 926 mm<br><br>Soil: Loam over medium clay, depth to clay 30 cm<br><br>Topography: 20% slope. |  | Previously planted with <i>Pinus radiata</i> .                    |  |
|   | See above   |         |              |   |      | 5        |         |                    | 42                 |   |  |   |  |
|   | See above   |         |              |   |      | 7        |         |                    | 108                |   |  |   |  |
|   | See above   |         |              |   |      | 9        |         |                    | 200                |   |  |   |  |
|   | See above   |         |              |   |      | 11       |         |                    | 32                 |   |  |   |  |
|   | See above   |         | N/A          | <i>E. nitens</i><br>Shining Gum           |      | 3        | N/A     | N/A                | 5                  |   |  |   |  |
|   | See above   |         |              |   |      | 5        |         |                    | 65                 |   |  |   |  |
|   | See above   |         |              |   |      | 7        |         |                    | 151                |   |  |   |  |
|   | See above   |         |              |   |      | 9        |         |                    | 268                |   |  |   |  |
|   | See above   |         |              |   |      | 11       |         |                    | 390                |   |  |   |  |
|   | See above   |         | N/A          | <i>E. viminalis</i><br>Manna Gum          |      | 3        | N/A     | N/A                | 1                  |   |  |   |  |
|   | See above   |         |              |   |      | 5        |         |                    | 32                 |   |  |   |  |
|   | See above   |         |              |   |      | 7        |         |                    | 89                 |   |  |   |  |
|   | See above   |         |              |   |      | 9        |         |                    | 173                |   |  |   |  |
|   | See above   |         |              |   |      | 11       |         |                    | 288                |   |  |   |  |
|   | See above   |         | N/A          | <i>E. botryoides</i><br>Southern Mahogany |      | 3        | N/A     | N/A                | 4                  |   |  |   |  |
|   | See above   |         |              |   |      | 5        |         |                    | 37                 |   |  |   |  |
|   | See above   |         |              |   |      | 7        |         |                    | 91                 |   |  |   |  |



| Site information (South Australia, Wong <i>et. al.</i> , 2000) |   |      |              |   |              |     |             |                             |                           |  |   |
|--|---|------|--------------|---|--------------|-----|-------------|-----------------------------|---------------------------|--|---|
| ID   | Name, Location, Coordinates                           | Est. | Area Planted | Species List & Purpose                  | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity                 | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation |
|  | RT123C, Mt. Lofty Ranges                              |      |              |   |              | 9   |             |                             | 164                       |  |   |
|  | See above   |      | N/A          | <i>E. grandis</i><br>Flooded Gum        |              | 3   | N/A         | N/A                         | 2                         |  |   |
|  | See above   |      |              |   |              | 5   |             |                             | 37                        |  |   |
|  | See above   |      |              |   |              | 7   |             |                             | 88                        |  |   |
|  | See above   |      |              |   |              | 9   |             |                             | 150                       |  |   |
|  | See above   |      |              |   |              | 11  |             |                             | 232                       |  |   |
|  | See above   |      | N/A          | <i>E. saligna</i><br>Sydney Blue Gum    |              | 3   | N/A         | N/A                         | 2                         |  |   |
|  | See above   |      |              |   |              | 5   |             |                             | 35                        |  |   |
|  | See above   |      |              |   |              | 7   |             |                             | 85                        |  |   |
|  | See above   |      |              |   |              | 9   |             |                             | 162                       |  |   |
|  | See above   |      |              |   |              | 11  |             |                             | 240                       |  |   |
| 2  | EP205, NW of Mt. Gambier<br><br>Lat/long: 3743, 14037 | 1988 | N/A          | <i>E. globulus</i><br>Southern Blue Gum |              | 2   | N/A         | N/A                         | 11                        | Rainfall: Av 766 mm<br><br>Soit: Sand over medium clay, depth to clay 70 cm<br><br>Topography: Flat. | Previously used as pasture for grazing.                                     |
|  | See above   |      |              |   |              | 3   |             |                             | 37                        |  |   |
|  | See above   |      |              |   |              | 4   |             |                             | 60                        |  |   |
|  | See above   |      |              |   |              | 6   |             |                             | 130                       |  |   |
|  | See above   |      |              |   |              | 9   |             |                             | 196                       |  |   |
|  | See above   |      |              |   |              | 11  |             |                             | 279                       |  |   |
|  | See above   |      | N/A          | <i>E. nitens</i><br>Shining Gum         |              | 2   | N/A         | N/A                         | 5                         |  |   |
|  | See above   |      |              |   |              | 3   |             |                             | 20                        |  |   |
|  | See above   |      |              |   |              | 4   |             |                             | 39                        |  |   |

A4 - Table 2a. Growth Data for Sites in South Australia, Wong *et al.* (2000) continued.

| Site ID   | Site information (South Australia, Wong <i>et al.</i> , 2000) | Area Est. | Species List & Purpose                    | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity                   | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation |
|-----------|---|-----------|---|--------------|-----|-------------|-----------------------------|---------------------------|--|---|
| See above |   |           |   |              | 6   |             |                             | 93                        |  |   |
| See above |   |           |   |              | 9   |             |                             | 147                       |  |   |
| See above |   | N/A       | <i>E. viminalis</i><br>Manna Gum          |              | 2   | N/A         | N/A                         | 4                         |  |   |
| See above |   |           |   |              | 3   |             |                             | 20                        |  |   |
| See above |   |           |   |              | 4   |             |                             | 37                        |  |   |
| See above |   |           |   |              | 6   |             |                             | 80                        |  |   |
| See above |   |           |   |              | 9   |             |                             | 123                       |  |   |
| See above |   | N/A       | <i>E. botryoides</i><br>Southern Mahogany |              | 2   | N/A         | N/A                         | 5                         |  |   |
| See above |   |           |   |              | 3   |             |                             | 20                        |  |   |
| See above |   |           |   |              | 4   |             |                             | 35                        |  |   |
| See above |   |           |   |              | 6   |             |                             | 92                        |  |   |
| See above |   |           |   |              | 9   |             |                             | 147                       |  |   |
| 3         | EM133, N of Mt. Gambier<br>Lat/long: 3745, 14047              | 1991      | <i>E. globulus</i><br>Southern Blue Gum   |              | 2   | N/A         | N/A                         | 3                         | Rainfall: Av 745 mm<br><br>Soil: Uniform sandy profile, depth to clay 200 cm+<br><br>Topography: Flat. | Previously used as pasture for grazing.                                     |
| See above |   |           |   |              | 3   |             |                             | 21                        |  |   |
| See above |   |           |   |              | 4   |             |                             | 48                        |  |   |
| See above |   |           |   |              | 5   |             |                             | 69                        |  |   |
| See above |   | N/A       | <i>E. saligna</i><br>Sydney Blue Gum      |              | 3   | N/A         | N/A                         | 4                         |  |   |
| See above |   |           |   |              | 4   |             |                             | 15                        |  |   |
| See above |   |           |   |              | 5   |             |                             | 24                        |  |   |

| Site information (South Australia, Wong <i>et. al.</i> , 2000) |                             |      |              |                                  |              |     |             |                             |                           |  |   |
|--|-----------------------------|------|--------------|----------------------------------|--------------|-----|-------------|-----------------------------|---------------------------|--|---|
| ID   | Name, Location, Coordinates | Est. | Area Planted | Species List & Purpose           | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation |
|  | See above                   |      | N/A          | <i>E. grandis</i><br>Flooded Gum |              | 2   | N/A         | N/A                         | 1                         |  |   |
|  | EM133, N of Mt. Gambier     |      |              |                                  |              | 3   |             |                             | 13                        |  |   |
|  | See above                   |      |              |                                  |              | 4   |             |                             | 37                        |  |   |
|  | See above                   |      |              |                                  |              | 5   |             |                             | 51                        |  |   |

**A4 - Table 2b. Growth Data for Sites in Victoria, Wong *et al.* (2000)**

| Site information (Victoria, Wong <i>et. al.</i> , 2000) |  |      |              |   |              |     |             |                             |                           |   |   |
|---|--|------|--------------|---|--------------|-----|-------------|-----------------------------|---------------------------|---|---|
| ID  | Name, Location, Coordinates                                    | Est. | Area Planted | Species List & Purpose                    | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity          | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 1   | Tostaree, near Lakes Entrance, Vic.<br>Lat/long: 3747S, 14811E | 1988 | N/A          | <i>E. globulus</i><br>Southern Blue Gum   |              | 2   | N/A         | 951                         | 13                        | Rainfall: Av 819 mm<br><br>Soil: Loamy sand over light medium clay<br>Topography: 3-6% slope. | Small scale plot (<1 ha).<br><br>Previously used for grazing on improved pasture with a history of relatively high rates of fertiliser application. Direct seeding. |
|   | See above  |      |              |   |              | 4   |             | 942                         | 74                        |   |   |
|   | See above  |      |              |   |              | 6   |             | 924                         | 157                       |   |   |
|   | See above  |      |              |   |              | 8   |             | 884                         | 262                       |   |   |
|   | See above  |      |              |   |              | 10  |             | 840                         | 334                       |   |   |
| 2   | Tostaree   |      |              | <i>E. nitens</i><br>Shining Gum           |              | 2   | N/A         | 920                         | 13                        |   |   |
|   | See above  |      |              |   |              | 4   |             | 916                         | 76                        |   |   |
|   | See above  |      |              |   |              | 6   |             | 900                         | 147                       |   |   |
|   | See above  |      |              |   |              | 8   |             | 860                         | 237                       |   |   |
|   | See above  |      |              |   |              | 10  |             | 832                         | 287                       |   |   |
| 3   | Tostaree   |      |              | <i>E. viminalis</i><br>Manna Gum          |              | 2   | N/A         | 980                         | 12                        |   |   |
|   | See above  |      |              |   |              | 4   |             | 904                         | 81                        |   |   |
|   | See above  |      |              |   |              | 6   |             | 892                         | 166                       |   |   |
|   | See above  |      |              |   |              | 8   |             | 876                         | 262                       |   |   |
|   | See above  |      |              |   |              | 10  |             | 872                         | 319                       |   |   |
| 4   | Tostaree   |      |              | <i>E. botryoides</i><br>Southern Mahogany |              | 2   | N/A         | 964                         | 8                         |   |   |
|   | See above  |      |              |   |              | 4   |             | 960                         | 49                        |   |   |
|   | See above  |      |              |   |              | 6   |             | 952                         | 103                       |   |   |
|   | See above  |      |              |   |              | 8   |             | 936                         | 171                       |   |   |
|   | See above  |      |              |   |              | 10  |             | 888                         | 216                       |   |   |

| Site information (Victoria, Wong <i>et al.</i> , 2000) |   |      |              |   |              |     |             |                             |                           |  |   |
|--|---|------|--------------|---|--------------|-----|-------------|-----------------------------|---------------------------|--|---|
| ID   | Name, Location, Coordinates                           | Est. | Area Planted | Species List & Purpose                  | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 5  | Tostaree  |      |              | <i>E. grandis</i><br>Flooded Gum        |              | 2   | N/A         | 980                         | 8                         |  |   |
|  | See above   |      |              |   |              | 4   |             | 972                         | 45                        |  |   |
|  | See above   |      |              |   |              | 6   |             | 972                         | 93                        |  |   |
|  | See above   |      |              |   |              | 8   |             | 960                         | 161                       |  |   |
|  | See above   |      |              |   |              | 10  |             | 940                         | 215                       |  |   |
| 6  | Tostaree  |      |              | <i>E. saligna</i><br>Sydney Blue Gum    |              | 2   | N/A         | 960                         | 8                         |  |   |
|  | See above   |      |              |   |              | 4   |             | 948                         | 46                        |  |   |
|  | See above   |      |              |   |              | 6   |             | 948                         | 95                        |  |   |
|  | See above   |      |              |   |              | 8   |             | 928                         | 154                       |  |   |
|  | See above   |      |              |   |              | 10  | N/A         | 908                         | 195                       |  |   |
| 7  | Waygara, near Orbost, Vic.<br>Lat/long: 3741S, 14819E | 1989 | N/A          | <i>E. globulus</i><br>Southern Blue Gum |              | 2   | N/A         | 938                         | 10                        | Rainfall: Av 870 mm<br>Soil: Loam over light medium clay<br>Topography: Flat.        | Small scale plot (<1 ha).<br>Previous land use degraded native forest severely affected by fungal pathogen <i>Phytophthora cinnamomi</i> . Cleared before direct seeding. |
|  | See above   |      |              |   |              | 4   |             | 938                         | 65                        |  |   |
|  | See above   |      |              |   |              | 6   |             | 917                         | 115                       |  |   |
|  | See above   |      |              |   |              | 8   |             | 892                         | 159                       |  |   |
|  | See above   |      |              |   |              | 10  |             | 858                         | 181                       |  |   |
| 8  | Waygara   |      |              | <i>E. nitens</i><br>Shining Gum         |              | 2   |             | 695                         | 5                         |  |   |
|  | See above   |      |              |   |              | 4   |             | 685                         | 41                        |  |   |
|  | See above   |      |              |   |              | 6   |             | 662                         | 70                        |  |   |
|  | See above   |      |              |   |              | 8   |             | 643                         | 87                        |  |   |
|  | See above   |      |              |   |              | 10  |             | 656                         | 114                       |  |   |

A4 - Table 2b. Growth Data for Sites in Victoria, Wong *et al.* (2000) continued.

| Site information (Victoria, Wong <i>et al.</i> , 2000) |                             | Species List & Purpose                    | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation |
|--|-----------------------------|---|--------------|-----|-------------|-----------------------------|---------------------------|--|---|
| ID   | Name, Location, Coordinates |   |              |     |             |                             |                           |  |   |
| 9  | Waygara                     | <i>E. viminalis</i><br>Manna Gum          | 2            | 2   | 713         | 5                           |                           |  |   |
|  | See above                   |   | 4            | 4   | 696         | 38                          |                           |  |   |
|  | See above                   |   | 6            | 6   | 688         | 74                          |                           |  |   |
|  | See above                   |   | 8            | 8   | 683         | 103                         |                           |  |   |
|  | See above                   |   | 10           | 10  | 679         | 109                         |                           |  |   |
| 10   | Waygara                     | <i>E. botryoides</i><br>Southern Mahogany | 2            | 2   | 963         | 5                           |                           |  |   |
|  | See above                   |   | 4            | 4   | 963         | 24                          |                           |  |   |
|  | See above                   |   | 6            | 6   | 958         | 64                          |                           |  |   |
|  | See above                   |   | 8            | 8   | 938         | 86                          |                           |  |   |
|  | See above                   |   | 10           | 10  | 938         | 102                         |                           |  |   |
| 11   | Waygara                     | <i>E. grandis</i><br>Flooded Gum          | 2            | 2   | 911         | 5                           |                           |  |   |
|  | See above                   |   | 4            | 4   | 911         | 30                          |                           |  |   |
|  | See above                   |   | 6            | 6   | 911         | 68                          |                           |  |   |
|  | See above                   |   | 8            | 8   | 894         | 92                          |                           |  |   |
|  | See above                   |   | 10           | 10  | 886         | 95                          |                           |  |   |
| 12   | Waygara                     | <i>E. saligna</i><br>Sydney Blue Gum      | 2            | 2   | 954         | 6                           |                           |  |   |
|  | See above                   |   | 4            | 4   | 946         | 30                          |                           |  |   |
|  | See above                   |   | 6            | 6   | 921         | 69                          |                           |  |   |
|  | See above                   |   | 8            | 8   | 904         | 92                          |                           |  |   |
|  | See above                   |   | 10           | 10  | 896         | 105                         |                           |  |   |



| Site information (Victoria, Wong <i>et. al.</i> , 2000) |                                    |                                   |              |   |   |     |             |                             |                           |  |   |
|---|------------------------------------|-----------------------------------|--------------|---|---|-----|-------------|-----------------------------|---------------------------|--|---|
| ID  | Name, Location, Coordinates        | Est.                              | Area Planted | Species List & Purpose                  | Biomass t/ha                            | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 13  | Mt Worth West, near Warragul, Vic. | 1986                              | N/A          | <i>E. globulus</i><br>Southern Blue Gum |   | 3.5 | N/A         | 800                         | 31                        | Rainfall: Av 1212 mm   | Small scale plot (<1 ha).   |
|   | Lat/long: 3818S, 14559E            |                                   |              |   |   |     |             |                             |                           | Soil: Clay loam over light medium clay   | Previously used for grazing on improved pasture with a history of relatively high rates of fertiliser application. Cleared before direct seeding. |
|   | See above                          |                                   |              |   |   | 6   |             | 831                         | 135                       |  |   |
|   | Mt Worth West                      |                                   |              |   |   | 8.5 |             | 800                         | 227                       |  |   |
|   | See above                          |                                   |              |   |   | 12  |             | 631                         | 401                       |  |   |
|   | See above                          |                                   |              | <i>E. nitens</i><br>Shining Gum         |   | 3.5 |             | 970                         | 48                        |  |   |
|   | See above                          |                                   |              |   |   | 6   |             | 954                         | 206                       |  |   |
|   | See above                          |                                   |              |   |   | 8.5 |             | 970                         | 356                       |  |   |
|   | See above                          |                                   |              |   |   | 12  |             | 818                         | 558                       |  |   |
|   | 14                                 | Narracan East, near Morwell, Vic. | 1986         | N/A                                     | <i>E. globulus</i><br>Southern Blue Gum |     | 3.5         | N/A                         | 927                       | 33   | Rainfall: Av 963 mm   |
| Lat/long: 3817S, 14616E                                 |                                    |                                   |              |   |   |     |             |                             |                           | Soil: Loam over medium clay  | Previous landuse <i>Pinus radiata</i> .   |
| See above   |                                    |                                   |              |   |   | 6   |             | 889                         | 94                        | Topography: 2-3% slope.  | Cleared before direct seeding.  |
| See above   |                                    |                                   |              |   |   | 12  |             | 800                         | 365                       |  |   |
| See above   |                                    |                                   |              | <i>E. nitens</i><br>Shining Gum         |   | 3.5 |             | 912                         | 29                        |  |   |
|   | See above                          |                                   |              |   |   | 6   |             | 902                         | 85                        |  |   |
|   | See above                          |                                   |              |   |   | 12  |             | 776                         | 305                       |  |   |
|   | 15                                 | Yinnar, near Morwell, Vic.        | 1986         | N/A                                     | <i>E. globulus</i><br>Southern Blue Gum |     | 3.5         | N/A                         | 970                       | 8  | Rainfall: Av 930 mm   |
| Lat/long: 3818S, 14618E                                 |                                    |                                   |              |   |   |     |             |                             |                           | Soil: Shallow sandy loam over medium heavy clay                                      | Previous landuse <i>Pinus radiata</i> .   |
|   |                                    |                                   |              |   |   |     |             |                             |                           | Topography: 0-5% slope.  | Low growth rates experienced at this site due to shallow surface soils.   |
|   |                                    |                                   |              |   |   |     |             |                             |                           |  | Cleared before direct seeding.  |
| See above   |                                    |                                   |              |   |   | 6   |             | 929                         | 38                        |  |   |
| See above   |                                    |                                   |              |   | 12                                      |     | 866         | 152                         |                           |  |   |

A4 - Table 2b. Growth Data for Sites in Victoria, Wong *et al.* (2000) continued.

| Site information (Victoria, Wong <i>et al.</i> , 2000) |   | Species List & Area Planted |              | Biomass                                 | Age     | MAI     | Planting           | Volume             | Site Factors  | Husbandry  |
|--|---|-----------------------------|--------------|---|---------|---------|--------------------|--------------------|---|--|
| ID   | Name, Location, Coordinates   | Est.                        | Area Planted | t/ha                                    | t/ha/yr | t/ha/yr | Density (trees/ha) | m <sup>3</sup> /ha | e.g. rainfall, soil type/quality, topography, access to water, salinity               | e.g. tubestock/direct seed, weed & pest control, fire, irrigation  |
|  | See above   |                             |              | <i>E. nitens</i><br>Shining Gum         | 3.5     |         | 800                | 6                  |   |  |
|  | See above   |                             |              |   | 6       |         | 803                | 28                 |   |  |
|  | See above   |                             |              |   | 12      |         | 707                | 81                 |   |  |
| 16   | Maryvale,<br>N of Morwell, Vic.<br><br>Lat/long: 3812S, 14628E                    | 1986                        | N/A          | <i>E. globulus</i><br>Southern Blue Gum | 3.5     | N/A     | 932                | 20                 | Rainfall: Av 768 mm<br><br>Soil: Sandy loam over medium clay<br><br>Topography: Flat. | Small scale plot (<1 ha).<br><br>Previous landuse <i>Pinus radiata</i> .<br><br>Cleared before direct seeding. |
|  | See above   |                             |              |   | 6       |         | 888                | 65                 |   |  |
|  | See above   |                             |              |   | 12      |         | 883                | 214                |   |  |
|  | See above   |                             |              | <i>E. nitens</i><br>Shining Gum         | 3.5     |         | 966                | 13                 |   |  |
|  | See above   |                             |              |   | 6       |         | 966                | 41                 |   |  |
|  | See above   |                             |              |   | 12      |         | 638                | 87                 |   |  |
| 17   | Gormondale,<br>between Sale and Traralgon,<br>Vic.<br><br>Lat/long: 3816S, 14642E | 1986                        | N/A          | <i>E. globulus</i><br>Southern Blue Gum | 3.5     | N/A     | 992                | 31                 | Rainfall: Av 830 mm<br><br>Soil: Sand over loamy sand<br><br>Topography: 0-5% slope.  | Small scale plot (<1 ha).<br><br>Previous landuse <i>Pinus radiata</i> .<br><br>Cleared before direct seeding. |
|  | See above   |                             |              |   | 6       |         | 992                | 86                 |   |  |
|  | See above   |                             |              |   | 12      |         | 868                | 216                |   |  |
|  | See above   |                             |              | <i>E. nitens</i><br>Shining Gum         | 3.5     |         | 979                | 20                 |   |  |
|  | See above   |                             |              |   | 6       |         | 966                | 58                 |   |  |
|  | See above   |                             |              |   | 12      |         | 916                | 135                |   |  |

| Site information (Victoria, Wong <i>et. al.</i> , 2000) |   |      |              |   |              |     |             |                             |                           |   |   |
|---|---|------|--------------|---|--------------|-----|-------------|-----------------------------|---------------------------|---|---|
| ID  | Name, Location, Coordinates                                       | Est. | Area Planted | Species List & Purpose                    | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity                  | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation   |
| 18  | Mt Worth East, near Warragul, Vic.<br><br>Lat/long: 3819S, 14559E | 1987 | N/A          | <i>E. globulus</i><br>Southern Blue Gum   |              | 3.5 | N/A         | 884                         | 27                        | Rainfall: Av 1224 mm<br><br>Soil: Clay loam over light to medium clay<br><br>Topography: 5-28% slope. | Small scale plot (<1 ha).<br><br>Previously used for grazing on improved pasture with a history of relatively high rates of fertiliser application. Unrealistically high productivity experienced at this site. Not considered representative of broad-scale plantations.<br><br>Cleared before direct seeding. |
|   | Mt Worth East   |      |              |   |              | 6   |             | 876                         | 114                       |   |   |
|   | See above   |      |              |   |              | 11  |             | 828                         | 340                       |   |   |
|   | See above   |      |              | <i>E. viminalis</i><br>Manna Gum          |              | 3.5 |             | 960                         | 32                        |   |   |
|   | See above   |      |              |   |              | 6   |             | 961                         | 138                       |   |   |
|   | See above   |      |              |   |              | 11  |             | 938                         | 435                       |   |   |
|   | See above   |      |              | <i>E. botryoides</i><br>Southern Mahogany |              | 3.5 |             | 884                         | 15                        |   |   |
|   | See above   |      |              |   |              | 6   |             | 857                         | 77                        |   |   |
|   | See above   |      |              |   |              | 11  |             | 764                         | 212                       |   |   |
|   | See above   |      |              | <i>E. grandis</i><br>Flooded Gum          |              | 3.5 |             | 846                         | 11                        |   |   |
|   | See above   |      |              |   |              | 6   |             | 856                         | 44                        |   |   |
|   | See above   |      |              |   |              | 11  |             | 720                         | 120                       |   |   |
|   | See above   |      |              | <i>E. saligna</i><br>Sydney Blue Gum      |              | 3.5 |             | 918                         | 23                        |   |   |
|   | See above   |      |              |   |              | 6   |             | 901                         | 101                       |   |   |
|   | See above   |      |              |   |              | 11  |             | 838                         | 281                       |   |   |

A4 - Table 2b. Growth Data for Sites in Victoria, Wong *et al.* (2000) continued.

| Site information (Victoria, Wong <i>et. al.</i> , 2000) |  |      |              |   |              |     |             |                             |                           |  |  |
|---|--|------|--------------|---|--------------|-----|-------------|-----------------------------|---------------------------|--|--|
| ID  | Name, Location, Coordinates                                  | Est. | Area Planted | Species List & Purpose                    | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity   | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation                                    |
| 19  | Delburn, near Leongatha, Vic.<br><br>Lat/long: 3821S, 14614E | 1987 | N/A          | <i>E. globulus</i><br>Southern Blue Gum   |              | 3.5 | N/A         | 935                         | 15                        | Rainfall: Av 1002 mm<br><br>Soil: Loam over medium clay<br><br>Topography: 0-8% slope. | Small scale plot (<1 ha).<br><br>Previous landuse <i>Pinus radiata</i> .<br><br>Cleared before direct seeding. |
|   | See above  |      |              |   |              | 6   |             | 935                         | 84                        |  |  |
|   | See above  |      |              |   |              | 11  |             | 917                         | 316                       |  |  |
|   | See above  |      |              | <i>E. nitens</i><br>Shining Gum           |              | 3.5 |             | 817                         | 11                        |  |  |
|   | See above  |      |              |   |              | 6   |             | 817                         | 70                        |  |  |
|   | See above  |      |              |   |              | 11  |             | 817                         | 259                       |  |  |
|   | See above  |      |              | <i>E. viminalis</i><br>Manna Gum          |              | 3.5 |             | 876                         | 11                        |  |  |
|   | See above  |      |              |   |              | 6   |             | 850                         | 63                        |  |  |
|   | See above  |      |              |   |              | 11  |             | 850                         | 224                       |  |  |
|   | See above  |      |              | <i>E. botryoides</i><br>Southern Mahogany |              | 3.5 |             | 972                         | 8                         |  |  |
|   | See above  |      |              |   |              | 6   |             | 943                         | 57                        |  |  |
|   | See above  |      |              |   |              | 11  |             | 926                         | 211                       |  |  |
|   | See above  |      |              | <i>E. grandis</i><br>Flooded Gum          |              | 3.5 |             | 951                         | 9                         |  |  |
|   | See above  |      |              |   |              | 6   |             | 935                         | 54                        |  |  |
|   | See above  |      |              |   |              | 11  |             | 892                         | 183                       |  |  |
|   | See above  |      |              | <i>E. saligna</i><br>Sydney Blue Gum      |              | 3.5 |             | 978                         | 10                        |  |  |
|   | See above  |      |              |   |              | 6   |             | 962                         | 59                        |  |  |
|   | See above  |      |              |   |              | 11  |             | 935                         | 215                       |  |  |

| Site information (Victoria, Wong <i>et. al.</i> , 2000) |   |      |              |   |              |     |             |                             |                           |   |  |
|---|---|------|--------------|---|--------------|-----|-------------|-----------------------------|---------------------------|---|--|
| ID  | Name, Location, Coordinates   | Est. | Area Planted | Species List & Purpose                    | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity            | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation                                    |
| 20  | Flynns Creek, between Sale and Traralgon, Vic.<br><br>Lat/long: 38°16S, 146°36E | 1987 | N/A          | <i>E. globulus</i><br>Southern Blue Gum   |              | 3.5 | N/A         | 919                         | 13                        | Rainfall: Av 1173 mm<br><br>Soil: Loamy sand over light to medium clay<br><br>Topography: Flat. | Small scale plot (<1 ha).<br><br>Previous landuse <i>Pinus radiata</i> .<br><br>Cleared before direct seeding. |
|   | See above   |      |              |   | 6            |     |             | 910                         | 52                        |   |  |
|   | See above   |      |              |   | 11           |     |             | 884                         | 179                       |   |  |
|   | See above   |      |              | <i>E. nitens</i><br>Shining Gum           |              | 3.5 |             | 744                         | 10                        |   |  |
|   | See above   |      |              |   | 6            |     |             | 695                         | 44                        |   |  |
|   | See above   |      |              |   | 11           |     |             | 678                         | 139                       |   |  |
|   | See above   |      |              | <i>E. viminalis</i><br>Manna Gum          |              | 3.5 |             | 860                         | 9                         |   |  |
|   | Flynns Creek  |      |              |   | 6            |     |             | 860                         | 37                        |   |  |
|   | See above   |      |              |   | 11           |     |             | 835                         | 105                       |   |  |
|   | See above   |      |              | <i>E. botryoides</i><br>Southern Mahogany |              | 3.5 |             | 910                         | 9                         |   |  |
|   | See above   |      |              |   | 6            |     |             | 902                         | 40                        |   |  |
|   | See above   |      |              |   | 11           |     |             | 885                         | 136                       |   |  |
|   | See above   |      |              | <i>E. grandis</i><br>Flooded Gum          |              | 3.5 |             | 886                         | 8                         |   |  |
|   | See above   |      |              |   | 6            |     |             | 877                         | 34                        |   |  |
|   | See above   |      |              |   | 11           |     |             | 860                         | 98                        |   |  |
|   | See above   |      |              | <i>E. saligna</i><br>Sydney Blue Gum      |              | 3.5 |             | 851                         | 9                         |   |  |
|   | See above   |      |              |   | 6            |     |             | 842                         | 38                        |   |  |
|   | See above   |      |              |   | 11           |     |             | 786                         | 113                       |   |  |

A4 - Table 2b. Growth Data for Sites in Victoria, Wong *et al.* (2000) continued.

| Site information (Victoria, Wong <i>et al.</i> , 2000) |  |      |              | Biomass                                   |      | MAI | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors  |  | Husbandry  |  |
|--|--|------|--------------|---|------|-----|-----------------------------|---------------------------|---|--|--|--|
| ID   | Name, Location, Coordinates                                  | Est. | Area Planted | Species List & Purpose                    | t/ha | Age |                             |                           | e.g. rainfall, soil type/quality, topography, access to water, salinity |  | e.g. tubestock/direct seed, weed & pest control, fire, irrigation  |  |
| 21   | Stradbroke, near Sale, Vic.<br><br>Lat/long: 38°16S, 147°03E | 1987 | N/A          | <i>E. globulus</i><br>Southern Blue Gum   |      | 3.5 | N/A                         | 18                        | Rainfall: Av 598 mm<br><br>Soil: Sand<br><br>Topography: 2-7% slope.    |  | Small scale plot (<1 ha).<br><br>Previous landuse <i>Pinus radiata</i> .<br><br>Cleared before direct seeding. |  |
|  | See above  |      |              |   | 6    |     | 919                         | 54                        |   |  |  |  |
|  | See above  |      |              |   | 11   |     | 888                         | 106                       |   |  |  |  |
|  | See above  |      |              | <i>E. nitens</i><br>Shining Gum           |      | 3.5 | 934                         | 16                        |   |  |  |  |
|  | See above  |      |              |   | 6    |     | 927                         | 49                        |   |  |  |  |
|  | See above  |      |              |   | 11   |     | 876                         | 86                        |   |  |  |  |
|  | See above  |      |              | <i>E. viminalis</i><br>Manna Gum          |      | 3.5 | 951                         | 20                        |   |  |  |  |
|  | See above  |      |              |   | 6    |     | 943                         | 59                        |   |  |  |  |
|  | See above  |      |              |   | 11   |     | 898                         | 107                       |   |  |  |  |
|  | See above  |      |              | <i>E. botryoides</i><br>Southern Mahogany |      | 3.5 | 959                         | 30                        |   |  |  |  |
|  | See above  |      |              |   | 6    |     | 959                         | 97                        |   |  |  |  |
|  | See above  |      |              |   | 11   |     | 947                         | 257                       |   |  |  |  |
|  | See above  |      |              | <i>E. grandis</i><br>Flooded Gum          |      | 3.5 | 934                         | 23                        |   |  |  |  |
|  | See above  |      |              |   | 6    |     | 927                         | 74                        |   |  |  |  |
|  | See above  |      |              |   | 11   |     | 921                         | 172                       |   |  |  |  |
|  | See above  |      |              | <i>E. saligna</i><br>Sydney Blue Gum      |      | 3.5 | 935                         | 28                        |   |  |  |  |
|  | See above  |      |              |   | 6    |     | 927                         | 78                        |   |  |  |  |
|  | See above  |      |              |   | 11   |     | 924                         | 183                       |   |  |  |  |



| Site information (Victoria, Wong <i>et. al.</i> , 2000) |   |      |              |   |              |     |             |                             |                           |   |  |
|---|---|------|--------------|---|--------------|-----|-------------|-----------------------------|---------------------------|---|--|
| ID  | Name, Location, Coordinates                                 | Est. | Area Planted | Species List & Purpose                    | Biomass t/ha | Age | MAI t/ha/yr | Planting Density (trees/ha) | Volume m <sup>3</sup> /ha | Site Factors e.g. rainfall, soil type/quality, topography, access to water, salinity        | Husbandry e.g. tubestock/direct seed, weed & pest control, fire, irrigation                                    |
| 22  | Stockdale, N of Sale, Vic.<br><br>Lat/long: 37°51S, 147°11E | 1987 | N/A          | <i>E. globulus</i><br>Southern Blue Gum   |              | 3.5 | N/A         | 919                         | 19                        | Rainfall: Av 1182 mm<br><br>Soil: Loamy sand over medium heavy clay<br><br>Topography: Flat | Small scale plot (<1 ha).<br><br>Previous landuse <i>Pinus radiata</i> .<br><br>Cleared before direct seeding. |
|   | See above   |      |              |   |              | 6   |             | 902                         | 61                        |   |  |
|   | See above   |      |              |   |              | 11  |             | 877                         | 141                       |   |  |
|   | See above   |      |              | <i>E. nitens</i><br>Shining Gum           |              | 3.5 |             | 910                         | 17                        |   |  |
|   | See above   |      |              |   |              | 6   |             | 893                         | 52                        |   |  |
|   | See above   |      |              |   |              | 11  |             | 679                         | 84                        |   |  |
|   | See above   |      |              | <i>E. viminalis</i><br>Manna Gum          |              | 3.5 |             | 959                         | 18                        |   |  |
|   | See above   |      |              |   |              | 6   |             | 934                         | 59                        |   |  |
|   | See above   |      |              |   |              | 11  |             | 883                         | 117                       |   |  |
|   | See above   |      |              | <i>E. botryoides</i><br>Southern Mahogany |              | 3.5 |             | 919                         | 15                        |   |  |
|   | See above   |      |              |   |              | 6   |             | 919                         | 49                        |   |  |
|   | Stockdale   |      |              |   |              | 11  |             | 900                         | 102                       |   |  |
|   | See above   |      |              | <i>E. grandis</i><br>Flooded Gum          |              | 3.5 |             | 901                         | 12                        |   |  |
|   | See above   |      |              |   |              | 6   |             | 901                         | 43                        |   |  |
|   | See above   |      |              |   |              | 11  |             | 876                         | 89                        |   |  |
|   | See above   |      |              | <i>E. saligna</i><br>Sydney Blue Gum      |              | 3.5 |             | 911                         | 15                        |   |  |
|   | See above   |      |              |   |              | 6   |             | 893                         | 48                        |   |  |
|   | See above   |      |              |   |              | 11  |             | 829                         | 98                        |   |  |








## ***Series 1 Publications***

1. Setting the Frame
2. Estimation of Changes in Soil Carbon Due to Changes in Land Use
3. Woody Biomass: Methods for Estimating Change
4. Land Clearing 1970-1990: A Social History
- 5a. Review of Allometric Relationships for Estimating Woody Biomass for Queensland, the Northern Territory and Western Australia
- 5b. Review of Allometric Relationships for Estimating Woody Biomass for New South Wales, the Australian Capital Territory, Victoria, Tasmania and South Australia
6. The Decay of Coarse Woody Debris
7. Carbon Content of Woody Roots: Revised Analysis and a Comparison with Woody Shoot Components (Revision 1)
8. Usage and Lifecycle of Wood Products
9. Land Cover Change: Specification for Remote Sensing Analysis
10. National Carbon Accounting System: Phase 1 Implementation Plan for the 1990 Baseline
11. International Review of the Implementation Plan for the 1990 Baseline (13-15 December 1999)

## ***Series 2 Publications***

12. Estimation of Pre-Clearing Soil Carbon Conditions
13. Agricultural Land Use and Management Information
14. Sampling, Measurement and Analytical Protocols for Carbon Estimation in Soil, Litter and Coarse Woody Debris
15. Carbon Conversion Factors for Historical Soil Carbon Data
16. Remote Sensing Analysis Of Land Cover Change - Pilot Testing of Techniques
17. Synthesis of Allometrics, Review of Root Biomass and Design of Future Woody Biomass Sampling Strategies
18. Wood Density Phase 1 - State of Knowledge
19. Wood Density Phase 2 - Additional Sampling
20. Change in Soil Carbon Following Afforestation or Reforestation
21. System Design
22. Carbon Contents of Above-Ground Tissues of Forest and Woodland Trees
23. Developing a National Forest Productivity Model
24. Analysis of Wood Product Accounting Options for the National Carbon Accounting System
25. Review of Unpublished Biomass-Related Information: Western Australia, South Australia, New South Wales and Queensland
26. CAMFor User Manual



The National Carbon Accounting System provides a complete accounting and forecasting capability for human-induced sources and sinks of greenhouse gas emissions from Australian land based systems. It will provide a basis for assessing Australia's progress towards meeting its international emissions commitments.