

0.1 Continuous-time Fourier transform

Let us start with the definition of this very important tool

Definition 1: Fourier transform of a continuous-time signal

The CTFT of a continuous-time signal $x_a(t)$ is given by:

$$X_a(j\Omega) = \int_{-\infty}^{\infty} x_a(t) e^{-j\Omega t} dt \quad (1)$$

often referred to as the Fourier spectrum or simply the spectrum of the continuous-time signal.

Definition 2: Inverse Fourier transform of a continuous-time signal

The inverse CTFT of a Fourier transform $X_a(j\Omega)$ is given by:

$$x_a(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X_a(j\Omega) e^{+j\Omega t} d\Omega \quad (2)$$

often referred to as the Fourier integral.

A CTFT pair will be denoted as:

$$x_a(t) \longleftrightarrow X_a(j\Omega) \quad (3)$$

Note that Ω is real and denotes the continuous-time angular frequency variable in radians. In general, the CTFT is a complex function of Ω in the range $-\infty < \Omega < \infty$. It can be expressed in the polar form as:

$$X_a(j\Omega) = |X_a(j\Omega)| e^{j\theta_a(\Omega)} \quad (4)$$

where $\theta_a(\Omega) = \arg X_a(j\Omega)$. The quantity $|X_a(j\Omega)|$ is called the magnitude spectrum and the quantity $\theta_a(\Omega)$ is called the phase spectrum. Both spectra are real function of Ω and in general the CTFT $X_a(j\Omega)$ exists if $x_a(t)$ satisfies the Dirichlet conditions:

- the signal $x_a(t)$ has a finite number of discontinuities and a finite number of maxima and minima in any finite interval;
- the signal is absolutely integrable, i.e.:

$$\int_{-\infty}^{\infty} |x_a(t)| dt < \infty \quad (5)$$

If the Dirichlet conditions are satisfied, then:

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} X_a(j\Omega) e^{+j\Omega t} d\Omega \quad (6)$$

converges to $x_a(t)$ except at values of t where $x_a(t)$ has discontinuities. Moreover, it can be showed that if $x_a(t)$ is absolutely integrable, then proving the existence of the CTFT reduces to proving:

$$|X_a(j\Omega)| < \infty \quad (7)$$

0.1.1 Energy density spectrum

The total energy E_x of a finite energy continuous-time complex signal $x_a(t)$ is given by:

$$\begin{aligned} E_x &= \int_{-\infty}^{\infty} |x_a(t)|^2 dt \\ &= \int_{-\infty}^{\infty} x_a(t) x_a^*(t) dt \\ &= \int_{-\infty}^{\infty} x_a(t) \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} X_a^*(j\Omega) e^{-j\Omega t} d\Omega \right] dt \end{aligned} \quad (8)$$

Interchanging the order of the integration we get:

$$\begin{aligned} E_x &= \frac{1}{2\pi} \int_{-\infty}^{\infty} X_a^*(j\Omega) \left[\int_{-\infty}^{\infty} x_a(t) e^{-j\Omega t} dt \right] d\Omega \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} X_a^*(j\Omega) X_a(j\Omega) d\Omega \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} |X_a(j\Omega)|^2 d\Omega \end{aligned} \quad (9)$$

Hence:

$$\int_{-\infty}^{\infty} |x(t)|^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} |X_a(j\Omega)|^2 d\Omega \quad (10)$$

The above relation is more commonly known as the Parseval's relation for finite-energy continuous-time signals. The quantity $|X_a(j\Omega)|^2$ is called the energy density spectrum of $x_a(t)$ and it is usually denoted as:

$$S_{xx}(\Omega) = |X_a(j\Omega)|^2 \quad (11)$$

The energy over a specified range of frequencies $\Omega_a \leq \Omega \leq \Omega_b$ can be computed using:

$$E_{x,r} = \frac{1}{2\pi} \int_{\Omega_a}^{\Omega_b} S_{xx}(\Omega) d\Omega \quad (12)$$

0.1.2 Band-limited continuous-time signals

A full-band, finite-energy, continuous-time signal has a spectrum occupying the whole frequency range $-\infty \leq \Omega \leq \infty$. A band-limited continuous-time signal has a spectrum that is limited to a portion of the frequency range $-\infty \leq \Omega \leq \infty$. An ideal band-limited signal has a spectrum that is zero outside a finite frequency range $\Omega_a \leq |\Omega| \leq \Omega_b$ can be computed using:

$$X_a(j\Omega) = \begin{cases} 0 & 0 \leq |\Omega| < \Omega_a \\ 0 & \Omega_b < |\Omega| < \infty \end{cases} \quad (13)$$

However, an ideal band-limited signal cannot be generated in practice.

Band-limited signals are classified according to the frequency range where most of the signal's is concentrated:

- a lowpass, continuous-time signal has a spectrum occupying the frequency range $0 < |\Omega| \leq \Omega_p < \infty$, where Ω_p is called the bandwidth of the signal;
- a highpass, continuous-time signal has a spectrum occupying the frequency range $0 < \Omega_p \leq |\Omega| < \infty$, where the bandwidth of the signal is from Ω_p to ∞ ;
- a bandpass, continuous-time signal has a spectrum occupying the frequency range $0 < \Omega_L \leq |\Omega| \leq \Omega_H < \infty$, where $\Omega_H - \Omega_L$ is the bandwidth.

0.1.3 Discrete-time fourier transform

Let us introduce the definition of this concept.

Definition 3: Discrete-time Fourier transform

The discrete-time Fourier transform (DTFT) $X(e^{j\omega})$ of a sequence $x[n]$ is given by:

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n} \quad (14)$$

where in general $X(e^{j\omega})$ is a complex function of the real variable ω and can be written as:

$$X(e^{j\omega}) = X_{\text{re}}(e^{j\omega}) + jX_{\text{im}}(e^{j\omega}) \quad (15)$$

$X_{\text{re}}(e^{j\omega})$ and $X_{\text{im}}(e^{j\omega})$ are respectively, the real and imaginary parts of $X(e^{j\omega})$, and are real functions of ω . $X(e^{j\omega})$ can alternately be expressed as:

$$X(e^{j\omega}) = |X(e^{j\omega})|e^{j\theta(\omega)} \quad (16)$$

where $\theta(\omega) = \arg X(e^{j\omega})$. $|X(e^{j\omega})|$ and $\arg X(e^{j\omega})$ are called respectively magnitude function and phase function. Both quantities are again real functions of ω . In many applications, the DTFT is called the Fourier spectrum. Likewise, $|X(e^{j\omega})|$ and $\theta(\omega)$ are called respectively the magnitude and phase spectra.

For a real sequence $x[n]$, $|X(e^{j\omega})|$ and $X_{\text{re}}(e^{j\omega})$ are even functions of ω , whereas, $\theta(\omega)$ and $X_{\text{im}}(e^{j\omega})$ are odd functions of ω . Note also that $X(e^{j\omega}) = |X(e^{j\omega})|e^{j\theta(\omega+2\pi k)} = |X(e^{j\omega})|e^{j\theta(\omega)}$ for any integer k . The phase function $\theta(\omega)$ cannot be uniquely specified for any DTFT. Unless otherwise stated, we shall assume that the phase function $\theta(\omega)$ is restricted to the range of values $-\pi \leq \theta(\omega) < \pi$, called the principal value.

Example 1: DTFT of the unit sample sequence

The DTFT of the unit sample sequence $\delta[n]$ is given by:

$$\Delta(e^{j\omega}) = \sum_{n=-\infty}^{\infty} \delta[n]e^{-j\omega n} = \delta[0] = 1 \quad (17)$$

Example 2: DTFT of a causal sequence


Consider the causal sequence:

$$x[n] = \alpha^n \mu[n], \quad |\alpha| < 1 \quad (18)$$

Its DTFT is given by:

$$\begin{aligned} X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} \alpha^n \mu[n]e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n \\ &= \frac{1}{1 - \alpha e^{-j\omega}} \end{aligned} \quad (19)$$

as $|\alpha e^{-j\omega}| = |\alpha| < 1$. If we take for example $\alpha = 0.5$, we get the plot below for the magnitude and phase of the DTFT.



The DTFT $X(e^{j\omega})$ of a sequence $x[n]$ is a continuous function of ω . It is also a periodic function of ω with a period 2π :

$$X(e^{j(\omega+2\pi k)}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}e^{-j2\pi kn} = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n} = X(e^{j\omega}) \quad (20)$$

Therefore:

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n} \quad (21)$$

represents the Fourier series representation of the periodic function. As a result, the Fourier coefficients $x[n]$ can be computed from $X(e^{j\omega})$ using the Fourier integral:

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega})e^{j\omega n}d\omega \quad (22)$$

Proof. Consider:

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\sum_{\ell=-\infty}^{\infty} x[\ell]e^{-j\omega \ell} \right) e^{j\omega n} d\omega \quad (23)$$

The order of integration and summation can be interchanged if the summation inside the brackets converges uniformly, i.e. $X(e^{j\omega})$ exists. Then:

$$\begin{aligned} \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\sum_{\ell=-\infty}^{\infty} x[\ell]e^{-j\omega \ell} \right) e^{j\omega n} d\omega &= \sum_{\ell=-\infty}^{\infty} x[\ell] \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{j\omega(n-\ell)} d\omega \right) \\ &= \sum_{\ell=-\infty}^{\infty} x[\ell] \frac{\sin(\pi(n-\ell))}{\pi(n-\ell)} \\ &= \sum_{\ell=-\infty}^{\infty} x[\ell] \delta[n-\ell] \\ &= x[n] \end{aligned} \quad (24)$$

For the convergence condition, an infinite series of the form:

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n} \quad (25)$$

may or may not converg. Therefore, let us consider:

$$X_k(e^{j\omega}) = \sum_{n=-k}^k x[n]e^{-j\omega n} \quad (26)$$

Then for uniform convergence of $X_k(e^{j\omega})$:

$$\lim_{k \rightarrow \infty} X_k(e^{j\omega}) = X(e^{j\omega}) \quad (27)$$

Now, if $x[n]$ is an absolutely summable sequence, i.e., if $\sum_{n=-\infty}^{\infty} |x[n]| < \infty$, then:

$$|X(e^{j\omega})| = \left| \sum_{n=-k}^k x[n]e^{-j\omega n} \right| \leq \sum_{n=-k}^k |x[n]| < \infty \quad (28)$$

for all values of ω . Thus, the absolute summability of $x[n]$ is a sufficient condition for the existence of the DTFT $X(e^{j\omega})$. ■

Example 3: Absolute summability condition

The sequence $x[n] = \alpha^n \mu[n]$ for $|\alpha| < 1$ is absolutely summable as:

$$\sum_{n=-k}^k |\alpha^n \mu[n]| = \sum_{n=0}^{\infty} |\alpha^n| = \frac{1}{1 - |\alpha|} < \infty \quad (29)$$

and its DTFT $X(e^{j\omega})$ therefore converges to $\frac{1}{1 - \alpha e^{j\omega}}$ uniformly.

Note that since:

$$\sum_{n=-\infty}^{\infty} |x[n]|^2 \leq \left(\sum_{n=-\infty}^{\infty} |x[n]| \right)^2 \quad (30)$$

an absolutely summable sequence has always a finite energy. However, a finite-energy sequence is not necessarily absolutely summable.