

# Latent Graph Models

Wilker Aziz  
University of Amsterdam


May 14, 2018

# Outline

- 1 Recap
- 2 Discrete variables

# Variational auto-encoder

Generative model with NN likelihood



- complex (non-linear) observation model  $p_{\theta}(x|z)$
- complex (non-linear) mapping from data to latent variables  $q_{\lambda}(z|x)$

Jointly optimise generative model  $p_{\theta}(x|z)$  and inference model  $q_{\lambda}(z|x)$  under the same objective (ELBO)

$$\log p_{\theta}(x) \geq \overbrace{\mathbb{E}_{q_{\lambda}(z|x)} [\log p_{\theta}(x|Z)] - \text{KL}(q_{\lambda}(z|x) || p(z))}^{\text{ELBO}}$$

$$\log p_{\theta}(x) \geq \overbrace{\mathbb{E}_{q_{\lambda}(z|x)} [\log p_{\theta}(x|Z)] - \text{KL}(q_{\lambda}(z|x) || p(z))}^{\text{ELBO}}$$

Parameter estimation

$$\arg \max_{\theta, \lambda} \mathbb{E}_{q(\epsilon)} \left[ \log p_{\theta}(x | \underbrace{h^{-1}(\epsilon, \lambda)}_{=z}) \right] - \text{KL}(q_{\lambda}(z|x) || p(z))$$

$$\log p_{\theta}(x) \geq \overbrace{\mathbb{E}_{q_{\lambda}(z|x)} [\log p_{\theta}(x|Z)] - \text{KL}(q_{\lambda}(z|x) || p(z))}^{\text{ELBO}}$$

Parameter estimation

$$\arg \max_{\theta, \lambda} \mathbb{E}_{q(\epsilon)} \left[ \log p_{\theta}(x | \underbrace{h^{-1}(\epsilon, \lambda)}_{=z}) \right] - \text{KL}(q_{\lambda}(z|x) || p(z))$$

- assume  $\text{KL}(q_{\lambda}(z|x) || p(z))$  analytical  
true for exponential families

$$\log p_{\theta}(x) \geq \overbrace{\mathbb{E}_{q_{\lambda}(z|x)} [\log p_{\theta}(x|Z)] - \text{KL}(q_{\lambda}(z|x) || p(z))}^{\text{ELBO}}$$

Parameter estimation

$$\arg \max_{\theta, \lambda} \mathbb{E}_{q(\epsilon)} \left[ \log p_{\theta}(x | \underbrace{h^{-1}(\epsilon, \lambda)}_{=z}) \right] - \text{KL}(q_{\lambda}(z|x) || p(z))$$

- assume  $\text{KL}(q_{\lambda}(z|x) || p(z))$  analytical  
true for exponential families
- approximate  $\mathbb{E}_{q(\epsilon)} [\log p_{\theta}(x | h^{-1}(\epsilon, \lambda))]$  by sampling  
requires a reparameterisation  
 $h^{-1}(\epsilon, \lambda) \sim q_{\lambda}(z|x) \Leftrightarrow h(z, \lambda) \sim q(\epsilon)$

# Outline

- 1 Recap
- 2 Discrete variables



# Discrete variables

Suppose  $z$  is a  $d$ -dimensional binary vector  
i.e.  $z_i \in \{0, 1\}$

# Discrete variables

Suppose  $z$  is a  $d$ -dimensional binary vector  
i.e.  $z_i \in \{0, 1\}$

Then let's define an inference model

$$q_{\lambda}(z|x) = \underbrace{\prod_{i=1}^d q_{\lambda}(z_i|x)}_{\text{mean field}}$$

# Discrete variables

Suppose  $z$  is a  $d$ -dimensional binary vector  
i.e.  $z_i \in \{0, 1\}$

Then let's define an inference model

$$q_{\lambda}(z|x) = \underbrace{\prod_{i=1}^d q_{\lambda}(z_i|x)}_{\text{mean field}} = \prod_{i=1}^d \text{Bern}(z_i | \underbrace{\text{sigmoid}(f_{\lambda}(x))}_{\text{NN}}) \quad (1)$$

# Discrete variables

Suppose  $z$  is a  $d$ -dimensional binary vector  
i.e.  $z_i \in \{0, 1\}$

Then let's define an inference model

$$q_{\lambda}(z|x) = \underbrace{\prod_{i=1}^d q_{\lambda}(z_i|x)}_{\text{mean field}} = \prod_{i=1}^d \text{Bern}(z_i | \underbrace{\text{sigmoid}(f_{\lambda}(x))}_{\text{NN}}) \quad (1)$$

Can we reparameterise  $q_{\lambda}(z|x)$ ?

# Bernoulli pmf

$$\text{Bern}(z_i | b_i) = b_i^{z_i} (1 - b_i)^{1-z_i}$$

# Bernoulli pmf

$$\begin{aligned}\text{Bern}(z_i|b_i) &= b_i^{z_i}(1-b_i)^{1-z_i} \\ &= \begin{cases} b_i & \text{if } z_i = 1 \\ 1-b_i & \text{if } z_i = 0 \end{cases} \end{aligned} \quad (2)$$

# Bernoulli pmf

$$\begin{aligned}\text{Bern}(z_i|b_i) &= b_i^{z_i}(1-b_i)^{1-z_i} \\ &= \begin{cases} b_i & \text{if } z_i = 1 \\ 1-b_i & \text{if } z_i = 0 \end{cases} \end{aligned} \quad (2)$$

Can we reparameterise a Bernoulli variable?

# Reparameterisation requires a Jacobian matrix

Not really :(

$$q(z; \lambda) = \underbrace{\phi(\epsilon = h(z, \lambda))}_{\text{change of density}} |\det J_{h(z, \lambda)}| \quad (3)$$

Elements in the Jacobian matrix

$$J_{h(z, \lambda)}[i, j] = \frac{\partial h_i(z, \lambda)}{\partial z_j}$$

are not defined for non-differentiable functions



# Relaxation

Let's redefine  $z_i$  to live in the interval  $(0, 1)$

- and find an alternative reparameterisable **density**

Examples

- $Z \sim \mathcal{LN}(u, s^2)$   
 $z = \text{sigmoid}(u + s\epsilon)$  with  $\epsilon \sim \mathcal{N}(0, 1)$
- $Z \sim \text{Concrete}(u, \tau)$   
 $z = \text{sigmoid}(\frac{u+\epsilon}{\tau})$  with  $\epsilon \sim \text{Gumbel}(0, 1)$

# Relaxation

Let's redefine  $z_i$  to live in the interval  $(0, 1)$

- and find an alternative reparameterisable **density**

Examples

- $Z \sim \mathcal{LN}(u, s^2)$   
 $z = \text{sigmoid}(u + s\epsilon)$  with  $\epsilon \sim \mathcal{N}(0, 1)$
- $Z \sim \text{Concrete}(u, \tau)$   
 $z = \text{sigmoid}(\frac{u+\epsilon}{\tau})$  with  $\epsilon \sim \text{Gumbel}(0, 1)$

But note that we **no longer** have a discrete variable

# Concrete or Gumbel-Softmax

An alternative parameterisation of a Categorical variable

$$A \sim \text{Cat}(\text{softmax}(\phi))$$

$$A = \arg \max_i [\phi_i + \epsilon_i]_{i=1}^K \quad \text{where } \epsilon \sim \text{Gumbel}(0, I) \quad (4)$$

# Concrete or Gumbel-Softmax

An alternative parameterisation of a Categorical variable

$$A \sim \text{Cat}(\text{softmax}(\phi))$$

$$A = \arg \max_i [\phi_i + \epsilon_i]_{i=1}^K \quad \text{where } \epsilon \sim \text{Gumbel}(0, 1) \quad (4)$$

We can sample a one-hot encoding of the categorical variable

$$B = \text{onehot} \left( \arg \max_i [\phi_i + \epsilon_i]_{i=1}^K \right) \quad (5)$$

# Concrete or Gumbel-Softmax

An alternative parameterisation of a Categorical variable

$$A \sim \text{Cat}(\text{softmax}(\phi))$$

$$A = \arg \max_i [\phi_i + \epsilon_i]_{i=1}^K \quad \text{where } \epsilon \sim \text{Gumbel}(0, 1) \quad (4)$$

We can sample a one-hot encoding of the categorical variable

$$B = \text{onehot} \left( \arg \max_i [\phi_i + \epsilon_i]_{i=1}^K \right) \quad (5)$$

And we get a continuous relaxation with softmax

$$B = \text{softmax}(\phi + \epsilon) \quad (6)$$

# Concrete or Gumbel-Softmax

An alternative parameterisation of a Categorical variable

$$A \sim \text{Cat}(\text{softmax}(\phi))$$

$$A = \arg \max_i [\phi_i + \epsilon_i]_{i=1}^K \quad \text{where } \epsilon \sim \text{Gumbel}(0, 1) \quad (4)$$

We can sample a one-hot encoding of the categorical variable

$$B = \text{onehot} \left( \arg \max_i [\phi_i + \epsilon_i]_{i=1}^K \right) \quad (5)$$

And we get a continuous relaxation with softmax


$$B = \text{softmax}(\phi + \epsilon) \quad (6)$$

Finally, with a temperature  $\tau$  we can approach a one-hot encoding of the most likely category as  $\tau \rightarrow 0$

$$B = \text{softmax} \left( \frac{\phi + \epsilon}{\tau} \right) \quad (7)$$

# Simplex

The tips of the simplex represent a one-hot encoding of a 3-way Categorical variable




- the softmax relaxes the variable to take on values in the interior of the simplex
- as we cool down the system we push most of the mass towards the tips

---

Illustrations from (Maddison et al., 2016).

# Concrete samples



Illustrations from (Jang et al., 2016).



# Summary

## Deep learning in NLP

- task-driven feature extraction
- models with more realistic assumptions

## Probabilistic modelling

- better (or at least more explicit) statistical assumptions
- compact models
- semi-supervised learning

# Literature I

- Eric Jang, Shixiang Gu, and Ben Poole. Categorical reparameterization with gumbel-softmax. *arXiv preprint arXiv:1611.01144*, 2016.
- Diederik P. Kingma and Max Welling. Auto-Encoding Variational Bayes. 2013. URL <http://arxiv.org/abs/1312.6114>.
- Chris J Maddison, Andriy Mnih, and Yee Whye Teh. The concrete distribution: A continuous relaxation of discrete random variables. *arXiv preprint arXiv:1611.00712*, 2016.