

Probability and Statistics: Lecture-23

Monsoon-2020

by Pawan Kumar (IIIT, Hyderabad)

on October 5, 2020

» Checklist for online class

1. Turn off your microphone, when you are listening
2. Turn on microphone only when you have question
3. Attend tutorials to practice problems or to discuss solutions or doubts
4. Chat is not always reliable, I may not look at chat

» Table of contents

1. Continuous Distributions

2. Mixed Random Variable

» Special Distribution: Exponential Distributions...

» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution

Let X be a continuous random variable. Here X is said to have **exponential distribution** with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its **PDF** is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$


» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution

Let X be a continuous random variable. Here X is said to have exponential distribution with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its PDF is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

$$\int_0^{\infty} f_X(x) dx = 1$$



» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution


Let X be a continuous random variable. Here X is said to have **exponential distribution** with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its **PDF** is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

$$F_X(x) = \int_{-\infty}^x f_X(u) du$$

The **CDF** is given by

$$F_X(x) = \int_0^x \lambda e^{-\lambda t} dt = 1 - e^{-\lambda x}$$



» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution


Let X be a continuous random variable. Here X is said to have **exponential distribution** with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its **PDF** is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

The **CDF** is given by

$$F_X(x) = \int_0^x \lambda e^{-\lambda t} dt = 1 - e^{-\lambda x}$$

The **expectation** is




» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution

Let X be a continuous random variable. Here X is said to have **exponential distribution** with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its **PDF** is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$



The **CDF** is given by

$$F_X(x) = \int_0^x \lambda e^{-\lambda t} dt = 1 - e^{-\lambda x}$$

Handwritten orange notes: A bracket under the integral term $\lambda e^{-\lambda t}$ is labeled $f_X(t)$. A bracket under the result $1 - e^{-\lambda x}$ is labeled $F_X(x)$.

The **expectation** is

$$\begin{aligned} E[X] &= \int_0^\infty x \lambda e^{-\lambda x} dx = \frac{1}{\lambda} \int_0^\infty y e^{-y} dy \\ &= \frac{1}{\lambda} [-e^{-y} - y e^{-y}]_0^\infty = \frac{1}{\lambda} \end{aligned}$$


Handwritten orange notes: The term $x \lambda e^{-\lambda x}$ is circled and labeled $f_X(x)$. The final result $\frac{1}{\lambda}$ is circled.

» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution

Let X be a continuous random variable. Here X is said to have **exponential distribution** with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its **PDF** is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$




» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution

Let X be a continuous random variable. Here X is said to have **exponential distribution** with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its **PDF** is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$



Var(X) is given by:

$E[X^2] - (E[X])^2$ $\lambda x = y$

$$E[X^2] = \int_0^{\infty} x^2 \lambda e^{-\lambda x} dx = \frac{1}{\lambda^2} \int_0^{\infty} y^2 e^{-y} dy$$

int. by parts

$$= \frac{1}{\lambda^2} \left[-2e^{-y} - 2ye^{-y} - y^2 e^{-y} \right]_0^{\infty} = \frac{2}{\lambda^2}$$


verify

» Special Distribution: Exponential Distributions...

Definition of Exponential Distribution

Let X be a continuous random variable. Here X is said to have **exponential distribution** with parameter $\lambda > 0$ shown as $X \sim \text{Exponential}(\lambda)$, if its **PDF** is given as follows

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$




$\text{Var}(X)$ is given by:

$$\begin{aligned} E[X^2] &= \int_0^{\infty} x^2 \lambda e^{-\lambda x} dx = \frac{1}{\lambda^2} \int_0^{\infty} y^2 e^{-y} dy \\ &= \frac{1}{\lambda^2} [-2e^{-y} - 2ye^{-y} - y^2 e^{-y}]_0^{\infty} = \frac{2}{\lambda^2} \end{aligned}$$

$$\text{Var}(X) = \underbrace{E[X^2]} - (\underbrace{E[X]})^2 = \frac{2}{\lambda^2} - \frac{1}{\lambda^2} = \frac{1}{\lambda^2}$$

» Exponential Distribution is Memoryless...



» Exponential Distribution is Memoryless...

Exponential distribution is memoryless

If X is **exponential** random variable with parameter $\lambda > 0$, then X is a **memoryless** variable:

» Exponential Distribution is Memoryless...



Exponential distribution is memoryless

If X is **exponential** random variable with parameter $\lambda > 0$, then X is a **memoryless** variable:

$$P(X > x + a \mid X > a) = \underbrace{P(X > x)}, \quad \text{for } a, x \geq 0.$$

» Exponential Distribution is Memoryless...

Exponential distribution is memoryless

If X is exponential random variable with parameter $\lambda > 0$, then X is a memoryless variable:

$$P(X > x + a \mid X > a) = P(X > x), \quad \text{for } a, x \geq 0.$$

$$\begin{aligned} \underline{P(X > x + a \mid X > a)} &= \frac{P(X > x + a, X > a)}{P(X > a)} \\ &= \frac{P(X > x + a)}{P(X > a)} = \frac{1 - F_X(x + a)}{1 - F_X(a)} \\ &= \frac{e^{-\lambda(x+a)}}{e^{-\lambda a}} = e^{-\lambda x} \\ &= \underline{P(X > x)} \end{aligned}$$

$$\begin{aligned} F_X(x) &= \int_{-\infty}^x \lambda e^{-\lambda t} dt \\ P(X \leq x) \\ \Rightarrow P(X > x + a) \\ &= 1 - P(X \leq x + a) \\ &= 1 - F_X(x + a) \\ &= 1 - e^{-\lambda(x+a)} \\ F_X(x) &= 1 - e^{-\lambda x} \\ e^{-\lambda x} &= 1 - F_X(x) \\ &= P(X > x) \end{aligned}$$

» Normal (Gaussian) Distribution...

» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a standard normal (standard Gaussian) random variable,

» Normal (Gaussian) Distribution...

Normal

$$Z \sim N(0, 1)$$

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal (standard Gaussian) random variable**, shown as $Z \sim (0, 1)$,

» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal (standard Gaussian) random variable**, shown as $Z \sim (0, 1)$, if its **PDF** is given by

» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal** (standard Gaussian) **random variable**, shown as $Z \sim (0, 1)$, if its **PDF** is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}, \quad \text{for all } z \in \mathbb{R}$$

Handwritten notes:

- $\int_{-\infty}^{\infty} f_Z(z) dz = 1$ check
- $\int_{-\infty}^{\infty} e^{-z^2/2} dz = \sqrt{2\pi}$
- Indefinite integral

» Normal (Gaussian) Distribution...


Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal** (standard Gaussian) **random variable**, shown as $Z \sim (0, 1)$, if its **PDF** is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}, \quad \text{for all } z \in \mathbb{R}$$

← even or odd

$$\underline{\underline{f(-z) = f(z)}}$$




» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a standard normal (standard Gaussian) random variable, shown as $Z \sim (0, 1)$, if its PDF is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}, \quad \text{for all } z \in \mathbb{R}$$




* Most important Probability Distribution!

» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal (standard Gaussian) random variable**, shown as $Z \sim (0, 1)$, if its **PDF** is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}, \quad \text{for all } z \in \mathbb{R}$$




- * Most important Probability Distribution!
- * **Central Limit Theorem** (TODO):

» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal (standard Gaussian) random variable**, shown as $Z \sim (0, 1)$, if its **PDF** is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}, \quad \text{for all } z \in \mathbb{R}$$




- * Most important Probability Distribution!
- * **Central Limit Theorem** (TODO):
 - * If we add **large** number of random variable, then the distribution of the **sum is normal** (proof later)

» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal (standard Gaussian) random variable**, shown as $Z \sim (0, 1)$, if its **PDF** is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}, \quad \text{for all } z \in \mathbb{R}$$



- * Most important Probability Distribution!
- * **Central Limit Theorem** (TODO):
 - * If we add **large** number of random variable, then the distribution of the **sum is normal** (proof later)


» Normal (Gaussian) Distribution...

Definition of Standard Normal Random Variable

A continuous random variable Z is said to be a **standard normal (standard Gaussian) random variable**, shown as $Z \sim (0, 1)$, if its **PDF** is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}, \quad \text{for all } z \in \mathbb{R}$$

$$\int_{-\infty}^{\infty} e^{-z^2/2} dz = \sqrt{2\pi}$$



- * Most important Probability Distribution!
- * **Central Limit Theorem** (TODO):
 - * If we add **large** number of random variable, then the distribution of the **sum is normal** (proof later)
- * Here $\frac{1}{\sqrt{2\pi}}$ is there to make area under curve 1

» Mean and Variance of Standard Normal Distribution...

» Mean and Variance of Standard Normal Distribution...

Mean and Variance of Standard Normal Distribution

Let Z be a **normal distribution**, i.e., $Z \sim N(0, 1)$, then $E[Z] = 0$ and $\text{Var}(Z) = 1$.

Recall

If $g(u) : \mathbb{R} \rightarrow \mathbb{R}$. If $g(u)$ is an **odd function**, i.e., $g(-u) = -g(u)$, and

$$\left| \int_0^{\infty} g(u) du \right| < \infty,$$

then

$$\int_{-\infty}^{\infty} g(u) du = 0.$$

» Mean and Variance of Standard Normal Distribution...

Mean and Variance of Standard Normal Distribution

Let Z be a normal distribution, i.e., $Z \sim N(0, 1)$, then $E[Z] = 0$ and $\text{Var}(Z) = 1$.

Recall

If $g(u) : \mathbb{R} \rightarrow \mathbb{R}$. If $g(u)$ is an **odd function**, i.e., $g(-u) = -g(u)$, and

$$\left| \int_0^{\infty} g(u) du \right| < \infty,$$

then

$$\int_{-\infty}^{\infty} g(u) du = 0.$$

» Answer to previous problem...

$$z \sim N(0,1)$$

$$E[z] = \int_{-\infty}^{\infty} x \cdot \underbrace{\frac{1}{\sqrt{2\pi}} e^{-x^2/2}}_{\text{odd fn.}} dx$$

= 0

$$E[z^2] = \int_{-\infty}^{\infty} x^2 \underbrace{\frac{1}{\sqrt{2\pi}} e^{-x^2/2}}_{\text{even fn.}} dx$$

Int. by parts

$$\int_{-\infty}^{\infty} x^2 e^{-x^2/2} dx = \sqrt{2\pi}$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x^2 e^{-x^2/2} dx$$

$$= \frac{1}{\sqrt{2\pi}} 2 \int_0^{\infty} x^2 e^{-x^2/2} dx$$

$$I = \int_0^{\infty} \underbrace{x^2}_{1^{st}} \underbrace{e^{-x^2/2}}_{2^{nd}} dx$$

$$= \left[x^2 \int_0^{\infty} e^{-x^2/2} dx \right]_0^{\infty} - \int_0^{\infty} 2x \int_0^{\infty} e^{-x^2/2} dx$$

» Answer to previous problem...

$$= 2 \int_0^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx =$$

$$= 2 \int_0^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$$

$$= \frac{2}{\sqrt{2\pi}} \int_0^{\infty} e^{-x^2/2} dx$$

$$= \frac{2}{\sqrt{2\pi}} \cdot \frac{1}{2} \int_{-\infty}^{\infty} e^{-x^2/2} dx = \frac{1}{\sqrt{2\pi}}$$

» CDF of Standard Normal Distribution...

» CDF of Standard Normal Distribution...

Definition of CDF of Standard Normal Distribution

The CDF of the **standard normal distribution** is denoted by Φ

» CDF of Standard Normal Distribution...

Definition of CDF of Standard Normal Distribution

The CDF of the **standard normal distribution** is denoted by Φ

$$\Phi(x) = \underbrace{P(Z \leq x)} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{u^2}{2}} du$$

» CDF of Standard Normal Distribution...

Definition of CDF of Standard Normal Distribution

The CDF of the **standard normal distribution** is denoted by Φ

$$\Phi(x) = P(Z \leq x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{u^2}{2}} du$$

* The integral **does not** have a **closed** form solution!

» CDF of Standard Normal Distribution...

Definition of CDF of Standard Normal Distribution

The CDF of the **standard normal distribution** is denoted by Φ

$Z \sim N(0,1)$

$$\Phi(x) = P(Z \leq x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{u^2}{2}} du$$

Handwritten notes: Arrows point from $\Phi(x)$ to $F(x)$ and from $P(Z \leq x)$ to $F(x)$. The term $e^{-\frac{u^2}{2}}$ is circled.

- * The integral **does not** have a **closed** form solution!
- * However, values of $F(Z)$ have been tabulated

» CDF of Standard Normal Distribution...

$$F_Z(x) = \underline{P(Z \leq x)}$$

Definition of CDF of Standard Normal Distribution


The CDF of the standard normal distribution is denoted by Φ

$$\Phi(x) = \underline{P(Z \leq x)} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{u^2}{2}} du$$

- * The integral **does not** have a closed form solution!
- * However, values of $F(Z)$ have been tabulated
- * The CDF of any normal distribution can be written in terms of Φ function


» CDF of Standard Normal Distribution...

» CDF of Standard Normal Distribution...



The Φ function satisfies the following properties:


» CDF of Standard Normal Distribution...



The Φ function satisfies the following properties:

$$\checkmark \lim_{x \rightarrow \infty} \Phi(x) = 1, \quad \lim_{x \rightarrow -\infty} \Phi(x) = 0$$


» CDF of Standard Normal Distribution...



The Φ function satisfies the following properties:

- * $\lim_{x \rightarrow \infty} \Phi(x) = 1$, $\lim_{x \rightarrow -\infty} \Phi(x) = 0$
- * $\Phi(0) = \frac{1}{2}$

» CDF of Standard Normal Distribution...



The Φ function satisfies the following properties:

- * $\lim_{x \rightarrow \infty} \Phi(x) = 1, \quad \lim_{x \rightarrow -\infty} \Phi(x) = 0$

- * $\Phi(0) = \frac{1}{2}$

- * $\Phi(-x) = 1 - \Phi(x)$ for all $x \in \mathbb{R}$

» Bound for Φ Function...

» Bound for Φ Function...

Bound for Φ Function

Let $Z \sim N(0, 1)$. We recall that

$$\Phi(x) = P(Z \leq x).$$

$$\begin{aligned} 1 - \Phi(x) &= 1 - P(Z \leq x) \\ &= P(Z > x). \end{aligned}$$

For all $x \geq 0$, the Φ -function satisfies the following bound

$$\frac{1}{\sqrt{2\pi}} \frac{x}{x^2 + 1} e^{-x^2/2} \leq \underbrace{1 - \Phi(x)}_{P(Z > x)} \leq \frac{1}{\sqrt{2\pi}} \frac{1}{x} e^{-x^2/2}$$

» Answer to previous problem...

To show upper bound:

$$P(z > x) = \frac{1}{\sqrt{2\pi}} \int_x^\infty \underbrace{e^{-u^2/2}}_{g(u)} du$$

$$\leq \frac{1}{\sqrt{2\pi}} \int_x^\infty \underbrace{\frac{u}{x}}_{g(u)} e^{-u^2/2} du$$


$$= \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{x} \int_x^\infty u e^{-u^2/2} du$$

Note: obviously $u \geq x \geq 0$

Recall

$$f(x) \leq g(x) \quad x \in [a, b]$$

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx$$



$$= \frac{1}{\sqrt{2\pi}} \frac{1}{x} \left[-e^{-u^2/2} \right]_x^\infty = \frac{1}{\sqrt{2\pi}} \frac{1}{x} e^{-x^2/2}$$