

# Probability and Statistics: Lecture-1

Monsoon-2020

by Pawan Kumar (IIIT, Hyderabad)

on August 10, 2020

## » Table of contents

### 1. Basic Counting

### 2. Sum Rule

### 3. Set Theory

### 4. Product Rule

### 5. Tuples

## » Why Counting?

## » Why Counting?

- \* counting is a basic task in mathematics

## » Why Counting?

- \* counting is a basic task in mathematics
- \* **Our objective:** To tell how many objects are there without actually counting

## » Why Counting?

- \* counting is a basic task in mathematics
- \* **Our objective:** To tell how many objects are there without actually counting
- \* **applications of counting:**

## » Why Counting?

- \* counting is a basic task in mathematics
- \* **Our objective:** To tell how many objects are there without actually counting
- \* **applications of counting:**
  - \* number of steps of an algorithm

## » Why Counting?

- \* counting is a basic task in mathematics
- \* **Our objective:** To tell how many objects are there without actually counting
- \* **applications of counting:**
  - \* number of steps of an algorithm
  - \* estimating probability of occurrence of an event (**in this course**)



## » Why Counting?

- \* counting is a basic task in mathematics
- \* **Our objective:** To tell how many objects are there without actually counting
- \* **applications of counting:**
  - \* number of steps of an algorithm
  - \* estimating probability of occurrence of an event (**in this course**)
  - \* proofs such as Pigeon Hole Principle (PHP)



## » Brief History of Counting



## » Brief History of Counting



- \* **Ishango Bone** is possibly the **oldest** mathematical artefact still in existence

## » Brief History of Counting



- \* **Ishango Bone** is possibly the **oldest** mathematical artefact still in existence
- \* It dates back to the Upper Paleolithic period of human history, and is approximately **20,000 years old**.

## » Brief History of Counting



- \* **Ishango Bone** is possibly the **oldest** mathematical artefact still in existence
- \* It dates back to the Upper Paleolithic period of human history, and is approximately **20,000 years old**.
- \* The bone is 10 cm long and contains a series of notches, which many scientists believe were used for counting.

## » Brief History of Counting



## » Brief History of Counting



- \* This tablet shows a **multiplication table** that was created around **2600 BCE** in the Sumerian city of Shuruppak



## » Brief History of Counting



- \* This tablet shows a **multiplication table** that was created around **2600 BCE** in the Sumerian city of Shuruppak
- \* The table has three columns. The dots in the first two columns represent distances ranging from around 6 meters to 3 kilometres. The third column contains the product of the first two

## » Brief History of Counting



- \* This tablet shows a **multiplication table** that was created around **2600 BCE** in the Sumerian city of Shuruppak
- \* The table has three columns. The dots in the first two columns represent distances ranging from around 6 meters to 3 kilometres. The third column contains the product of the first two
- \* Sumer was a region of ancient Mesopotamia in the Middle East

## » How do we count?

## » How do we count?

### Question-2

How many numbers are there between 21 and 31?

## » How do we count?

### Question-2

How many numbers are there between 21 and 31?

### Answer

11

## » How do we count?

### Question-2

How many numbers are there between 21 and 31?

### Answer

11

### Question-3

How many numbers are there between  $m$  and  $n$ ,  $m < n$ ?

## » How do we count?

### Question-2

How many numbers are there between 21 and 31?

### Answer

11

### Question-3

How many numbers are there between  $m$  and  $n$ ,  $m < n$ ?

### Answer

$m - n + 1$

## » How do we count?

### Question-1

How many numbers between 33 and 67 are divisible by 4?



## » Sum Rule: Statement and Example

## » Sum Rule: Statement and Example

### Sum Rule

If there are  $m$  objects of first type and there are  $n$  objects of second type, then there are  $n + m$  objects of one of the two types.

## » Sum Rule: Statement and Example

### Sum Rule

If there are  $m$  objects of first type and there are  $n$  objects of second type, then there are  $n + m$  objects of one of the two types.

### Sum Rule Example

How many of the Pizza or Burgers places are there?

Pizza places



Burger places



## » Sum Rule: Statement and Example

### Sum Rule

If there are  $m$  objects of first type and there are  $n$  objects of second type, then there are  $n + m$  objects of one of the two types.

### Sum Rule Example

How many of the Pizza or Burgers places are there?



### Answer

There are 7 Pizzas and there are 5 Burgers, hence, by sum rule, we have  $7 + 5 = 12$

## » Another Sum Rule Example

## » Another Sum Rule Example


### Example

A piece starts in the bottom left corner of a chessboard. In one move it can move one step to the right or one step up. How many moves are needed to get to the position on the picture?

## » Another Sum Rule Example

### Example

A piece stays in the bottom left corner of a chessboard. In one move it can move one step to the right or one step up. How many moves are needed to get to the position on the picture?




## » Another Sum Rule Example

### Example

A piece stays in the bottom left corner of a chessboard. In one move it can move one step to the right or one step up. How many moves are needed to get to the position on the picture?

- \* There are **two** types of moves:






## » Another Sum Rule Example

### Example

A piece stays in the bottom left corner of a chessboard. In one move it can move one step to the right or one step up. How many moves are needed to get to the position on the picture?




- \* There are **two** types of moves: move right and move up

## » Another Sum Rule Example

### Example

A piece stays in the bottom left corner of a chessboard. In one move it can move one step to the right or one step up. How many moves are needed to get to the position on the picture?




- \* There are **two** types of moves: move right and move up
- \* To get to the column 4, we need 3 moves to the right

## » Another Sum Rule Example

### Example

A piece stays in the bottom left corner of a chessboard. In one move it can move one step to the right or one step up. How many moves are needed to get to the position on the picture?




- \* There are **two** types of moves: move right and move up
- \* To get to the column 4, we need 3 moves to the right
- \* To get to the row 6, we need 5 moves up

## » Another Sum Rule Example

### Example

A piece stays in the bottom left corner of a chessboard. In one move it can move one step to the right or one step up. How many moves are needed to get to the position on the picture?



- \* There are **two** types of moves: move right and move up
- \* To get to the column 4, we need 3 moves to the right
- \* To get to the row 6, we need 5 moves up
- \* **Applying sum rule:** In total, we need  $3+5=8$  moves

» **Caution using Sum Rule!**

## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

### Answer

\* There are 5 numbers divisible by 2: 2, 4, 6, 8, 10

## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

### Answer

- \* There are 5 numbers divisible by 2: 2, 4, 6, 8, 10
- \* There are 3 numbers divisible by 3: 3, 6, 9



## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

### Answer

- \* There are 5 numbers divisible by 2: 2, 4, 6, 8, 10
- \* There are 3 numbers divisible by 3: 3, 6, 9
- \* Hence, by sum rule, the answer is:  $5+3 = 8$

## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

### Answer

- \* There are 5 numbers divisible by 2: 2, 4, 6, 8, 10
- \* There are 3 numbers divisible by 3: 3, 6, 9
- \* Hence, by sum rule, the answer is:  $5+3 = 8$

Is this correct answer?

## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

### Answer

- \* There are 5 numbers divisible by 2: 2, 4, 6, 8, 10
- \* There are 3 numbers divisible by 3: 3, 6, 9
- \* Hence, by sum rule, the answer is:  $5+3 = 8$

Is this correct answer? **No**

## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

### Answer

- \* There are 5 numbers divisible by 2: 2, 4, 6, 8, 10
- \* There are 3 numbers divisible by 3: 3, 6, 9
- \* Hence, by sum rule, the answer is:  $5+3 = 8$

Is this correct answer? **No**

Let us count directly: 2,3,4,6,8,9,10, that is the answer is 7

## » Caution using Sum Rule!

### Question

Count all integers from 1 to 10 that are divisible by 2 or by 3

### Answer

- \* There are 5 numbers divisible by 2: 2, 4, 6, 8, 10
- \* There are 3 numbers divisible by 3: 3, 6, 9
- \* Hence, by sum rule, the answer is:  $5+3 = 8$

Is this correct answer? **No**

Let us count directly: 2,3,4,6,8,9,10, that is the answer is 7

### Caution with Sum Rule

In the rule of sum, **no object should belong to both types!**

## » Review of Set theory

## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

### Remarks and Examples

- \* ordering of elements in set **does not** matter



## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

### Remarks and Examples

- \* ordering of elements in set **does not** matter
- \* set of **natural** numbers  $\mathbb{N} = \{1, 2, 3, \dots\}$

## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

### Remarks and Examples

- \* ordering of elements in set **does not** matter
- \* set of **natural** numbers  $\mathbb{N} = \{1, 2, 3, \dots\}$
- \* Set of **integers**  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3\}$

## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

### Remarks and Examples

- \* ordering of elements in set **does not** matter
- \* set of **natural** numbers  $\mathbb{N} = \{1, 2, 3, \dots\}$
- \* Set of **integers**  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3\}$
- \* Set of **rational** numbers  $\mathbb{Q}$ , set of **real** numbers  $\mathbb{R}$

## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

### Remarks and Examples

- \* ordering of elements in set **does not** matter
- \* set of **natural** numbers  $\mathbb{N} = \{1, 2, 3, \dots\}$
- \* Set of **integers**  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3\}$
- \* Set of **rational** numbers  $\mathbb{Q}$ , set of **real** numbers  $\mathbb{R}$
- \*  $[2, 3] = \{x : 2 \leq x \leq 3\}$ , a **closed** interval on real line

## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

### Remarks and Examples

- \* ordering of elements in set **does not** matter
- \* set of **natural** numbers  $\mathbb{N} = \{1, 2, 3, \dots\}$
- \* Set of **integers**  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3\}$
- \* Set of **rational** numbers  $\mathbb{Q}$ , set of **real** numbers  $\mathbb{R}$
- \*  $[2, 3] = \{x : 2 \leq x \leq 3\}$ , a **closed** interval on real line
- \*  $(-1, 2) = \{x : -1 < x < 2\}$ , an **open** interval on real line

## » Review of Set theory

### Definition of a Set

A set is a collection of things (called elements).

### Remarks and Examples

- \* ordering of elements in set **does not** matter
- \* set of **natural** numbers  $\mathbb{N} = \{1, 2, 3, \dots\}$
- \* Set of **integers**  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3\}$
- \* Set of **rational** numbers  $\mathbb{Q}$ , set of **real** numbers  $\mathbb{R}$
- \*  $[2, 3] = \{x : 2 \leq x \leq 3\}$ , a **closed** interval on real line
- \*  $(-1, 2) = \{x : -1 < x < 2\}$ , an **open** interval on real line
- \*  $(1, 3] = \{x : 1 < x \leq 3\}$ , a **half open/closed** interval on real line



## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$



## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$

## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$
- \* A set with no elements is called **empty set or null set**, denoted by  $\phi$

## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$
- \* A set with no elements is called **empty set or null set**, denoted by  $\phi$
- \* The **universal set** is the set of all things that we could possibly consider in the context we are studying

## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$
- \* A set with no elements is called **empty set or null set**, denoted by  $\phi$
- \* The **universal set** is the set of all things that we could possibly consider in the context we are studying
  - \* every set  $A$  is a **subset** of the **universal** set

## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$
- \* A set with no elements is called **empty set or null set**, denoted by  $\phi$
- \* The **universal set** is the set of all things that we could possibly consider in the context we are studying
  - \* every set  $A$  is a **subset** of the **universal set**
- \* A set  $S$  is called **countable**, if there exists a **bijective** function

$$f: S \rightarrow \mathbb{N}$$

## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$
- \* A set with no elements is called **empty set or null set**, denoted by  $\phi$
- \* The **universal set** is the set of all things that we could possibly consider in the context we are studying
  - \* every set  $A$  is a **subset** of the **universal** set
- \* A set  $S$  is called **countable**, if there exists a **bijective** function

$$f: S \rightarrow \mathbb{N}$$

- \* The sets  $\mathbb{N}$ ,  $\mathbb{Q}$  are **countable**

## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$
- \* A set with no elements is called **empty set or null set**, denoted by  $\phi$
- \* The **universal set** is the set of all things that we could possibly consider in the context we are studying
  - \* every set  $A$  is a **subset** of the **universal** set
- \* A set  $S$  is called **countable**, if there exists a **bijective** function

$$f: S \rightarrow \mathbb{N}$$

- \* The sets  $\mathbb{N}, \mathbb{Q}$  are **countable**
- \* A set with finite number of elements is called **finite** set

## » Review of Set Theory Continued

- \* A set  $A$  is called **subset** of a set  $B$  if every element of  $A$  is also an element of  $B$ . It is denoted by  $A \subseteq B$
- \* Two sets  $A$  and  $B$  are said to be **equal** if  $A \subseteq B$  and  $B \subseteq A$
- \* A set with no elements is called **empty set or null set**, denoted by  $\phi$
- \* The **universal set** is the set of all things that we could possibly consider in the context we are studying
  - \* every set  $A$  is a **subset** of the **universal** set
- \* A set  $S$  is called **countable**, if there exists a **bijective** function

$$f: S \rightarrow \mathbb{N}$$

- \* The sets  $\mathbb{N}, \mathbb{Q}$  are **countable**
- \* A set with finite number of elements is called **finite** set
- \* A set which is countable and not finite is called **countably infinite**






## » Set Theory: Venn Diagram

- \* Venn diagrams are useful in analysing the relationship between sets


## » Set Theory: Venn Diagram

- \* Venn diagrams are useful in analysing the relationship between sets




## » Set Theory: Venn Diagram

- \* Venn diagrams are useful in analysing the relationship between sets



- \* Venn diagram showing subset relationship



## » Set Operations: Union

## » Set Operations: Union


- \* The union of two sets is a set containing all elements that are in  $A$  or in  $B$ . Here  $A$  union  $B$  is denoted by  $A \cup B$

## » Set Operations: Union

- \* The union of two sets is a set containing all elements that are in  $A$  or in  $B$ . Here  $A$  union  $B$  is denoted by  $A \cup B$ 
  - \* Example:  $\{1, 2\} \cup \{2, 3\} = \{1, 2, 3\}$

## » Set Operations: Union


- \* The union of two sets is a set containing all elements that are in  $A$  or in  $B$ . Here  $A$  union  $B$  is denoted by  $A \cup B$ 
  - \* Example:  $\{1, 2\} \cup \{2, 3\} = \{1, 2, 3\}$





## » Set Operations: Union

- \* The union of two sets is a set containing all elements that are in  $A$  or in  $B$ . Here  $A$  union  $B$  is denoted by  $A \cup B$ 
  - \* Example:  $\{1, 2\} \cup \{2, 3\} = \{1, 2, 3\}$



- \* Similarly, we define union of three or more sets as follows

$$A_1 \cup A_2 \cup \dots \cup A_k = \bigcup_{i=1}^k A_i$$

## » Quiz

- \* If  $A$  and  $B$  are countable, then  $A \cup B$  is also countable

- \* Countable union of countable sets is countable

## » Set Operations: Intersection

## » Set Operations: Intersection


- \* The intersection of two sets is a set containing all elements that are in  $A$  and in  $B$ . Here  $A$  intersection  $B$  is denoted by  $A \cap B$

## » Set Operations: Intersection

- \* The intersection of two sets is a set containing all elements that are in  $A$  and in  $B$ . Here  $A \cap B$  is denoted by  $A \cap B$ 
  - \* Example:  $\{1, 2\} \cap \{2, 3\} = \{2\}$


## » Set Operations: Intersection

- \* The union of two sets is a set containing all elements that are in  $A$  and in  $B$ . Here  $A$  intersection  $B$  is denoted by  $A \cap B$ 
  - \* Example:  $\{1, 2\} \cap \{2, 3\} = \{2\}$



## » Set Operations: Intersection

- \* The union of two sets is a set containing all elements that are in  $A$  and in  $B$ . Here  $A$  intersection  $B$  is denoted by  $A \cap B$ 
  - \* Example:  $\{1, 2\} \cap \{2, 3\} = \{2\}$




- \* Similarly, we define intersection of three or more sets as follows



## » Set Operations: Intersection

- \* The union of two sets is a set containing all elements that are in  $A$  and in  $B$ . Here  $A$  intersection  $B$  is denoted by  $A \cap B$ 
  - \* Example:  $\{1, 2\} \cap \{2, 3\} = \{2\}$



- \* Similarly, we define intersection of three or more sets as follows

$$A_1 \cap A_2 \cap \cdots \cap A_k = \bigcap_{i=1}^k A_i$$


## » Set Operations: Complement

## » Set Operations: Complement

- \* The **complement** of a set  $A$  denoted by  $A^c$  is the set of all elements that are in the universal set  $S$ , but **not** in  $A$

## » Set Operations: Complement

- \* The **complement** of a set  $A$  denoted by  $A^c$  is the set of all elements that are in the universal set  $S$ , but **not** in  $A$



## » Set Operations: Set Difference

## » Set Operations: Set Difference


- \* The **set difference** denoted by  $A - B$  consists of elements that are in  $A$ , but **not** in  $B$

## » Set Operations: Set Difference

- \* The **set difference** denoted by  $A - B$  consists of elements that are in  $A$ , but **not** in  $B$
- \* For example, if  $A = \{1, 2, 3\}$  and  $B = \{3, 5\}$ , then  $A - B = \{1, 2\}$

## » Set Operations: Set Difference


- \* The **set difference** denoted by  $A - B$  consists of elements that are in  $A$ , but **not** in  $B$
- \* For example, if  $A = \{1, 2, 3\}$  and  $B = \{3, 5\}$ , then  $A - B = \{1, 2\}$





## » Set Operations: Set Difference

- \* The **set difference** denoted by  $A - B$  consists of elements that are in  $A$ , but **not** in  $B$
- \* For example, if  $A = \{1, 2, 3\}$  and  $B = \{3, 5\}$ , then  $A - B = \{1, 2\}$



- \* Two sets  $A$  and  $B$  are **mutually exclusive or disjoint** if they have no shared element, i.e.,  $A \cap B = \phi$

## » Cartesian Product of Sets

### Define Cartesian Product of Sets

Cartesian product of two sets  $A = \{a_1, a_2, \dots, a_m\}$  and  $B = \{b_1, b_2, \dots, b_n\}$  denoted by  $A \times B$  is defined as follows

$$A \times B = \cup_{i,j} \{(a_i, b_j)\}$$




## » Set Theory: Partition of a Set

- \* A collection of nonempty sets  $A_1, A_2, \dots$ , is a **partition** of a set  $A$  if they are **disjoint** and their union is  $A$

## » Set Theory: Partition of a Set

- \* A collection of nonempty sets  $A_1, A_2, \dots$ , is a **partition** of a set  $A$  if they are **disjoint** and their union is  $A$



## » DeMorgan and Distributive Laws

## » DeMorgan and Distributive Laws

- \* De Morgan's Law

## » DeMorgan and Distributive Laws

- \* De Morgan's Law

- \* For any sets  $A_1, A_2, \dots, A_n$ , we have



## » DeMorgan and Distributive Laws

### \* De Morgan's Law

\* For any sets  $A_1, A_2, \dots, A_n$ , we have

$$(A_1 \cup A_2 \cdots \cup A_n)^c = A_1^c \cap A_2^c \cap \cdots \cap A_n^c$$

$$(A_1 \cap A_2 \cdots \cap A_n)^c = A_1^c \cup A_2^c \cup \cdots \cup A_n^c$$

## » DeMorgan and Distributive Laws

### \* De Morgan's Law

\* For any sets  $A_1, A_2, \dots, A_n$ , we have

$$(A_1 \cup A_2 \cdots \cup A_n)^c = A_1^c \cap A_2^c \cap \cdots \cap A_n^c$$

$$(A_1 \cap A_2 \cdots \cap A_n)^c = A_1^c \cup A_2^c \cup \cdots \cup A_n^c$$

### \* Distributive Law

## » DeMorgan and Distributive Laws

### \* De Morgan's Law

\* For any sets  $A_1, A_2, \dots, A_n$ , we have

$$(A_1 \cup A_2 \cdots \cup A_n)^c = A_1^c \cap A_2^c \cap \cdots \cap A_n^c$$

$$(A_1 \cap A_2 \cdots \cap A_n)^c = A_1^c \cup A_2^c \cup \cdots \cup A_n^c$$

### \* Distributive Law

$$* A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

## » DeMorgan and Distributive Laws

### \* De Morgan's Law

\* For any sets  $A_1, A_2, \dots, A_n$ , we have

$$(A_1 \cup A_2 \cdots \cup A_n)^c = A_1^c \cap A_2^c \cap \cdots \cap A_n^c$$

$$(A_1 \cap A_2 \cdots \cap A_n)^c = A_1^c \cup A_2^c \cup \cdots \cup A_n^c$$

### \* Distributive Law

$$* A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$* A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

## » Rule of Sum using Set Language

## » Rule of Sum using Set Language


### Rule of Sum

If there is a set  $A$  with  $k$  elements, a set  $B$  with  $n$  elements and these sets do not have common elements, then the set  $A \cup B$  has  $n + k$  elements

## » Rule of Sum using Set Language

### Rule of Sum

If there is a set  $A$  with  $k$  elements, a set  $B$  with  $n$  elements and these sets do not have common elements, then the set  $A \cup B$  has  $n + k$  elements



## » Remark on Rule of Sum



## » Remark on Rule of Sum


### Rule of sum

Can we apply rule of sum when  $A$  and  $B$  intersect as follows?

## » Remark on Rule of Sum

### Rule of sum


Can we apply rule of sum when  $A$  and  $B$  intersect as follows?



## » Remark on Rule of Sum

### Rule of sum

Can we apply rule of sum when  $A$  and  $B$  intersect as follows?




- \* If we consider  $|A| + |B|$  as in sum rule, then we will be wrong

## » Remark on Rule of Sum

### Rule of sum

Can we apply rule of sum when  $A$  and  $B$  intersect as follows?



- \* If we consider  $|A| + |B|$  as in sum rule, then we will be wrong
- \* We will count elements that belong to both  $A$  and  $B$  twice
- \*  $|A \cup B| = |A| + |B| - |A \cap B|$  (Inclusion-Exclusion Principle)

## » Applications of sum rule

## » Applications of sum rule

Sum rule

Count all integers from 1 to 10 that are divisible by 2 or by 3

## » Applications of sum rule

### Sum rule

Count all integers from 1 to 10 that are divisible by 2 or by 3

\* Let us count all the numbers from 1 to 10:

1, 2, 3, 4, 5, 6, 7, 8, 9, 10

## » Applications of sum rule

### Sum rule

Count all integers from 1 to 10 that are divisible by 2 or by 3

- \* Let us count all the numbers from 1 to 10:

1, 2, 3, 4, 5, 6, 7, 8, 9, 10

- \* Here 6 is divisible both by 2 and by 3. Hence, rule of sum can't be applied!



## » Sum Rule: Example

## » Sum Rule: Example


### Number of Paths

Suppose there are several points connected by arrows. There is a starting point  $s$  (called source) and a final point  $t$  (called sink). How many different ways are there to get from  $s$  to  $t$ ?

## » Sum Rule: Example

### Number of Paths


Suppose there are several points connected by arrows. There is a starting point  $s$  (called source) and a final point  $t$  (called sink). How many different ways are there to get from  $s$  to  $t$ ?



## » Sum Rule: Example

### Number of Paths

Suppose there are several points connected by arrows. There is a starting point  $s$  (called source) and a final point  $t$  (called sink). How many different ways are there to get from  $s$  to  $t$ ?




\* counting can be done recursively

## » Sum Rule: Example

### Number of Paths

Suppose there are several points connected by arrows. There is a starting point  $s$  (called source) and a final point  $t$  (called sink). How many different ways are there to get from  $s$  to  $t$ ?



- \* counting can be done recursively
- \* for each node count the number of paths from  $s$  to this node
  - \* sum rule will be used



## » Product Rule

### Product Rule

If there are  $k$  object of the first type and there are  $n$  object of the second type, then there are  $k \times n$  pairs of objects, the first of the first type and the second of the second type

## » Product Rule

### Product Rule

If there are  $k$  object of the first type and there are  $n$  object of the second type, then there are  $k \times n$  pairs of objects, the first of the first type and the second of the second type

Pizza options



Soda options





## » Product Rule

### Product Rule

If there are  $k$  object of the first type and there are  $n$  object of the second type, then there are  $k \times n$  pairs of objects, the first of the first type and the second of the second type

Pizza options




Soda options



\* Hence, there are  $4 \times 3 = 12$  options

## » All possible pairs

## » All possible pairs



## » Rule of Product Using Sets

## » Rule of Product Using Sets

### Product Rule

If there is a finite set  $A$  and a finite set  $B$ , then there are  $|A| \times |B|$  pairs of objects, the first from  $A$ , and the second from  $B$

## » Tuples: Application of Product Rule

## » Tuples: Application of Product Rule

### Tuples

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

## » Tuples: Application of Product Rule

### Tuples

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

- \* How many different 1-letter passwords are possible?



## » Tuples: Application of Product Rule

### Tuples

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

- \* How many different 1-letter passwords are possible?
- \* What about 2-letters?

## » Tuples: Application of Product Rule

### Tuples

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

- \* How many different 1-letter passwords are possible?
- \* What about 2-letters?
- \* How many different 3-letter words are possible?

## » Tuples: Application of Product Rule

### Tuples

How many different 5-symbol passwords can we create using lower case Latin letters only? (the size of the alphabet is 26)

- \* How many different 1-letter passwords are possible?
- \* What about 2-letters?
- \* How many different 3-letter words are possible?
- \* Can you now answer the question above?

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

- \* can apply the same argument as above! (product rule!)

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

- \* can apply the same argument as above! (product rule!)
- \* there are  $n$  possibilities to pick the first letter

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

- \* can apply the same argument as above! (product rule!)
- \* there are  $n$  possibilities to pick the first letter
- \* similarly, There are  $n$  possibilities to pick the second letter, and so on...

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols?

- \* can apply the same argument as above! (product rule!)
- \* there are  $n$  possibilities to pick the first letter
- \* similarly, There are  $n$  possibilities to pick the second letter, and so on...
- \* thus, the answer is a product of  $n$  by itself  $k$  times, that is  $n^k$

## » Number of Number Plates



## » Number of Number Plates

Consider the typical vehicle number plate in India

## » Number of Number Plates

Consider the typical vehicle number plate in India



## » Number of Number Plates

Consider the typical vehicle number plate in India



- \* the first two letters denote state

## » Number of Number Plates

Consider the typical vehicle number plate in India



- \* the first two letters denote state
- \* the following two-digit number stands for district number

## » Number of Number Plates

Consider the typical vehicle number plate in India



- \* the first two letters denote state
- \* the following two-digit number stands for district number
- \* the following two-letter is RTO series

## » Number of Number Plates

Consider the typical vehicle number plate in India



- \* the first two letters denote state
- \* the following two-digit number stands for district number
- \* the following two-letter is RTO series
- \* the last four digits is the vehicle number ranging from 0000 to 9999

## » Number of Number Plates

Consider the typical vehicle number plate in India



- \* the first two letters denote state
- \* the following two-digit number stands for district number
- \* the following two-letter is RTO series
- \* the last four digits is the vehicle number ranging from 0000 to 9999

### Question

How many vehicles are there?



## » Tuples with Restrictions (Combine Sum and Product Rule)



## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

- \* Numbers between 0 and 9999 are sequences of digits of length 4

## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

- \* Numbers between 0 and 9999 are sequences of digits of length 4
- \* Three digital numbers correspond to sequences starting with 0

## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

- \* Numbers between 0 and 9999 are sequences of digits of length 4
- \* Three digital numbers correspond to sequences starting with 0
- \* We can place the unique 5 at any of four positions

## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

- \* Numbers between 0 and 9999 are sequences of digits of length 4
- \* Three digital numbers correspond to sequences starting with 0
- \* We can place the unique 5 at any of four positions
- \* This gives us 4 cases; if we compute the number of sequences in all four cases, we can get the answer by the rule of sum

## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

- \* Numbers between 0 and 9999 are sequences of digits of length 4
- \* Three digital numbers correspond to sequences starting with 0
- \* We can place the unique 5 at any of four positions
- \* This gives us 4 cases; if we compute the number of sequences in all four cases, we can get the answer by the rule of sum
- \* If we fix 5 in one place, then there are  $5 \times 5 \times 5 = 125$  sequences



## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

- \* Numbers between 0 and 9999 are sequences of digits of length 4
- \* Three digital numbers correspond to sequences starting with 0
- \* We can place the unique 5 at any of four positions
- \* This gives us 4 cases; if we compute the number of sequences in all four cases, we can get the answer by the rule of sum
- \* If we fix 5 in one place, then there are  $5 \times 5 \times 5 = 125$  sequences
- \* There are 4 ways to arrange 5 among 4 places

## » Tuples with Restrictions (Combine Sum and Product Rule)

### Question

How many integer numbers are there between 0 and 9999 that have exactly one 5 digit?

- \* Numbers between 0 and 9999 are sequences of digits of length 4
- \* Three digit numbers correspond to sequences starting with 0
- \* We can place the unique 5 at any of four positions
- \* This gives us 4 cases; if we compute the number of sequences in all four cases, we can get the answer by the rule of sum
- \* If we fix 5 in one place, then there are  $5 \times 5 \times 5 = 125$  sequences
- \* There are 4 ways to arrange 5 among 4 places
- \* Hence, there are  $4 \times 125 = 500$  four digit numbers below 10,000 with exactly one 5





## » Permutations

### Definition

Tuples of length  $k$  without repetitions are called  $k$ —permutations

## » Permutations

### Definition

Tuples of length  $k$  without repetitions are called  $k$ —permutations

### Question

## » Permutations

### Definition

Tuples of length  $k$  without repetitions are called  $k$ -permutations

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

## » Permutations

### Definition

Tuples of length  $k$  without repetitions are called  $k$ -permutations

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

- \* Observe that if  $n < k$  then there are no  $k$ -permutations: there are simply not enough different letters

## » Permutations

### Definition

Tuples of length  $k$  without repetitions are called  $k$ –permutations

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

- \* Observe that if  $n < k$  then there are no  $k$ –permutations: there are simply not enough different letters
- \* So it is enough to solve the problem for the case  $k \leq n$

## » Permutations

### Definition

Tuples of length  $k$  without repetitions are called  $k$ -permutations

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

- \* Observe that if  $n < k$  then there are no  $k$ -permutations: there are simply not enough different letters
- \* So it is enough to solve the problem for the case  $k \leq n$

1	2	3	...	$k$
*	*	*	...	*
$n$	$n-1$	$n-2$		

## » Permutations

### Definition

Tuples of length  $k$  without repetitions are called  $k$ -permutations

### Question

Suppose we have a set of  $n$  symbols. How many different sequences of length  $k$  we can form out of these symbols if we are not allowed to use the same symbol twice?

- \* Observe that if  $n < k$  then there are no  $k$ -permutations: there are simply not enough different letters
- \* So it is enough to solve the problem for the case  $k \leq n$

1	2	3	...	$k$
*	*	*	...	*
$n$	$n-1$	$n-2$		

- \* Hence there are

$$n \times (n - k) \times \cdots (n - k + 1)$$

$k$ -permutations, which is  $n!/(n - k)!$



## » Permutation Examples

## » Permutation Examples

### Question

In how many ways we can arrange  $n$  different books in  $n$  different bins on shelf?

## » Permutation Examples

### Question

In how many ways we can arrange  $n$  different books in  $n$  different bins on shelf?

### Answer

Hint: Use previous result with  $k = n$ .