

Probability and Statistics: Lecture-27

Monsoon-2020

by Dr. Pawan Kumar (IIIT, Hyderabad)

on October 14, 2020

» Mixed Random Variable...

» Mixed Random Variable...

Example of mixed random variable

Let X be a continuous random variable with the following PDF


$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Let


$$Y = g(X) = \begin{cases} X & 0 \leq X \leq \frac{1}{2} \\ \frac{1}{2} & X > \frac{1}{2} \end{cases}$$

Find the CDF of Y .

» Plot of the Mixed Random Variable Example




» Plot of the Mixed Random Variable Example




- * the CDF is not continuous, so Y is not a continuous random variable

» Plot of the Mixed Random Variable Example




- * the CDF is not continuous, so Y is not a continuous random variable
- * the CDF is not in the staircase form, so it is not a discrete random variable either

» Plot of the Mixed Random Variable Example




- * the CDF is not continuous, so Y is not a continuous random variable
- * the CDF is not in the staircase form, so it is not a discrete random variable either
- * It is indeed a mixed random variable

» Plot of the Mixed Random Variable Example




- * the CDF is not continuous, so Y is not a continuous random variable
- * the CDF is not in the staircase form, so it is not a discrete random variable either
- * It is indeed a mixed random variable
- * there is jump at $y = 1/2$

» Plot of the Mixed Random Variable Example



- * the CDF is not continuous, so Y is not a continuous random variable
- * the CDF is not in the staircase form, so it is not a discrete random variable either
- * It is indeed a mixed random variable
- * there is jump at $y = 1/2$
- * amount of jump is $1 - 1/4 = 3/4$

» Plot of the Mixed Random Variable Example



- * the CDF is not continuous, so Y is not a continuous random variable
- * the CDF is not in the staircase form, so it is not a discrete random variable either
- * It is indeed a mixed random variable
- * there is jump at $y = 1/2$
- * amount of jump is $1 - 1/4 = 3/4$
- * CDF is continuous at other points

» CDF of mixed RV as a sum of continuous and discrete RV...

» CDF of mixed RV as a sum of continuous and discrete RV...

- * In the previous example, CDF of Y was

» CDF of mixed RV as a sum of continuous and discrete RV...

* In the previous example, CDF of Y was

$$F_Y(y) = \begin{cases} 1 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & \text{otherwise} \end{cases}$$

» CDF of mixed RV as a sum of continuous and discrete RV...

- * In the previous example, CDF of Y was

$$F_Y(y) = \begin{cases} 1 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & \text{otherwise} \end{cases}$$

- * The CDF can be written as a sum of continuous and discrete part

» CDF of mixed RV as a sum of continuous and discrete RV...

- * In the previous example, CDF of Y was

$$F_Y(y) = \begin{cases} 1 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & \text{otherwise} \end{cases}$$

- * The CDF can be written as a sum of continuous and discrete part

$$F_Y(y) = \underbrace{C(y)} + \underbrace{D(y)},$$

» CDF of mixed RV as a sum of continuous and discrete RV...

- * In the previous example, CDF of Y was

$$F_Y(y) = \begin{cases} 1 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & \text{otherwise} \end{cases}$$

- * The CDF can be written as a sum of continuous and discrete part

$$F_Y(y) = C(y) + D(y),$$

where the continuous part is

» CDF of mixed RV as a sum of continuous and discrete RV...

- * In the previous example, CDF of Y was

$$F_Y(y) = \begin{cases} 1 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & \text{otherwise} \end{cases}$$

- * The CDF can be written as a sum of continuous and discrete part

$$F_Y(y) = C(y) + D(y),$$

where the continuous part is

$$C(y) = \begin{cases} 1/4 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & y < 0 \end{cases}$$

» CDF of mixed RV as a sum of continuous and discrete RV...

- * In the previous example, CDF of Y was

$$F_Y(y) = \begin{cases} 1 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & \text{otherwise} \end{cases}$$

- * The CDF can be written as a sum of continuous and discrete part

$$F_Y(y) = C(y) + D(y),$$

where the continuous part is

$$C(y) = \begin{cases} 1/4 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & y < 0 \end{cases}$$

and the discrete part is

» CDF of mixed RV as a sum of continuous and discrete RV...

* In the previous example, CDF of Y was

$$F_Y(y) = \begin{cases} 1 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & \text{otherwise} \end{cases}$$

* The CDF can be written as a sum of continuous and discrete part

$$F_Y(y) = C(y) + D(y),$$

where the continuous part is

$$C(y) = \begin{cases} 1/4 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & y < 0 \end{cases}$$

and the discrete part is

$$D(y) = \begin{cases} 3/4 & y \geq 1/2 \\ 0 & y < 1/2 \end{cases}$$

← jump point

» CDF of a mixed RV as a sum of Continuous and Discrete CDF..

» CDF of a mixed RV as a sum of Continuous and Discrete CDF...

CDF of Mixed Random Variable

» CDF of a mixed RV as a sum of Continuous and Discrete CDF...

CDF of Mixed Random Variable

The CDF of mixed random variable Y can be written as a sum of continuous and discrete/staircase function

$$F_Y(y) = \underline{C(y)} + \underline{D(y)}.$$

» CDF of a mixed RV as a sum of Continuous and Discrete CDF...

CDF of Mixed Random Variable

The CDF of mixed random variable Y can be written as a sum of continuous and discrete/staircase function

$$F_Y(y) = \underbrace{C(y)} + D(y).$$

We differentiate the continuous part $\underbrace{c(y)} = \frac{dC(y)}{dy}$, where ever $\underbrace{C(y)}$ is differentiable.
↑ additional assumption

» CDF of a mixed RV as a sum of Continuous and Discrete CDF..

CDF of Mixed Random Variable

The CDF of mixed random variable Y can be written as a sum of continuous and discrete/staircase function

$$F_Y(y) = C(y) + D(y).$$

We differentiate the continuous part $c(y) = \frac{dC(y)}{dy}$, where ever $C(y)$ is differentiable.

Let $\{y_1, y_2, \dots\}$ be the set of jump points of $D(y)$, i.e., the points for which $P(Y = y_k) > 0$.

» CDF of a mixed RV as a sum of Continuous and Discrete CDF..

CDF of Mixed Random Variable

The CDF of mixed random variable Y can be written as a sum of continuous and discrete/staircase function

$$F_Y(y) = C(y) + D(y).$$

$$\int \frac{f_X(x)}{f_X(x)} dx = 1 \quad \sum_i P_X(x_i) = 1$$

We differentiate the continuous part $c(y) = \frac{dC(y)}{dy}$, where ever $C(y)$ is differentiable.

Let $\{y_1, y_2, \dots\}$ be the set of jump points of $D(y)$, i.e., the points for which $P(Y = y_k) > 0$. We have

$$\int_{-\infty}^{\infty} \underbrace{c(y)}_{\text{PDF of } C} dy + \sum_{y_k} \underbrace{P(Y = y_k)}_{\text{PMF of } D} = 1$$

» CDF of a mixed RV as a sum of Continuous and Discrete CDF...

CDF of Mixed Random Variable

The CDF of mixed random variable Y can be written as a sum of continuous and discrete/staircase function

$$F_Y(y) = C(y) + D(y).$$

We differentiate the continuous part $c(y) = \frac{dC(y)}{dy}$, where ever $C(y)$ is differentiable.

Let $\{y_1, y_2, \dots\}$ be the set of jump points of $D(y)$, i.e., the points for which $P(Y = y_k) > 0$. We have

$$\int_{-\infty}^{\infty} c(y) dy + \sum_{y_k} P(Y = y_k) = 1$$

The expected value of Y can be obtained as

» CDF of a mixed RV as a sum of Continuous and Discrete CDF..

CDF of Mixed Random Variable

The CDF of mixed random variable Y can be written as a sum of continuous and discrete/staircase function

$$F_Y(y) = C(y) + D(y).$$

We differentiate the continuous part $c(y) = \frac{dC(y)}{dy}$, where ever $C(y)$ is differentiable.

Let $\{y_1, y_2, \dots\}$ be the set of jump points of $D(y)$, i.e., the points for which $P(Y = y_k) > 0$. We have


$$\int_{-\infty}^{\infty} c(y) dy + \sum_{y_k} P(Y = y_k) = 1$$

The expected value of Y can be obtained as


$$E[Y] = \int_{-\infty}^{\infty} \underbrace{y c(y)} dy + \sum_{y_k} \underbrace{y_k P(Y = y_k)}$$

» Check the Validity of CDF of Mixed RV...

» Check the Validity of CDF of Mixed RV...




» Check the Validity of CDF of Mixed RV...



$$F_Y(y) = \underline{C(y)} + \underline{D(y)},$$


» Check the Validity of CDF of Mixed RV...



$$F_Y(y) = C(y) + D(y),$$

where the continuous part is

» Check the Validity of CDF of Mixed RV...




$$F_Y(y) = C(y) + D(y),$$

where the continuous part is

← best class

$$C(y) = \begin{cases} 1/4 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & y < 0 \end{cases}$$

» Check the Validity of CDF of Mixed RV...




$$F_Y(y) = C(y) + D(y),$$

where the continuous part is

$$C(y) = \begin{cases} 1/4 & y \geq 1/2 \\ y^2 & 0 \leq y < 1/2 \\ 0 & y < 0 \end{cases}$$

and the discrete part is

» Check the Validity of CDF of Mixed RV...



Check that

$$\int_{-\infty}^{\infty} c(y) dy + \sum_{y_k} P(Y = y_k) = 1$$

$$\frac{1}{4} + \left(\frac{3}{4}\right) = 1 = \int_{-\infty}^{\infty} c(y) dy = \int_{-\infty}^{\infty} 2y dy = 2 \left[\frac{y^2}{2} \right]_0^{1/2} = \left(\frac{1}{4}\right)$$

Handwritten notes:

PMP: $\frac{3}{4}$

PDF: $\frac{1}{4}$

$F_Y(y) = C(y) + D(y)$

where the continuous part is

$c(y) = 0$

$C(y) = \begin{cases} y^2 & 0 \leq y < 1/2 \\ 0 & y < 0 \end{cases}$

$c(y) = 2y$

$C(y) = 0$

$D(y) = \begin{cases} 3/4 & y \geq 1/2 \\ 0 & y < 1/2 \end{cases}$

and the discrete part is

$R_D(y) = \{1/2\}$

$C(y) = \begin{cases} 0 & y > 1/2 \\ 2y & 0 \leq y < 1/2 \\ 0 & y < 0 \end{cases}$

» Computing Probabilities of Mixed Random Variable...

» Computing Probabilities of Mixed Random Variable...

Problem: Computing probabilities of mixed RV

Let X be a continuous random variable with the following PDF

» Computing Probabilities of Mixed Random Variable...

Problem: Computing probabilities of mixed RV

Let X be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

» Computing Probabilities of Mixed Random Variable...

Problem: Computing probabilities of mixed RV

Let X be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Let

$$Y = g(X) = \begin{cases} X & 0 \leq X \leq \frac{1}{2} \\ \frac{1}{2} & X > \frac{1}{2} \end{cases}$$

» Computing Probabilities of Mixed Random Variable...

Problem: Computing probabilities of mixed RV

Let X be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Let

$$Y = g(X) = \begin{cases} X & 0 \leq X \leq \frac{1}{2} \\ \frac{1}{2} & X > \frac{1}{2} \end{cases}$$

* Find $P(1/4 \leq Y \leq 3/8)$

» Computing Probabilities of Mixed Random Variable...

Problem: Computing probabilities of mixed RV

Let X be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Let

$$Y = g(X) = \begin{cases} X & 0 \leq X \leq \frac{1}{2} \\ \frac{1}{2} & X > \frac{1}{2} \end{cases}$$

- * Find $P(1/4 \leq Y \leq 3/8)$
- * Find $P(Y \geq 1/4)$

» Computing Probabilities of Mixed Random Variable...

Problem: Computing probabilities of mixed RV

Let X be a continuous random variable with the following PDF

$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Let

$$Y = g(X) = \begin{cases} X & 0 \leq X \leq \frac{1}{2} \\ \frac{1}{2} & X > \frac{1}{2} \end{cases}$$

* Find $P(1/4 \leq Y \leq 3/8)$

* Find $P(Y \geq 1/4)$

* Find $E[Y]$

» Answer to previous problem...

$$\textcircled{a} P\left(\frac{1}{4} \leq Y \leq \frac{3}{8}\right) = \underbrace{F_Y\left(\frac{3}{8}\right) - F_Y\left(\frac{1}{4}\right)}_{P\left(\frac{1}{4} < Y \leq \frac{3}{8}\right)} + \underbrace{P_Y\left(\frac{1}{4}\right)}_{\substack{\text{add} \\ \text{back} \\ \text{because} \\ \text{discr. r.v.}}} \\ = \left(\frac{3}{8}\right)^2 - \left(\frac{1}{4}\right)^2 + 0$$

Rk: For $P_Y\left(\frac{1}{4}\right)$ we look into $D(Y)$.

$$\textcircled{b} P\left(Y > \frac{1}{4}\right) = 1 - P\left(Y < \frac{1}{4}\right) \\ = 1 - F_Y\left(\frac{1}{4}\right) + P\left(Y = \frac{1}{4}\right) \xrightarrow{0} \\ = 1 - \left(\frac{1}{4}\right)^2 = \frac{15}{16}$$

$$\textcircled{c} E[Y] \quad c(y) = \frac{dL(y)}{dy} = \begin{cases} 2y & 0 \leq y \leq \frac{1}{2} \\ 0 & \text{otherwise} \end{cases}$$

$$E[Y] = \int_0^{1/2} y \cdot 2y \, dy + \frac{1}{2} \cdot P\left(Y = \frac{1}{2}\right) = \frac{2}{3} \cdot \frac{1}{8} + \frac{1}{2} \cdot \frac{3}{4} = \frac{1}{12} + \frac{3}{8} = \frac{11}{24}$$

$$F_Y(a) = P(Y \leq a)$$

Assuming a mixed r.v.

$$F_Y(a) = P(Y < a) + P(Y = a) \quad \left[\begin{array}{l} \text{need to} \\ \text{be careful} \\ \text{in Discr.} \end{array} \right]$$

$$\Rightarrow P(Y < a) = F_Y(a) - P(Y = a)$$

If X is mixed r.v., then.

$$P(a < X \leq b) \neq P(a \leq X \leq b)$$

Recall: How was $F_Y(a)$

$$F_Y(a) = P(Y \leq a) \leftarrow \underbrace{P(Y < a)}_{+ P(Y = a)}$$

» Solved Problem 1

» Solved Problem 1

Problem 1

Let X be a random variable with CDF

» Solved Problem 1

Problem 1

Let X be a random variable with CDF

$$F_X(x) = \begin{cases} 1 & x \geq 1 \\ \frac{1}{2} + \frac{x}{2} & 0 \leq x < 1 \\ 0 & x < 0 \end{cases}$$


» Solved Problem 1

Problem 1

Let X be a random variable with CDF

$$F_X(x) = \begin{cases} 1 & x \geq 1 \\ \frac{1}{2} + \frac{x}{2} & 0 \leq x < 1 \\ 0 & x < 0 \end{cases}$$

1. Is the RV X discrete, continuous, or mixed type?



» Solved Problem 1

Problem 1

Let X be a random variable with CDF

$$F_X(x) = \begin{cases} 1 & x \geq 1 \\ \frac{1}{2} + \frac{x}{2} & 0 \leq x < 1 \\ 0 & x < 0 \end{cases}$$

1. Is the RV X discrete, continuous, or mixed type?
2. Find the PDF of X , $f_X(x)$

» Solved Problem 1

Problem 1

Let X be a random variable with CDF

$$F_X(x) = \begin{cases} 1 & x \geq 1 \\ \frac{1}{2} + \frac{x}{2} & 0 \leq x < 1 \\ 0 & x < 0 \end{cases}$$

1. Is the RV X discrete, continuous, or mixed type?
2. Find the PDF of X , $f_X(x)$
3. Find $E(e^X)$

» Solved Problem 1

Problem 1

Let X be a random variable with CDF

$$F_X(x) = \begin{cases} 1 & x \geq 1 \\ \frac{1}{2} + \frac{x}{2} & 0 \leq x < 1 \\ 0 & x < 0 \end{cases}$$

1. Is the RV X discrete, continuous, or mixed type?
2. Find the PDF of X , $f_X(x)$
3. Find $E(e^X)$
4. Find $P(X = 0 \mid X \leq 0.5)$

» Answer to previous problem...

» Answer to previous problem...

» Solved Problem 2

Problem 2

Let $X \sim \text{Uniform}(-2, 2)$ be a continuous random variable. Let $Y = g(X)$ where

$$g(x) = \begin{cases} 1 & x > 1 \\ x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \\ & x < 0 \end{cases}$$

Find the CDF of Y .

$$R_Y = [0, 1]$$

$$F_Y(0) = 0$$
$$F_Y(1) = 1$$

$$y < 0$$
$$y \geq 1$$

» Answer to previous problem...

For $0 < y < 1$

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(X \leq y) \\ &= \cancel{F_X}(y) = \int_{-2}^y \frac{1}{4} dx \\ &= \frac{1}{4} [x]_{-2}^y = \frac{1}{4}(y+2) \end{aligned}$$

CDF of Y

$$F_Y(y) = \begin{cases} \frac{1}{4}(y+2), & 0 \leq y < 1 \\ 1, & y \geq 1 \\ 0, & \text{otherwise} \end{cases}$$

Recall

$$X \sim U(a, b)$$

$$f_X(x) = \begin{cases} \frac{1}{b-a} \\ 0 \end{cases}$$

for $a \leq x \leq b$
 $x < a$ or $x > b$

$$R_X = [-2, 2]$$

$$F_Y(a) = \int_{-\infty}^a f_X(x) dx$$

» Definition of Joint Probability Mass Function...

$$P(X=x, Y=y)$$

» Definition of Joint Probability Mass Function...

Definition of Joint Probability Mass Function

The **joint probability mass function** of two **discrete** random variables X and Y is defined as follows

» Definition of Joint Probability Mass Function...

Definition of Joint Probability Mass Function

The **joint probability mass function** of two **discrete** random variables X and Y is defined as follows

$$\underbrace{P_{XY}(x, y)} = P(X = x, Y = y).$$

» Definition of Joint Probability Mass Function...

Definition of Joint Probability Mass Function

The **joint probability mass function** of two **discrete** random variables X and Y is defined as follows

$$P_{XY}(x, y) = P(X = x, Y = y).$$

* The **joint range** for X and Y is

» Definition of Joint Probability Mass Function...

Definition of Joint Probability Mass Function

The **joint probability mass function** of two **discrete** random variables X and Y is defined as follows

$$P_{XY}(x, y) = P(X = x, Y = y).$$

* The **joint range** for X and Y is

$$\underline{R_{XY}} = \{ \underline{(x, y)} \mid \underline{P_{XY}} > 0 \}$$

» Definition of Joint Probability Mass Function...

Definition of Joint Probability Mass Function

The **joint probability mass function** of two **discrete** random variables X and Y is defined as follows

$$P_{XY}(x, y) = P(X = x, Y = y).$$

- * The **joint range** for X and Y is

$$R_{XY} = \{(x, y) \mid P_{XY} > 0\}$$

- * In particular, if $R_X = \{x_1, x_2, \dots\}$,

» Definition of Joint Probability Mass Function...

Definition of Joint Probability Mass Function

The **joint probability mass function** of two discrete random variables X and Y is defined as follows

$$P_{XY}(x, y) = P(X = x, Y = y).$$

- * The **joint range** for X and Y is

$$R_{XY} = \{(x, y) \mid P_{XY} > 0\}$$

- * In particular, if $R_X = \{\underline{x_1}, \underline{x_2}, \dots\}$, $R_Y = \{\underline{y_1}, \underline{y_2}, \dots\}$, then

» Definition of Joint Probability Mass Function...

Definition of Joint Probability Mass Function

The joint probability mass function of two discrete random variables X and Y is defined as follows

$$P_{XY}(x, y) = P(X = x, Y = y).$$

* The joint range for X and Y is

$$R_{XY} = \{(x, y) \mid P_{XY} > 0\}$$

* In particular, if $R_X = \{x_1, x_2, \dots\}$, $R_Y = \{y_1, y_2, \dots\}$, then

$$R_{XY} \subset R_X \times R_Y = \{(x_i, y_j) \mid x_i \in R_X, y_j \in R_Y\}$$

* Sum of joint probabilities must sum to 1: $\sum_{(x_i, y_j) \in R_{XY}} P_{XY}(x_i, y_j) = 1$

» Computing Probabilities with Joint Distributions...

» Computing Probabilities with Joint Distributions...

Computing Joint Probability

For any $A \subset \mathbb{R}^2$, we have

» Computing Probabilities with Joint Distributions...

Computing Joint Probability

For any $A \subset \mathbb{R}^2$, we have

$$P(\underbrace{(X, Y)}_{\in A}) = \sum_{\underbrace{(x_i, y_j)}_{\in (A \cap R_{XY})}} P_{XY}(x_i, y_j)$$

» Computing Probabilities with Joint Distributions...

Computing Joint Probability

For any $A \subset \mathbb{R}^2$, we have

$$P((X, Y) \in A) = \sum_{(x_i, y_j) \in (A \cap R_{XY})} P_{XY}(x_i, y_j)$$

Marginal PMF: Computing Individual Probabilities from Joint PMF

» Computing Probabilities with Joint Distributions...

Computing Joint Probability

For any $A \subset \mathbb{R}^2$, we have

$$P((X, Y) \in A) = \sum_{(x_i, y_j) \in (A \cap R_{XY})} P_{XY}(x_i, y_j)$$

Marginal PMF: Computing Individual Probabilities from Joint PMF

We can obtain PMF of X from its joint PMF with Y as follows

» Computing Probabilities with Joint Distributions...

Computing Joint Probability

For any $A \subset \mathbb{R}^2$, we have

$$P((X, Y) \in A) = \sum_{(x_i, y_j) \in (A \cap R_{XY})} P_{XY}(x_i, y_j)$$

Marginal PMF: Computing Individual Probabilities from Joint PMF

We can obtain **PMF** of X from its **joint PMF** with Y as follows

$$P_X(x) = P(X = x) = \sum_{y_j \in R_Y} P(X = x, Y = y_j) = \sum_{y_j \in R_Y} P_{XY}(x, y_j), \quad \text{for any } x \in R_X$$

Handwritten notes: "fixed" with an arrow pointing to x in $P_{XY}(x, y_j)$; a bracket under $y_j \in R_Y$ with an arrow pointing to the summation symbol.

» Computing Probabilities with Joint Distributions...

Computing Joint Probability

For any $A \subset \mathbb{R}^2$, we have

$$P((X, Y) \in A) = \sum_{(x_i, y_j) \in (A \cap R_{XY})} P_{XY}(x_i, y_j)$$

Marginal PMF: Computing Individual Probabilities from Joint PMF

We can obtain **PMF** of X from its **joint PMF** with Y as follows

$$P_X(x) = P(X = x) = \sum_{y_j \in R_Y} P(X = x, Y = y_j) = \sum_{y_j \in R_Y} P_{XY}(x, y_j), \quad \text{for any } x \in R_X$$

We call $P_X(x)$ the **marginal PMF** of X . Similarly, we have

» Computing Probabilities with Joint Distributions...

Computing Joint Probability

For any $A \subset \mathbb{R}^2$, we have

$$P((X, Y) \in A) = \sum_{(x_i, y_j) \in (A \cap R_{XY})} P_{XY}(x_i, y_j)$$

Marginal PMF: Computing Individual Probabilities from Joint PMF

We can obtain **PMF** of X from its **joint PMF** with Y as follows

$$\underbrace{P_X(x)} = P(X = x) = \sum_{y_j \in R_Y} P(X = x, Y = y_j) = \sum_{y_j \in R_Y} \underbrace{P_{XY}(x, y_j)}, \quad \text{for any } x \in R_X$$

We call $P_X(x)$ the **marginal PMF** of X . Similarly, we have

$$\underbrace{P_Y(y)} = \sum_{x_i \in R_X} \underbrace{P_{XY}(x_i, y)}, \quad \text{for any } y \in R_Y$$

Handwritten notes: An arrow points from the y in the joint PMF to the y in the marginal PMF. The sum is over x_i .

» Solved Example

Example

Let X and Y be two random variables with joint PMF as follows:

» Solved Example

Example

Let X and Y be two random variables with **joint PMF** as follows:


	$Y=0$	$Y=1$	$Y=2$
$X=0$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{8}$
$X=1$	$\frac{1}{8}$	$\frac{1}{6}$	$\frac{1}{6}$

» Solved Example

Example

Let X and Y be two random variables with **joint PMF** as follows:

	$Y=0$	$Y=1$	$Y=2$
$X=0$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{8}$
$X=1$	$\frac{1}{8}$	$\frac{1}{6}$	$\frac{1}{6}$




1. Find $P(X=0, Y \leq 1)$

» Solved Example

Example

Let X and Y be two random variables with **joint PMF** as follows:

	$Y=0$	$Y=1$	$Y=2$
$X=0$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{8}$
$X=1$	$\frac{1}{8}$	$\frac{1}{6}$	$\frac{1}{6}$




1. Find $P(X=0, Y \leq 1)$
2. Find the **marginal PMFs** of X and Y

» Solved Example

Example

Let X and Y be two random variables with **joint PMF** as follows:

	$Y=0$	$Y=1$	$Y=2$
$X=0$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{8}$
$X=1$	$\frac{1}{8}$	$\frac{1}{6}$	$\frac{1}{6}$



1. Find $P(X=0, Y \leq 1)$
2. Find the **marginal PMFs** of X and Y
3. Find $P(Y=1 \mid X=0)$

» Solved Example


Example

Let X and Y be two random variables with **joint PMF** as follows:

	$Y=0$	$Y=1$	$Y=2$
$X=0$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{8}$
$X=1$	$\frac{1}{8}$	$\frac{1}{6}$	$\frac{1}{6}$

Sum the row to get $P_X(0)$

Sum to get $P_X(1)$



1. Find $P(X=0, Y \leq 1)$
2. Find the **marginal PMFs** of X and Y
3. Find $P(Y=1 | X=0)$
4. Are X and Y **independent**?

» Answer to previous problem...

$$\textcircled{1} P(X=0, Y \leq 1) = P_{XY}(0,0) + P_{XY}(0,1)$$

$$= \frac{1}{6} + \frac{1}{4} = \frac{5}{12}$$

② Find $P_X(x)$ & $P_Y(y)$ (Marginals)

$$R_X = \{0, 1\}, \quad R_Y = \{0, 1, 2\}$$

$$P_X(0) = P_{XY}(0,0) + P_{XY}(0,1) + P_{XY}(0,2)$$

$$= \left(\frac{1}{6} + \frac{1}{12} + \frac{1}{6} \right) = \frac{2}{3}$$

$$P_X(1) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2}$$

$$\textcircled{3} P(Y=1 | X=0) = \frac{P(X=0, Y=1)}{P(X=0)}$$

$$= \frac{P_{XY}(0,1)}{P_X(X=0)} = \frac{1/4}{2/3} = \frac{3}{8}$$

④ X & Y ind?

$$P(Y=1 | X=0) = \frac{P_{XY}(0,1)}{P_X(0)}$$

$$= \frac{1/4}{1/6 + 1/4 + 1/8} = \frac{1/4}{3/4} = \frac{1}{3}$$

check this

If they are equal, then try other conditional

Recall for X, Y ind

$$P(Y=y_i | X=x_j) = P(Y=y_i)$$

$\forall i, j$