
Probability and Statistics: Lecture-35

Monsoon-2020

```
by Dr. Pawan Kumar (IIIT, Hyderabad)
on November 4, 2020
```



Example (Solved Example

Example (Solved Example

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x, y \ge 0\\ 0 & \text{otherwise} \end{cases}$$

Example (Solved Example

Let X, Y be jointly continuous RVs with joint PDF

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x,y \ge 0 \\ 0 & \text{otherwise} \end{cases}$$

• Are X and Y independent?

Example (Solved Example

Let X, Y be jointly continuous RVs with joint PDF

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x,y \ge 0 \\ 0 & \text{otherwise} \end{cases}$$

ullet Are ${\it X}$ and ${\it Y}$ independent? ${\it \bullet}$ Find ${\it E}[{\it Y} \,|\, {\it X} > 2]$

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x,y \ge 0 \\ 0 & \text{otherwise} \end{cases}$$
• Are X and Y independent? • Find $E[Y \mid X > 2]$ • Find $P(X > Y)$

» Answer to previous problem...

32

Example (Solved example

Let \boldsymbol{X} be a continuous RV with PDF

Example (Solved example)

Let X be a continuous RV with PDF

$$f_{X}(x) = egin{cases} 2x & 0 \leq x \leq 1 \\ 0 & ext{otherwise} \end{cases}$$

Example (Solved example

Let X be a continuous RV with PDF

$$f_{\mathcal{X}}(\mathbf{x}) = \begin{cases} 2\mathbf{x} & 0 \le \mathbf{x} \le 1 \\ 0 & \text{otherwise} \end{cases}$$

We are also know that given X = x, the RV Y is uniformly distributed on [-x, x].


Let X be a continuous RV with PDF

$$f_{X}(x) = \begin{cases} 2x & 0 \le x \le 1 \\ 0 & \text{otherwise} \end{cases}$$

We are also know that given X = x, the RV Y is uniformly distributed on [-x, x].

• Find the joint PDF
$$f_{XY}(x,y)$$
 • Find $P_Y(y)$ • Find $P(|Y| < X^3)$











Example (Solved example

Example (Solved example

$$f_{\mathcal{X},\mathcal{Y}}(\mathbf{x},\mathbf{y}) = egin{cases} 6\mathbf{x}\mathbf{y} & 0 \leq \mathbf{x} \leq 1, \ 0 \leq \mathbf{y} \leq \sqrt{\mathbf{x}} \\ 0 & ext{otherwise} \end{cases}$$

Example (Solved example

Let X, Y be two jointly continuous RVs with joint PDF

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

1. Plot R_{XY}

Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

- 1. Plot R_{XY}
- 2. Find $f_X(x)$ and $f_Y(y)$

Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & & ext{otherwise} \end{cases}$$

- 1. Plot R_{XY}
- 2. Find $f_X(x)$ and $f_Y(y)$
- 3. Are X and Y independent?

Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

- 1. Plot R_{XY}
- 2. Find $f_X(x)$ and $f_Y(y)$
- 3. Are X and Y independent?
- 4. Find the conditional PDF of X given Y = y, $f_{X|Y}(x \mid y)$

Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

- 1. Plot R_{XY}
- 2. Find $f_X(x)$ and $f_Y(y)$
- 3. Are X and Y independent?
- 4. Find the conditional PDF of X given Y = y, $f_{X|Y}(x \mid y)$
- 5. Find $E[X | Y = y], 0 \le y \le 1$

Example (Solved example

$$f_{X,Y}(x,y) = \begin{cases} 6xy & 0 \le x \le 1, \ 0 \le y \le \sqrt{x} \\ 0 & \text{otherwise} \end{cases}$$

- ightharpoonupPlot R_{XY}
- Find $f_X(x)$ and $f_Y(y)$
- Are X and Y independent?
- 4. Find the conditional PDF of X given Y = y, $f_{X|Y}(x \mid y)$
- **5.** Find E[X | Y = y], 0 ≤ y ≤ 1
- 6 Find $Var(X \mid Y = y)$ for $0 \le y \le 1$

*Answer to previous problem...

Frith:
$$\int fxy(x)(x) dx$$

Frith: $\int fxy(x)(x) dx$

Frith: $\int fxy(x) dx$

» Answer to previous problem...

(a)
$$\int x|y|(x|y) = \int x|y|(x|y)$$

$$= \int \frac{2}{3}x^{3}|y|^{2} = \frac{2}{1-y^{4}} \left[\frac{2}{3}\right]^{2}$$


$$= \int \frac{2}{3}x^{3}|y|^{2} = \frac{2}{1-y^{4}} \left[\frac{2}{3}\right]^{2}$$

$$= \int \frac{2}{3}x^{3}|y|^{2} = \frac{2}{1-y^{4}} \left[\frac{2}{3}\right]^{2}$$

$$= \int \frac{2}{3}x^{3} = \int \frac{2}{3}x^{3} =$$

» Answer to previous problem...

9/132



» Definition of Covariance...

Definition of Covariance

Let X and Y be two random variables.

» Definition of Covariance...

Definition of Covariance

Let *X* and *Y* be two random variables. The covariance between *X* and *Y* is defined as


» Definition of Covariance...

Definition of Covariance

Let X and Y be two random variables. The covariance between X and Y is defined as

$$Cov(X, Y) = \underbrace{E[(X - E[X])(Y - E[Y])]}_{E[XY]} = \underbrace{E[XY] - (E[X])(E[Y])}_{E[XY]}$$

Derivation



Solved Example

Example (Solved Example)


Suppose
$$X \sim \text{Uniform}(1,2)$$
 and given $X = x$, Y is exponential with parameter $\lambda = x$.

Find $Cov(X, Y)$.

Cov $(Y,Y) = E[YY] - E[Y] = E[Y]$

We have $E[Y] = \frac{3}{2}$

We have $E[Y] = E[XY] = \frac{3}{2}$
 $E[Y] = \frac{3}{2}$



» Properties of Covariance...

Properties of Covariance

1.
$$Cov(X, X) = Var(X)$$

» Properties of Covariance...

Properties of Covariance

 $\mathsf{L}.\mathsf{Cov}(X,X) = \mathsf{Var}(X)$



 \nearrow If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]

» Properties of Covariance...

Properties of Covariance

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- 4. Cov(aX, Y) = aCov(X, Y)

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- \mathcal{S} . $\mathsf{Cov}(X,Y) = \mathsf{Cov}(Y,X)$
- 4. Cov(aX, Y) = aCov(X, Y)
- 5. Cov(X + c, Y) = Cov(X, Y)


- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- 4. Cov(aX, Y) = aCov(X, Y)
- 5. Cov(X + c, Y) = Cov(X, Y)
- 6. Cov(X + Y, Z) = Cov(X, Z) + Cov(Y, Z)

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- 4. Cov(aX, Y) = aCov(X, Y)
- 5. Cov(X + c, Y) = Cov(X, Y)
- 6. Cov(X + Y, Z) = Cov(X, Z) + Cov(Y, Z)
- 7. More generally, we have

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- 4. Cov(aX, Y) = aCov(X, Y)
- $6. \operatorname{Cov}(X + c, Y) = \operatorname{Cov}(X, Y)$
- 6. Cov(X + Y, Z) = Cov(X, Z) + Cov(Y, Z)
- 7. More generally, we have

$$\operatorname{Cov}\left(\sum_{i=1}^{m}a_{i}X_{i},\sum_{j=1}^{n}b_{j}Y_{j}\right)=\sum_{i=1}^{m}\sum_{j=1}^{n}a_{i}b_{j}\operatorname{Cov}(X_{i},Y_{j})$$





» Solved Example...

Let X and Y be two independent random variables following standard normal distribution, and

$$Z = 1 + X + XY^{2}$$

$$W = 1 + X$$

Find
$$Cov(Z, W)$$
.
Solution

Find
$$Cov(2, W)$$
.

Solution
$$Cov(2, W) = Cov(1+x+xy^2, 1+x) = Cov(x+xy, x)$$

$$= Cov(x,x) + Cov(x,x) + (ov(x,x), x) (Roop 6)$$


$$= Vov(x) + E(xy^2) - E(xy^2) E(x) + (Sinu x,y ind x)$$

$$= 1 + E(xy^2) - E(xy^2) E(yy^2)$$

$$= 1 + E(xy^2) E(yy^2) + E(yy^2) E(yy^2)$$

$$= 1 + E(xy^2) E(yy^2) + E(yy^2)$$

$$= 1 + E(x$$



Example (Variance of a Sum)

Let Z = X + Y, then

Example (Variance of a Sum)

Let Z = X + Y, then

$$\mathsf{Var}(\mathbf{Z}) = \mathsf{Var}(\mathbf{X}) + \mathsf{Var}(\mathbf{Y}) + 2\mathsf{Cov}(\mathbf{X},\mathbf{Y})$$

Example (Variance of a Sum)

Let Z = X + Y, then

$$\mathsf{Var}(\mathbf{Z}) = \mathsf{Var}(\mathbf{X}) + \mathsf{Var}(\mathbf{Y}) + 2\mathsf{Cov}(\mathbf{X},\mathbf{Y})$$

More generally, for $a,b\in\mathbb{R},$ and Z=aX+bY, we have

Example (Variance of a Sum)
Let
$$Z = X + Y$$
, then

$$Var(Z) = Var(X) + Var(Y) + 2Cov(X, Y)$$

More generally, for $a, b \in \mathbb{R}$, and $Z = aX + bY$, we have

$$Var(Z) = a^2 Var(X) + b^2 Var(Y) + 2abCov(X, Y)$$

Solution

$$Var(Z) = a^2 Var(X) + b^2 Var(Y) + 2abCov(X, Y)$$

$$= Cov(X, Y) + Cov(X, Y) + Cov(X, Y) + Cov(Y, Y) + Cov(Y, Y)$$

$$= Var(X) + Cov(X, Y) + Cov(Y, Y) + Var(Y)$$

$$= Var(X) + Cov(X, Y) + Var(Y) + Var(Y)$$

Correlation coefficient

The correlation coefficient ρ_{XY} or $\rho(X, Y)$ is defined as follows

Correlation coefficient

The correlation coefficient ρ_{XY} or $\rho(X, Y)$ is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}} = \frac{\text{Cov}(X, Y)}{\sqrt{\rho_X \rho_Y}}$$

Correlation coefficient

The correlation coefficient ρ_{XY} or $\rho(X, Y)$ is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X\,\rho_Y}$$

* Given two RVs X and Y, define standardized versions of these as follows

Correlation coefficient

The correlation coefficient ρ_{XY} or $\rho(X, Y)$ is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X\,\rho_Y}$$

* Given two RVs X and Y, define standardized versions of these as follows

$$U = \frac{X - E[X]}{\rho_X}, \quad V = \frac{Y - E[Y]}{\rho_Y}$$

Correlation coefficient

The correlation coefficient ρ_{XY} or $\rho(X, Y)$ is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X\,\rho_Y}$$

* Given two RVs X and Y, define standardized versions of these as follows

$$U = \frac{X - E[X]}{\rho_X}, \quad V = \frac{Y - E[Y]}{\rho_Y}$$

* With this by computing the covariance Cov(U,V), we have

Correlation coefficient

The correlation coefficient ρ_{XY} or $\rho(X, Y)$ is defined as follows


$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X\,\rho_Y}$$

st Given two RVs X and Y, define standardized versions of these as follows

$$U = \frac{X - E[X]}{\rho_X}, \quad V = \frac{Y - E[Y]}{\rho_Y}$$

* With this by computing the covariance Cov(U, V), we have

$$\begin{array}{c}
\rho_{XY} = \text{Cov}(U, V) = \text{Cov}\left(\frac{X - E[X]}{\rho_X}, \frac{Y - E[Y]}{\rho_X}\right) \\
= \text{Cov}\left(\frac{X}{\rho_X}, \frac{Y}{\rho_Y}\right) = \frac{\text{Cov}(X, Y)}{\rho_X \rho_Y} = \frac{\text{Cov}(X, Y)}$$



Properties of correlation coefficient

Let X, Y be two RVs.

Properties of correlation coefficient

Properties of correlation coefficient

1.
$$-1 \le \rho(X, Y) \le 1$$

Properties of correlation coefficient

- 1. $-1 \le \rho(X, Y) \le 1$
- 2. If $\rho(X, Y) = 1$,

Properties of correlation coefficient

- 1. $-1 \le \rho(X, Y) \le 1$
- 2. If $\rho(X, Y) = 1$, then Y = aX + b, where a > 0

Properties of correlation coefficient

1.
$$-1 \le \rho(X, Y) \le 1$$

2. If
$$\rho(X, Y) = 1$$
, then $Y = aX + b$, where $a > 0$

3. If
$$\rho(X, Y) = 1$$
,

Properties of correlation coefficient

1.
$$-1 \le \rho(X, Y) \le 1$$

2. If
$$\rho(X, Y) = 1$$
, then $Y = aX + b$, where $a > 0$

3. If
$$\rho(X, Y) = 1$$
, then $Y = aX + b$, where $a < 0$

Properties of correlation coefficient

$$1/-1 \le \rho(X, Y) \le 1$$

If
$$\rho(X, Y) = 1$$
, then $Y = aX + b$, where $a > 0$

If
$$\rho(X, Y) = 1$$
, then $Y = aX + b$, where $a < 0$

$$\rho(aX+b,cY+d)=\rho(X,Y) \text{ for } a,c>0$$





8/1

» Positive Correlation, Negative Correlation, Uncorrelation...

Definition of positive, negative correlation

Let X and Y be two RVs.

Definition of positive, negative correlation

Let X and Y be two RVs.

1. If $\rho(X, Y) = 0$, we say that X and Y are uncorrelated

Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If $\rho(X, Y) = 0$, we say that X and Y are uncorrelated
- 2. If $\rho(\mathbf{X},\mathbf{Y})>0$, we say that \mathbf{X} and \mathbf{Y} are positively correlated

Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If $\rho(X, Y) = 0$, we say that X and Y are uncorrelated
- 2. If $\rho(\mathbf{X},\mathbf{Y})>0$, we say that \mathbf{X} and \mathbf{Y} are positively correlated
- 3. If $\rho(\mathbf{X},\mathbf{Y})<0$, we say that \mathbf{X} and \mathbf{Y} are negatively correlated

Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If $\rho(X, Y) = 0$, we say that X and Y are uncorrelated
- 2. If $\rho(X, Y) > 0$, we say that X and Y are positively correlated
- 3. If $\rho(X,Y) < 0$, we say that X and Y are negatively correlated

Pairwise uncorrelation and Variance

If X and Y are uncorrelated, then

$$Var(X + Y) = Var(X) + Var(Y)$$

More generally, if X_1, X_2, \dots, X_n , are pairwise uncorrelated, i.e., $\rho(X_i, X_j) = 0$ when $i \neq j$, then

Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If $\rho(X, Y) = 0$, we say that X and Y are uncorrelated
- 2. If $\rho(X, Y) > 0$, we say that X and Y are positively correlated
- 3. If ho(X,Y) < 0, we say that X and Y are negatively correlated

Pairwise uncorrelation and Variance

If X and Y are uncorrelated, then

$$Var(X + Y) = Var(X) + Var(Y)$$

More generally, if X_1, X_2, \dots, X_n , are pairwise uncorrelated, i.e., $\rho(X_i, X_j) = 0$ when $i \neq j$, then

$$Var(X_1 + X_2 + \cdots + X_n) = Var(X_1) + Var(X_2) + \cdots + Var(X_n)$$