# Part I

Abstract Algebra

## Definition (Binary Operations)

- lack A binary operation on a set A is an everywhere defined function

## Note: A binary operation must satisfy

- f assigns an element f(a,b) of A to each ordered pair (a,b) in  $A\times A.$ 
  - Since a binary operation is a function, only one element of A is assigned to each ordered pair.

## Definition (Binary Operations)

- lack A binary operation on a set A is an everywhere defined function

## Note: A binary operation must satisfy

- ▶ f assigns an element f(a,b) of A to each ordered pair (a,b) in  $A \times A$ .
- Since a binary operation is a function, only one element of A is assigned to each ordered pair.

#### Notation

- It's customary to denote binary operations by a symbol such as
  - a \* b, instead of \*(a,b)
    - a \* b, instead of \*(a)
    - T a \* b: the product of a and
  - ▶ A is closed under the operation \*, if a and b are elements in A,  $a*b \in A$ .

#### Notation

- ♪ It's customary to denote binary operations by a symbol such as
  - \* a\*b, instead of \*(a,b)
    - \*: multiplication
    - $\bullet$  a\*b: the product of a and b
- ▶ A is closed under the operation \*, if a and b are elements in A,  $a*b \in A$ .

#### Notation

- ♪ It's customary to denote binary operations by a symbol such as
  - \*
    - a\*b, instead of \*(a,b)→ \*: multiplication
    - $\bullet a * b$ : the product of a and b
- ▶ A is closed under the operation \*, if a and b are elements in A,  $a*b \in A$ .

## Example

- ▶ Let  $A = \mathbb{Z}$ . Define a \* b as a + b.
  - $\bullet$  \* is a binary operation on  $\mathbb{Z}$ .

### Example

- ▶ Let  $A = \mathbb{R}$ . Define a \* b as a/b.
  - $\mathfrak{I}$  \* is not a binary operation, since it is not defined for every ordered pair of elements of  $\mathbb{R}$ .
  - $\blacksquare$  For example, 3\*0 is not defined, since we can not divide by zero.

## Example

- ♪ Let  $A = \mathbb{Z}$ . Define a \* b as a + b.
  - $\bullet$  \* is a binary operation on  $\mathbb{Z}$ .

## Example

- ▶ Let  $A = \mathbb{R}$ . Define a \* b as a/b.
  - $\mathfrak{I}_{*}$  is not a binary operation, since it is not defined for every ordered pair of elements of  $\mathbb{R}$ .

## Definition

▶ If  $A = \{a_1, a_2, ..., a_n\}$  is a finite set, a binary operation on A can be defined by means of a multiplication table.

*	$ a_1 $	$a_2$	 $a_{j}$	 $a_n$
$a_1$				
$a_2$				
:				
$a_i$			$a_i * a_j$	
:				
$a_n$				

## Example ( $\vee$ and $\wedge$ )

♪ Let

$$A = \{0, 1\}$$

$$\begin{array}{c|cccc} \lor & 0 & 1 \\ \hline 0 & 0 & 1 \\ 1 & 1 & 1 \\ \end{array}$$

## Example (How Many Operations Can Be Defined on A?)

- ♪ Let
  - $A = \{a, b\}$
- ightharpoonup How many binary operations can be defined on A.
  - $\blacksquare$  Every binary operation \* on A can be described by a table

♣ Then what?

## Example (How Many Operations Can Be Defined on A?)

- ♪ Let
  - $A = \{a, b\}$
- ightharpoonup How many binary operations can be defined on A.
  - $lack {f P}$  Every binary operation \* on A can be described by a table

$$\begin{array}{c|cccc} * & a & b \\ \hline a & & \\ b & & \end{array}$$

■ Then what?

## Example (How Many Operations Can Be Defined on A?)

- ♪ Let
  - $A = \{a, b\}$
- ▶ How many binary operations can be defined on A.
  - $lack {f P}$  Every binary operation \* on A can be described by a table

$$\begin{array}{c|cccc} * & a & b \\ \hline a & & \\ b & & \end{array}$$

♪ Then what?

## Definition (Properties of Binary Operations)

- ightharpoonup For all elements a, b, and c in A
  - lacktriangle Commutative a\*b=b\*a
  - Associative a \* (b \* c) = (a \* b) \* c

## Definition (Properties of Binary Operations)

- ightharpoonup For all elements a, b, and c in A
  - $\bullet$  Commutative a\*b=b\*a
  - lacktriangle Associative a\*(b\*c)=(a\*b)\*c

## Definition (Properties of Binary Operations)

- ightharpoonup For all elements a, b, and c in A
  - $\bullet$  Commutative a \* b = b \* a
  - Associative a \* (b \* c) = (a \* b) \* c
  - Idempotent a \* a = a

## Definition (Identity)

- ▶ An element e in A is called an identity element if  $\forall a \in A$ 
  - $\bullet * a = a * e = a$

#### Note:

♪ An identity element must be unique.

### Definition (Inverse)

- ▶ An element  $a' \in A$  is called an inverse of a and written as  $a^{-1}$  if
  - a \* a' = a' \* a = e, or
  - $a * a^{-1} = a^{-1} * a = e$

## Definition (Identity)

- ▶ An element e in A is called an identity element if  $\forall a \in A$ 
  - $\bullet * a = a * e = a$

#### Note:

An identity element must be unique.

### Definition (Inverse)

- ▶ An element  $a' \in A$  is called an inverse of a and written as  $a^{-1}$  if
  - a \* a' = a' \* a = e, or
  - $a * a^{-1} = a^{-1} * a = e$

## Definition (Identity)

- ▶ An element e in A is called an identity element if  $\forall a \in A$ 
  - $\bullet * a = a * e = a$

#### Note:

An identity element must be unique.

## Definition (Inverse)

- ▶ An element  $a' \in A$  is called an inverse of a and written as  $a^{-1}$  if
  - a \* a' = a' \* a = e, or
  - $a * a^{-1} = a^{-1} * a = e$

#### **Theorem**

- Let \* be a binary operation on a set A, and suppose that \* satisfies the following properties for any a, b, and c in A:
  - a\*a=a
  - a\*b=b\*a
  - a\*(b\*c) = (a\*b)\*c
- ▶ Define a relation  $\leq$  on A by
  - $a \leq b$  if and only if a = a \* b
- ▶ Then  $(A, \leq)$  is a poset, and  $\forall a, b \in A$ , GLB(a, b) = a \* b.

### Proof.

- We must show that
  - ightharpoonup  $lap{1}{4}$   $lap{1}{4}$  is reflexive, antisymmetric and transitive.
  - $\bullet$   $a*b=a \wedge b$  for all a and b in A.



- ightharpoonup Since a = a \* a
  - $\bullet$   $a \leqslant a$  for all a in A
    - $\bullet$   $\triangleleft$   $\triangleleft$  is reflexive.



Proof: ≤ is reflexive

- ightharpoonup Since a = a \* a
  - $a \leq a$  for all a in A
  - is reflexive.



- Proof. Def  $a \leqslant b$  if and only if a = a \* b.
  - Now suppose that
    - $a \leqslant b$  and  $b \leqslant a$ 
      - a = a \* b = b \* a = b, by definition and property 2
      - a = b
    - **↑** Thus  $\leq$  is antisymmetric.



- Proof. Def  $a \leqslant b$  if and only if a = a \* b.
  - Now suppose that
    - $a \leqslant b \text{ and } b \leqslant a$
    - a = a \* b = b \* a = b, by definition and property 2
    - $\bullet$  so a=b
    - **.** Thus ≤ is antisymmetric.



- Proof. Def  $a \leqslant b$  if and only if a = a \* b.
  - Now suppose that
    - $a \leqslant b \text{ and } b \leqslant a$
    - a = a \* b = b \* a = b, by definition and property 2
    - so a=b
    - **.** Thus ≤ is antisymmetric.



- ♪ Now suppose that
  - $a \leqslant b \text{ and } b \leqslant a$
  - a = a \* b = b \* a = b, by definition and property 2
  - $\bullet$  so a=b
  - Thus  $\leq$  is antisymmetric.



Proof: 

≪ is transitive

- ♪ If  $a \leqslant b$  and  $b \leqslant c$ 
  - then a = a \* b = a \* (b \* c) = (a \* b) \* c = a \* c,
  - $\bullet$  so  $a \leqslant c$
  - √ ≤ is transitive.



Proof: ≤ is transitive

- ♪ If  $a \leqslant b$  and  $b \leqslant c$ 
  - 1 then a = a \* b = a \* (b \* c) = (a \* b) \* c = a \* c,
  - ightharpoonup so  $a\leqslant c$



#### Proof: ≤ is transitive

- ♪ If  $a \leqslant b$  and  $b \leqslant c$ 
  - 1 then a = a \* b = a \* (b \* c) = (a \* b) \* c = a \* c,
  - $\bullet$  so  $a \leqslant c$



- $\bullet$  a\*b is a lower bound for a and b
  - a \* b = a \* (b \* b) = (a \* b) \* b
    - $a * b \leq b$
  - **♪** similarly, a \* b ≤ a
  - a\*b is a lower bound for a and b
- - c = c \* a and c = c \* b by definition
  - c = (c \* a) \* b = c \* (a \* b)
  - $\bullet$  So,  $c \leqslant a * b$
  - ↑ Therefore, a \* b is the greatest lower bound of a and b.



- $\bullet$  a\*b is a lower bound for a and b
  - a \* b = a \* (b \* b) = (a \* b) \* b
  - so  $a*b \leqslant b$
  - **♪** similarly, a \* b ≤ a
  - a \* b is a lower bound for a and b
- - c = c \* a and c = c \* b by definition
  - c = (c \* a) \* b = c \* (a \* b)
  - $\bullet$  So,  $c \leqslant a * b$
  - ▶ Therefore, a \* b is the greatest lower bound of a and b.



- $\bullet$  a\*b is a lower bound for a and b
  - a \* b = a \* (b \* b) = (a \* b) \* b

  - ightharpoonup similarly,  $a*b \leqslant a$
  - a\*b is a lower bound for a and b
- 2 If  $c \leqslant a$  and  $c \leqslant b$ , then  $c \leqslant a * b$ 
  - c = c \* a and c = c \* b by definition
  - c = (c \* a) \* b = c \* (a \* b)
  - $\bullet$  So,  $c \leqslant a * b$
  - ▶ Therefore, a \* b is the greatest lower bound of a and b.



- $\bullet$  a\*b is a lower bound for a and b
  - a \* b = a \* (b \* b) = (a \* b) \* b
  - $\bullet$  so  $a*b \leqslant b$
  - $\bullet$  similarly,  $a * b \leq a$
  - $\bullet$  a\*b is a lower bound for a and b
- 2 If  $c \leqslant a$  and  $c \leqslant b$ , then  $c \leqslant a * b$ 
  - c = c \* a and c = c \* b by definition
  - c = (c \* a) \* b = c \* (a \* b)
  - $\bullet$  So,  $c \leqslant a * b$
  - ↑ Therefore, a \* b is the greatest lower bound of a and b.



- $\bullet$  a\*b is a lower bound for a and b
  - a \* b = a \* (b \* b) = (a \* b) \* b
  - $\bullet$  so  $a*b \leqslant b$
  - $\bullet$  similarly,  $a*b \leqslant a$
  - $\bullet$  a\*b is a lower bound for a and b
- 2 If  $c \leqslant a$  and  $c \leqslant b$ , then  $c \leqslant a * b$ 
  - c = c \* a and c = c \* b by definition
    - c = (c \* a) \* b = c \* (a \* b)
    - $\bullet$  So,  $c \leqslant a * b$
- ↑ Therefore, a \* b is the greatest lower bound of a and b.



- $\bullet$  a\*b is a lower bound for a and b
  - a \* b = a \* (b \* b) = (a \* b) \* b
  - $\bullet$  so  $a*b \leqslant b$
  - **♪** similarly, a \* b ≤ a
  - $\bullet$  a\*b is a lower bound for a and b
- 2 If  $c \leqslant a$  and  $c \leqslant b$ , then  $c \leqslant a * b$ 
  - c = c \* a and c = c \* b by definition
  - c = (c \* a) \* b = c \* (a \* b)

  - ↑ Therefore, a \* b is the greatest lower bound of a and b.



- $\bullet$  a\*b is a lower bound for a and b
  - a \* b = a \* (b \* b) = (a \* b) \* b
  - $\bullet$  so  $a*b \leq b$
  - $\bullet$  similarly,  $a*b \leqslant a$
  - $\bullet$  a\*b is a lower bound for a and b
- 2 If  $c \leqslant a$  and  $c \leqslant b$ , then  $c \leqslant a * b$ 
  - c = c \* a and c = c \* b by definition
  - c = (c \* a) \* b = c \* (a \* b)
  - ightharpoonup So,  $c \leqslant a * b$
  - ▶ Therefore, a \* b is the greatest lower bound of a and b.

