

1 Lecture 23

Slide 1

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Spatial and temporal coherence Optics and Laser Physics T. Cesca

Spatial and temporal coherence are independent concepts: a wave can have perfect spatial coherence but partial temporal coherence, and viceversa.

Spatial coherence is associated with the wave's properties **transverse** to the direction of propagation.

Spatial coherence is associated with a **distribution of propagation vectors** \vec{k} associated with the wave, i.e., with a departure of the wave from the ideal plane wave.

Temporal coherence is associated with the wave's properties **along** the direction of propagation.

It is associated with the **frequency distribution** of the source.

$$\tau_c \Delta\nu \approx 1 \Rightarrow \tau_c \approx \frac{1}{\Delta\nu} = \frac{\lambda^2}{c\Delta\lambda}$$

474

Temporal coherence is more related to monochromaticity.

Slide 2

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA First-order auto-correlation function Optics and Laser Physics T. Cesca

Degree of first-order longitudinal (temporal) coherence

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

Let's consider a **quasi-monochromatic** em wave, with central frequency ω_0 and bandwidth $\Delta\omega$:

$$E(t) = E_0 e^{-i(\omega_0 t + \phi(t))}$$

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle} = \frac{\langle E_0 e^{i(\omega_0 t + \phi(t))} E_0 e^{-i(\omega_0(t+\tau) + \phi(t+\tau))} \rangle}{E_0^2}$$

$$= \underbrace{e^{-i\omega_0\tau}}_{\text{real part}} \langle e^{-i[\phi(t+\tau) - \phi(t)]} \rangle$$

The real part of $g^{(1)}(\tau)$ is an oscillating function with period $\frac{2\pi}{\omega_0}$

475

The **first-order auto-correlation function** gives information on temporal coherence.

T is the period of oscillation of our wave.

The real part of $g^{(1)}$ is an oscillating function! The amplitude of this function will be modulated by the term in the brackets.

Slide 3

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA First-order auto-correlation function Optics and Laser Physics T. Cesca

Degree of first-order longitudinal (temporal) coherence

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

The variation of the modulus of $g^{(1)}(\tau)$ contains information on the coherence of the light:

$$|g^{(1)}(\tau)| = |\langle e^{-i[\phi(t+\tau) - \phi(t)]} \rangle|$$

$$|g^{(1)}(0)| = 1 \quad \text{always}$$

The real part of $g^{(1)}(\tau)$ is an oscillating function with period $\frac{2\pi}{\omega_0}$

476

τ_c is the correlation time.

Slide 4

800 ANNI UNIVERSITÀ DELL'ANNO DI PADOVA

First-order auto-correlation function

Optics and Laser Physics
T. Cesca

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

$0 < \tau \ll \tau_c \Rightarrow \phi(t+\tau) \approx \phi(t)$

$\Rightarrow |g^{(1)}(\tau)| \cong 1$

If τ increases, $|g^{(1)}(\tau)|$ decreases since it increases the probability of the phase relation to become random.

$\tau \gg \tau_c \Rightarrow \phi(t+\tau), \phi(t)$ become totally uncorrelated

$\Rightarrow |g^{(1)}(\tau)| = |\langle e^{-i[\phi(t+\tau)-\phi(t)]} \rangle| = 0$ on average

Degree of first-order **longitudinal (temporal)** coherence

477

If τ is increasing the relation between the phases become random.

Slide 5

800 ANNI UNIVERSITÀ DELL'ANNO DI PADOVA

First-order auto-correlation function

Optics and Laser Physics
T. Cesca

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

$\Rightarrow |g^{(1)}(\tau)|$ varies from 1 → 0 on a time scale of the order of τ_c

The functional shape of $g^{(1)}(\tau)$ for partially coherent light depends on the spectral broadening

Lorentzian lineshape (homogeneous broadening)

$$g^{(1)}(\tau) = e^{-i\omega_0\tau} e^{-|\tau|/\tau_c} \quad \Delta\omega_0 = 2\pi\Delta\nu_0 \quad \tau_c = \frac{2\pi}{\Delta\omega_0} = \frac{1}{\Delta\nu_0}$$

Degree of first-order **longitudinal (temporal)** coherence

478

The shape of $g^{(1)}$ depends on the spectral broadening. If we have homogeneous broadening, we have that the first-order correlation function can be calculated in this way.

Slide 6

800 ANNI UNIVERSITÀ DELL'ANNO DI PADOVA

First-order auto-correlation function

Optics and Laser Physics
T. Cesca

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

Gaussian lineshape (inhomogeneous broadening)

$$g^{(1)}(\tau) = e^{-i\omega_0\tau} e^{-(\pi/2)(\tau/\tau_c)^2}$$

$$\tau_c = \frac{2\sqrt{2\ln 2}}{\Delta\omega_0}$$

Lorentzian lineshape (homogeneous broadening)

$$g^{(1)}(\tau) = e^{-i\omega_0\tau} e^{-|\tau|/\tau_c} \quad \Delta\omega_0 = 2\pi\Delta\nu_0 \quad \tau_c = \frac{2\pi}{\Delta\omega_0} = \frac{1}{\Delta\nu_0}$$

Degree of first-order **longitudinal (temporal)** coherence

If we have inhomogeneous broadening, we have a function like this.

479

Slide 7

UNIVERSITÀ
DEGLI STUDI
DI PADOVA

First-order auto-correlation function

Optics and Laser Physics
T. Cesca

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

Degree of first-order **longitudinal (temporal)** coherence

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

From the experimental point of view the first-order auto-correlation function $g^{(1)}(\tau)$ can be measured by using a **Michelson's interferometer**

480

Slide 8

UNIVERSITÀ
DEGLI STUDI
DI PADOVA

Michelson's interferometer

Optics and Laser Physics
T. Cesca

$k = \frac{2\pi}{\lambda_0}$

$\Delta L = L_2 - L_1$

$E_{out} = E_1 + E_2$

$$\begin{aligned} &= \frac{E_0}{2} e^{i2kL_1} + \frac{E_0}{2} e^{i2kL_2} e^{i\Delta\phi} \\ &= \frac{E_0}{2} e^{i2kL_1} (1 + e^{i2k\Delta L} e^{i\Delta\phi}) \\ &= \frac{E_0}{2} e^{i2kL_1} (1 + e^{i\delta}) \end{aligned}$$

$\delta = \frac{4\pi}{\lambda_0} \Delta L + \Delta\phi$

$I_{out} \propto |E_{out}|^2$

$I_{out} = I_0^2 \cos^2 \left(\frac{\delta}{2} \right)$

$\cos\delta = 1 - 2 \left(\sin \frac{\delta}{2} \right)^2$

481

Slide 9

UNIVERSITÀ
DEGLI STUDI
DI PADOVA

Michelson's interferometer

Optics and Laser Physics
T. Cesca

$k = \frac{2\pi}{\lambda_0}$

$\Delta L = L_2 - L_1$

$I_{out} = I_0^2 \cos^2 \left(\frac{\delta}{2} \right)$

Interference **maxima**

$\delta = \frac{4\pi}{\lambda_0} \Delta L + \Delta\phi = 2m\pi$

Interference **minima**

$\delta = \frac{4\pi}{\lambda_0} \Delta L + \Delta\phi = (2m+1)\pi$

Fringe visibility

$$V = \frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}}$$

482

The problem is how to obtain τ_c from an experimental point of view. To determine the auto-correlation function we can use a Michelson interferometer.

It can be schematized in this way. We send the input beam on a beam splitter which divide the amplitude of the beam in 50%. We have one mirror in each arm. We can vary the distance of the second mirror.

The output will produce fringe of interference. If we move the second mirror, the fringes are moving. By analyzing the evolution of the fringes as a function of L_2 we can characterize the beam.

$\Delta\Phi$ is an extra phase which take into account the phase-shift introduced by the path or any other possibility (for instance reflection at the beam splitter or the presence of coatings and so on...).

The intensity can be described by this cosin function of δ . We apply the condition to get maxima of interference and minima.

We can introduce also the **fringe visibility**.

Slide 10

800 ANNI UNIVERSITÀ DELL'ANNO DI PADOVA

First-order auto-correlation function

Optics and
Laser Physics
T. Cesca

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

...using a Michelson's interferometer

$$E_{out}(t) = E(t) + E(t+\tau)$$

$$I(\tau) \propto \langle E_{out}^*(t)E_{out}(t) \rangle =$$

$$= \langle E^*(t)E(t) \rangle + \langle E^*(t+\tau)E(t+\tau) \rangle +$$

$$+ \langle E^*(t)E(t+\tau) \rangle + \langle E(t)E^*(t+\tau) \rangle$$

$$= 2\langle E^*(t)E(t) \rangle + 2Re(\langle E^*(t+\tau)E(t+\tau) \rangle)$$

$$= 2\langle E^*(t)E(t) \rangle [1 + Re(g^{(1)}(\tau))]$$

Degree of first-order **longitudinal (temporal)** coherence

$$\tau = \frac{2\Delta L}{c}$$

483

Let us see how we can link the Michelson interferometer to compute the auto-correlation function.

The intensity that we can measure as a function of τ is related to the first-order auto-correlation function.

Slide 11

800 ANNI UNIVERSITÀ DELL'ANNO DI PADOVA

First-order auto-correlation function

Optics and
Laser Physics
T. Cesca

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle |E(t)|^2 \rangle}$$

$$\langle E^*(t)E(t+\tau) \rangle = \frac{1}{T} \int_T E^*(t)E(t+\tau) dt$$

...using a Michelson's interferometer

$$E_{out}(t) = E(t) + E(t+\tau)$$

$$I(\tau) = I_0 [1 + Re(g^{(1)}(\tau))]$$

$$I_{max,min} = I_0(1 \pm |g^{(1)}(\tau)|)$$

$$V = \frac{I_{max} - I_{min}}{I_{max} + I_{min}} = |g^{(1)}(\tau)|$$

Degree of first-order **longitudinal (temporal)** coherence

$$\tau = \frac{2\Delta L}{c}$$

484

We can relate the fringe visibility with the first-order auto-correlation function.

By changing L_2 , we change τ so we can compute $|g_1(\tau)|$.

Slide 12

800 ANNI UNIVERSITÀ DELL'ANNO DI PADOVA

First-order auto-correlation function

Optics and
Laser Physics
T. Cesca

$$g_{12}^{(1)}(\tau) = \frac{\langle E_1(t)^*E_2(t+\tau) \rangle}{\sqrt{\langle |E_1(t)|^2 \rangle \langle |E_2(t)|^2 \rangle}}$$

If the amplitudes in the two arms are different:

$$E_{out}(t) = E_1(t) + E_2(t+\tau)$$

$$I(\tau) = I_1 + I_2 + 2\sqrt{I_1 I_2} Re(g_{12}^{(1)}(\tau))$$

$$V = \frac{2\sqrt{I_1 I_2}}{I_1 + I_2} |g_{12}^{(1)}(\tau)|$$

Degree of first-order **longitudinal (temporal)** coherence

$$\tau = \frac{2\Delta L}{c}$$

485

If the amplitudes of the electric field are different we have this more complex formula.

Slide 13

Temporal coherence of light		Optics and Laser Physics T. Cesca
Perfectly monochromatic light	<ul style="list-style-type: none"> • $\Delta\omega = 0$ • Perfect temporal coherence • Coherence time: $\tau_c = \infty$ • $g^{(1)}(\tau) = 1$ 	
Chaotic light	<ul style="list-style-type: none"> • $\Delta\omega \neq 0$ (finite bandwidth) • Partial temporal coherence • Coherence time: $\tau_c \propto \frac{1}{\Delta\omega}$ • $0 < g^{(1)}(\tau) < 1$ 	In this way we can distinguish the property of our light beam. The most real case is chaotic light or incoherent light .
Incoherent light	<ul style="list-style-type: none"> • $\Delta\omega = \infty$ • No temporal coherence • Coherence time: $\tau_c = 0$ • $g^{(1)}(\tau) = 0$ 	


487

Slide 14

Temporal coherence of light						Optics and Laser Physics T. Cesca
Source	Wavelength (nm)	Frequency (10 ¹⁴ Hz)	Bandwidth (Δν)	Coherence time, τ _c (s)	Coherence length, L _c	
white light	400-600	5-7.5	2.5 × 10 ¹⁴ Hz	4 × 10 ⁻¹⁵	1.2 μm (a few λ)	
Hg lamp with single isotope	546.1	5.49	300 MHz	3 × 10 ⁻⁹	1 m	
He-Ne laser	632.8	4.74	1 GHz	10 ⁻⁹	0.3 m	
Stabilized He-Ne laser	632.8	4.74	10 kHz	10 ⁻⁴	30 km	

488

Slide 15



489

In the same way, we can define a **first-order cross-correlation function** for the spatial coherence. The only difference is that we have to evaluate the electric field at the position x_1 and x_2 .

This is the same way for a Young interference. Interference between two waves that are passing in the positions x_1 and x_2 .

Slide 16

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA First-order cross-correlation function Optics and Laser Physics T. Cesca

Degree of first-order **transverse (spatial)** coherence

$$g_{12}^{(1)}(\tau) \equiv g_{12}^{(1)}(x_1, x_2, \tau) = \frac{\langle E(x_1, t)^* E(x_2, t + \tau) \rangle}{\sqrt{\langle |E(x_1, t)|^2 \rangle \langle |E(x_2, t)|^2 \rangle}}$$

$$\langle E(x_1, t)^* E(x_2, t + \tau) \rangle = \frac{1}{T} \int_T E(x_1, t)^* E(x_2, t + \tau) dt$$

$|g_{12}^{(1)}(\tau = 0)| = 1$ Full coherence
 $0 < |g_{12}^{(1)}(\tau = 0)| < 1$ Partial coherence
 $|g_{12}^{(1)}(\tau = 0)| = 0$ Incoherence

between the wavetrains passing in x_1 and x_2

490

Slide 17

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

Degree of second-order **(temporal)** coherence

$$g^{(2)}(\tau) = \frac{\langle E^*(t) E(t + \tau) E(t + \tau) E(t) \rangle}{\langle E^*(t) E(t) \rangle \langle E^*(t + \tau) E(t + \tau) \rangle} = \frac{\langle I(t) I(t + \tau) \rangle}{\langle I(t) \rangle \langle I(t + \tau) \rangle}$$

Let's assume a source with constant average intensity: $\langle I(t) \rangle = \langle I(t + \tau) \rangle = \langle I \rangle$

$$I(t) = \langle I \rangle + \Delta I(t) \quad \text{with} \quad \langle \Delta I(t) \rangle = 0$$

$\tau \gg \tau_c$

$$\begin{aligned} \langle I(t) I(t + \tau) \rangle &= \\ &= \langle (\langle I \rangle + \Delta I(t)) (\langle I \rangle + \Delta I(t + \tau)) \rangle = \\ &= \langle I \rangle^2 + \cancel{\langle I \rangle \langle \Delta I(t) \rangle} + \cancel{\langle I \rangle \langle \Delta I(t + \tau) \rangle} + \cancel{\langle \Delta I(t) \rangle \langle \Delta I(t + \tau) \rangle} = \langle I \rangle^2 \\ &= 0 \quad = 0 \quad = 0 \quad \tau \gg \tau_c \end{aligned}$$

496

We can have also in this case **full coherence**, **partial coherence** and **incoherence**.

You have to calculate **at the same time** the cross-correlation function!

While for the temporal coherence was **at the same position**!

Now, let us look at the **second-order correlation function**. We obtain the information about the degree of second-order temporal coherence: we are looking at the second power of the electric field. So, we have considering the correlation between the intensity of the beam. How do the intensity are correlated at different time?

Let us assume a **constant average intensity**. So, we can write the intensity at each time as the average value and fluctuations around the average value.

If $\tau \gg \tau_c$ the fluctuations of the intensity are uncorrelated.

Slide 18

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

Degree of second-order **(temporal)** coherence

$$g^{(2)}(\tau) = \frac{\langle E^*(t) E(t + \tau) E(t + \tau) E(t) \rangle}{\langle E^*(t) E(t) \rangle \langle E^*(t + \tau) E(t + \tau) \rangle} = \frac{\langle I(t) I(t + \tau) \rangle}{\langle I(t) \rangle \langle I(t + \tau) \rangle}$$

Let's assume a source with constant average intensity: $\langle I(t) \rangle = \langle I(t + \tau) \rangle = \langle I \rangle$

$$I(t) = \langle I \rangle + \Delta I(t) \quad \text{with} \quad \langle \Delta I(t) \rangle = 0$$

$\tau \gg \tau_c$

$$\begin{aligned} \langle I(t) I(t + \tau) \rangle_{\tau \gg \tau_c} &= \langle I \rangle^2 \\ g^{(2)}(\tau \gg \tau_c) &= \frac{\langle I(t) I(t + \tau) \rangle}{\langle I(t) \rangle^2} = 1 \end{aligned}$$

If we calculate $g^{(2)}$ it is equal to one.

497

Slide 19

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

Degree of second-order (**temporal**) coherence

$$g^{(2)}(\tau) = \frac{\langle E^*(t)E^*(t+\tau)E(t+\tau)E(t) \rangle}{\langle E^*(t)E(t) \rangle \langle E^*(t+\tau)E(t+\tau) \rangle} = \frac{\langle I(t)I(t+\tau) \rangle}{\langle I(t) \rangle \langle I(t+\tau) \rangle}$$

Let's assume a source with constant average intensity: $\langle I(t) \rangle = \langle I(t+\tau) \rangle = \langle I \rangle$

$$I(t) = \langle I \rangle + \Delta I(t) \quad \text{with} \quad \langle \Delta I(t) \rangle = 0$$

$\tau \gg \tau_c$	$\tau \ll \tau_c$	The intensity fluctuations at time t and $t+\tau$ might be correlated
-------------------	-------------------	---

$$\langle I(t)I(t+\tau) \rangle_{\tau \gg \tau_c} = \langle I \rangle^2 \quad \rightarrow \quad \tau = 0$$

$$g^{(2)}(\tau \gg \tau_c) = \frac{\langle I(t)I(t+\tau) \rangle}{\langle I(t) \rangle^2} = 1 \quad g^{(2)}(\tau = 0) = \frac{\langle I(t)^2 \rangle}{\langle I(t) \rangle^2} \geq 1$$

498

Slide 20

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

Degree of second-order (**temporal**) coherence

$$\forall I(t) \quad g^{(2)}(0) \geq 1 \quad \text{and} \quad g^{(2)}(0) \geq g^{(2)}(\tau)$$

Perfectly coherent light: $g^{(2)}(\tau) = 1 \quad \forall \tau$

Partially coherent (**chaotic**) light:

Gaussian lineshape: $g^{(2)}(\tau) = 1 + e^{-\pi(\tau/\tau_c)^2}$

Lorentzian lineshape: $g^{(2)}(\tau) = 1 + e^{-2|\tau|/\tau_c}$

$$g^{(2)}(\tau) = 1 + |g^{(1)}(\tau)|^2$$

For every **classical** source of light!

499

Slide 21

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

COHERENT light: $g^{(2)}(0) = 1$	Perfectly coherent light	} classical
BUNCHED light: $g^{(2)}(0) > 1$	Partially coherent light	
ANTIBUNCHED light: $g^{(2)}(0) < 1$	Quantum nature of light!	

Bunched light: Photon bunches, Fewer photons

Coherent (random): Random distribution of photons over time

Bunched: Bunched distribution of photons

Antibunched: Regular distribution of photons over time

500

Now, let us consider the case in which $\tau \ll \tau_c$. If $\tau = 0$, we see that $g^{(2)} \geq 1$.

We have that $I(t)^2$ is always positive so we have that the numerator is larger than the denominator!

We can summarize it here.

The trend of the chaotic depends on the lineshape (homogeneous or inhomogeneous broadening).

In general the second-order correlation function is related to the first-order correlation function. This is **true for any classical source of light!**

We can use the second-order correlation function to distinguish between **coherent light** and **noncoherent light**. This is valid for a **classical** source of light.

We can have **antibunched light**. This is possible only for a **quantum** nature of light!

These are the distribution of number of photons over time (each black dot is a photon). Coherent light correspond to a distribution of random photons. Bunched light means that we have a bunched distribution of photons. For antibunched we have a regular distribution of photons over time, with a given distance between each other.

So, for second-order correlation function we can distinguish between classical and quantum light beam!

Slide 22

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

...in terms of photons: $g^{(2)}(\tau) = \frac{\langle n_1(t)n_2(t+\tau) \rangle}{\sqrt{\langle n_1(t) \rangle \langle n_2(t+\tau) \rangle}}$

$g^{(2)}(\tau)$ is proportional to the **conditional probability** of detecting a second photon at time $t = \tau$, given that we detected one at $t = 0$.

Hanbury-Brown & Twiss (HBT) configuration

50:50 beam splitter
Photons → D1
D2
Antibunched light
Flux of photons with long temporal intervals between two consecutive photons
 $g^{(2)}(0) \approx 0 \quad g^{(2)}(0) \leq g^{(2)}(\tau)$
in contrast with the results for classical light:
 $g^{(2)}(0) \geq 1 \quad g^{(2)}(0) \geq g^{(2)}(\tau)$

501

So, $g^{(2)}$ is 0 at $\tau = 0$ (we cannot have at the same time the photon reaching the second detector). But, if we wait long time the probability to have a photon in the second detector is increasing and so you have the stop event.

Slide 23

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

...in terms of photons: $g^{(2)}(\tau) = \frac{\langle n_1(t)n_2(t+\tau) \rangle}{\sqrt{\langle n_1(t) \rangle \langle n_2(t+\tau) \rangle}}$

$g^{(2)}(\tau)$ is proportional to the **conditional probability** of detecting a second photon at time $t = \tau$, given that we detected one at $t = 0$.

Hanbury-Brown & Twiss (HBT) configuration

50:50 beam splitter
Photons → D1
D2
Antibunched light
Time delay τ (ns)
 $g^{(2)}(\tau)$

502

We can compute the second-order correlation function in terms of photons. It represent the condition probability of detecting a second photon at time $t = \tau$, given that we detected one at $t = 0$.

It is possible realizing the second-order correlation function with the HBT configuration.

We have a beam of photons impinging on a 50:50 beam splitter. We have two detectors. We can evaluate how long after the start the stop occurs.

Let us send to this setup an **antibunched beam** (long temporal intervals between two consecutive photons).

We have 50:50 probability that the beam goes to the first or to the second detector.

Let us suppose that it arrives in D1, the counter starts. If we wait a lot of time we have a second photon which can reach the detector D2. The timer is stopped.

If we measure the second-order correlation function (how many stop event occur after a given start event and you create a histogram for time delay τ) we have this trend here.

Slide 24

800 ANNI UNIVERSITÀ DEGLI STUDI DI PADOVA Second-order correlation function Optics and Laser Physics T. Cesca

...in terms of photons: $g^{(2)}(\tau) = \frac{\langle n_1(t)n_2(t+\tau) \rangle}{\sqrt{\langle n_1(t) \rangle \langle n_2(t+\tau) \rangle}}$

$g^{(2)}(\tau)$ is proportional to the **conditional probability** of detecting a second photon at time $t = \tau$, given that we detected one at $t = 0$.


Hanbury-Brown & Twiss (HBT) configuration

50:50 beam splitter
Photons → D1
D2
Bunched light
Number of events
Time interval (arb. units)

503


For bunched light, we have a large probability if we detect an event of start an event after a very short delay. If you wait time, the probability is decreasing: out of the bunch we get no more photons.

Slide 25



504

Slide 26



505

The excited a Na beam with a laser.

To obtain antibunching light we need the emission from a single atom (so obtaining a single photon). If we have a single atom which is emitting with a radiative lifetime τ_R , the probability of getting the emission of two photons with $\tau \ll \tau_R$ is very low.

There is large interesting in realizing single photons sources. We need a parameter that we can control from the outside to trigger the emission of a single photon. Single-photon sources can be used in quantum information. The idea can be to have single-photon sources which are able to emit at room temperature and that can be triggered.