

Copyright

```
close all; format compact; clc;  
fprintf("Engineer: Rodrigo Becerril Ferreyra\n");  
fprintf("Company: California State University, Long Beach\n");  
fprintf("Project Name: Task 5\n");  
fprintf("Date: 20 October 2020\n");
```

```
Engineer: Rodrigo Becerril Ferreyra  
Company: California State University, Long Beach  
Project Name: Task 5  
Date: 20 October 2020
```

Problem 1

```
fprintf("\nProblem 1\n");
clear variables;

x = [0, 2, 5];
y = [0, -1, 5];
domain = linspace(0, 5, 101);
f = @(z) (z.^2 - 3.*z)./2; % See picture for work

figure(); hold on;
line = plot(domain, f(domain), 'r-');
points = plot(x, y, 'b.', "MarkerSize", 20);
title("Manual Interpolation Calculation");
legend([points, line], ["Data points", "Interpolation"], "Location",
"northwest");
text(3.1, 0, "$$\leftarrow f(z) = \frac{z^2 - 3z}{2}$$", "Interpreter",
"latex");
grid on;
hold off;

fprintf("The interpolating function is (z^2 - 3*z)/2.\n");
```


Problem 1

The interpolating function is $(z^2 - 3z)/2$.

i	x_i	y_i
1	(0,	0)
2	(2,	-1)
3	(5,	5)

$$f(z) = \sum_{i=1}^N y_i \prod_{\substack{j=1 \\ j \neq i}}^N \frac{z - x_j}{x_i - x_j}$$

$$\begin{aligned} f(z) &= -\frac{z(z-5)}{2(2-5)} + 5\frac{z(z-2)}{5(5-2)} \\ &= \frac{1}{6}(z^2 - 5z) + \frac{1}{3}(z^2 - 2z) \\ &= \frac{1}{6}z^2 - \frac{5}{6}z + \frac{1}{3}z^2 - \frac{2}{3}z \\ &= \frac{1}{2}z^2 - \frac{3}{2}z \\ f(z) &= \frac{z^2 - 3z}{2} \end{aligned}$$



Problem 2

```
fprintf("\nProblem 2\n");
clear variables;

% Given
x = 0:10;
y = x.^2 + 3.*x + 2;
sample_points = [1/2, 3/2, 5/2];

% See fencepost problem; we need N-1 lines
% Also we need to put function handles in a cell array (not a matrix)
lines = cell(1, length(x)-1);
for i = 1:(length(x)-1)

    % Lagrange (1st degree) polynomial
    lines{i} = @(z) (z - x(i+1))./(x(i) - x(i+1)) .* y(i) + (z -
x(i))./(x(i+1) - x(i)) .* y(i+1);

end

figure(); hold on;

% Plot all of the lines
for i = 1 : length(lines)

    domain = linspace(x(i), x(i+1), 11);
    l = plot(domain, lines{i}(domain), 'r-');

end
p = plot(x, y, 'b.', "MarkerSize", 20);
grid on;
title("Linear Interpolation Between Points");
legend([p, l], ["Data points", "Interpolating lines"], "Location",
"northwest");

hold off;

for i = 1:3

    experimental = lines{i}(sample_points(i));
    actual = sample_points(i).^2 + 3.*sample_points(i) + 2;
    fprintf("Relative error of approximating f(%3.1f): %f%%\n",
sample_points(i), 100*abs(experimental - actual)/actual);


end
```

Problem 2

Relative error of approximating $f(0.5)$: 6.666667%

Relative error of approximating $f(1.5)$: 2.857143%

Relative error of approximating $f(2.5)$: 1.587302%



Problem 3

```
fprintf("\nProblem 3\n");
clear variables;

c = polyfit([0, pi/2, pi], [0, 1, 0], 2);
fprintf("The polynomial that goes through the points (0, 0), (pi/2, 1), and
(pi, 0) is\n");
fprintf("f(x) = %fx^2 + %fx + %f\n", c(1), c(2), c(3));
fprintf("More precisely, f(x) = (-4/pi^2)x^2 + (4/pi)x; see attached image");

figure(); hold on;

domain = linspace(0, pi, 101);
line = plot(domain, polyval(c, domain), 'r-');
points = plot([0, pi/2, pi], [0, 1, 0], 'b.', "MarkerSize", 20);
legend([points, line], ["Data points", "Interpolation"], "Location", "south");
grid on;
title("Interpolation Using polyfit");
axis([0, pi, -0.2, 1.0625]);
set(gca, 'XTick', 0:pi/4:pi); set(gca, 'XTickLabel', {'0', '\pi/4', '\pi/2',
'3\pi/4', '\pi'});
text(pi/8, polyval(c, pi/8), "$\leftarrow f(z) = \frac{-4\left(z^2 - \pi
z\right)}{\pi^2}$", "Interpreter", "latex");

hold off;
```

Problem 3

The polynomial that goes through the points $(0, 0)$, $(\pi/2, 1)$, and $(\pi, 0)$ is $f(x) = -0.405285x^2 + 1.273240x + -0.000000$

More precisely, $f(x) = (-4/\pi^2)x^2 + (4/\pi)x$; see attached image


i	x_i	y_i
1	0,	0)
2	$(\pi/2,$	1)
3	$(\pi,$	0)

$$f(z) = \sum_{i=1}^N y_i \prod_{\substack{j=1 \\ j \neq i}}^N \frac{z - x_j}{x_i - x_j}$$

$$f(z) = \frac{(z-0)(z-\pi)}{(\pi/2-0)(\pi/2-\pi)}$$

$$= \frac{z^2 - \pi z}{-\pi^2/4}$$

$$f(z) = \frac{-4(z^2 - \pi z)}{\pi^2} = -\frac{4}{\pi^2}z^2 + \frac{4}{\pi}z + 0$$



Problem 4

```
fprintf("\nProblem 4\n");
clear variables;

figure(); hold on;
interpol = @(z) (-8.*z.^3 + 8*pi^2.*z)./(3*pi^3); % see attached image
domain = linspace(-pi/2, pi, 101);
lines = plot(domain, interpol(domain), 'r-');
points = plot([-pi/2, 0, pi/2, pi], [-1, 0, 1, 0], 'b.', "MarkerSize", 20);
set(gca, 'XTick', -pi/2:pi/4:pi); set(gca, 'XTickLabel', {'-\pi/2', '-\pi/4', '0',
'\pi/4', '\pi/2', '3\pi/4', '\pi'});
axis([-pi/2, pi, -interpol(pi/sqrt(3)), interpol(pi/sqrt(3))]);
legend([points, lines], ["Data points", "Interpolation"], "Location",
"southeast");
grid on;
title("Manual Interpolation Calculation")
text(-pi/4, interpol(-pi/4), "$$\leftarrow f(z) = \frac{-8z^3+8\pi^2z}{3\pi^3}$$", "Interpreter", "latex");

hold off;

fprintf("The interpolating function is f(z) = (-8z^3 + 8pi^2*z)/(3pi^3).\n");
```


Problem 4


The interpolating function is $f(z) = (-8z^3 + 8\pi^2 z)/(3\pi^3)$.

i	x_i	y_i
1	$(-\pi/2, -1)$	
2	$(0, 0)$	
3	$(\pi/2, 1)$	
4	$(\pi, 0)$	

$$f(z) = \sum_{i=1}^N y_i \prod_{\substack{j=1 \\ j \neq i}}^N \frac{z - x_j}{x_i - x_j}$$

$$\begin{aligned} f(z) &= -\frac{(z-0)(z-\pi/2)(z-\pi)}{(-\pi/2-0)(-\pi/2-\pi/2)(-\pi/2-\pi)} \\ &\quad + \frac{(z+\pi/2)(z-0)(z-\pi)}{(\pi/2+\pi/2)(\pi/2-0)(\pi/2-\pi)} \\ &= \frac{-8z^3 + 8\pi^2 z}{3\pi^3} \end{aligned}$$

(Reduced using a computer algebra system)



Problem 5

```

fprintf("\nProblem 5\n");
clear variables;

x = [-pi, -pi/2, 0, pi/2, pi];
y = [0, -1, 0, 1, 0];
figure(); hold on;


domain = -pi:pi/20:pi;
lines = plot(domain, spline(x, y, domain), 'r-');
points = plot(x, y, 'b.', "MarkerSize", 20);

grid on;
legend([points, lines], ["Data points", "Interpolation"], "Location",
"southeast");
title("Interpolation Using spline");
set(gca, 'XTick', -pi:pi/2:pi); set(gca, 'XTickLabel', {'-\pi', '-\pi/2', '0',
'\pi/2', '\pi'});

hold off;

```

Problem 5



Problem 6

```
fprintf("\nProblem 6\n");
clear variables;

x = 0 : 0.1 : 1;
y = [3.16, 3.01, 2.73, 2.47, 2.13, 1.82, 1.52, 1.21, 0.76, 0.43, 0.03];


ab = [sum(sin(x).^2), sum(sin(x).*cos(x)); sum(cos(x).*sin(x)),
sum(cos(x).^2)] \ [sum(y.*sin(x)); sum(y.*cos(x))];
a = ab(1); b = ab(2); clear ab;

figure(); hold on;

f = @(x) a.*sin(x) + b.*cos(x);
func = plot(x, f(x), 'r-');
points = plot(x, y, 'b.', "MarkerSize", 20);
grid on;
legend([points, func], ["Data points", "$$y=a\sin(x) + b\cos(x)$$"],
"Location", "southwest", "Interpreter", "latex")
text([0.025, 0.025], [1.5, 1], [sprintf("$$a = %f$$", a), sprintf("$$b = %f$$", b)], "Interpreter", "latex");
title("Line of Best Fit Approximation");

hold off;
```

Problem 6



Problem 7


```
fprintf("\nProblem 7\n");
clear variables;

x = [-pi, -pi/2, 0, pi/2, pi];
y = [0, -1, 0, 1, 0];
figure(); hold on;

domain = -pi:pi/20:pi;
s = spline(x, y, domain);
f = sin(domain);
spline_ = plot(domain, s, 'r-');
func = plot(domain, f, 'g-');
points = plot(x, y, 'b.', "MarkerSize", 20);
grid on;
legend([points, spline_, func], ["Data points", "Interpolation",
"$y=\sin(x)$"], "Location", "southeast", "Interpreter", "latex");
title("Interpolation Using spline Compared to Original Function");
set(gca, 'XTick', -pi:pi/2:pi); set(gca, 'XTickLabel', {'-\pi', '-\pi/2', '0',
'\pi/2', '\pi'});
hold off;
fprintf("The sum of the squared errors over the (discretely-defined) domain is
%f\n", sum((s - f).^2));
```

Problem 7

The sum of the squared errors over the (discretely-defined) domain is 0.389921



Problem 8

```
fprintf("\nProblem 8\n");
clear variables;

fprintf("f(x) = x^3 + sin(x) - 1\n");
x = 1:11+1;
f = x.^3 + sin(x) - 1;
c = polyfit(x, f, length(x)-1);

fprintf("f(4.5) = %f\n", polyval(c, 4.5));
fprintf("f(15) = %f\n", polyval(c, 15));

figure(); hold on;

domain = linspace(0, 15, 101);
lines = plot(domain, polyval(c, domain), 'r-');
points = plot(x, f, 'b.', "MarkerSize", 20);
wanted = plot([4.5, 15], [polyval(c, 4.5), polyval(c, 15)], 'g.',
"MarkerSize", 20);

grid on; axis([0, 15, -1, polyval(c, 15)]);
title("Interpolation and Extrapolation");
legend([points, lines, wanted], ["Data points", "Interpolation/Extrapolation",
"Interesting points"], "Location", "northwest");

hold off;
```

Problem 8

$$f(x) = x^3 + \sin(x) - 1$$

$$f(4.5) = 89.147466$$

$$f(15) = 3336.280580$$

