

TABLE 2.1 Estimates of genetic diversity and total inbreeding for each dataset that used **microsatellite** as marker. Species: *A. germinans* (Ag), *A. schaueriana* (As), *L. racemosa* (Lr), and *R. mangle* (Rm); Allelic richness (Ar); Expected heterozygosity (H_E); Observed heterozygosity (H_o); Inbreeding coefficient (F_{IS}); and the reference for each dataset. Results are shown as mean calculated across loci of all populations of each dataset \pm standard deviation.

Species	Ar	H_E	H_o	F_{IS}	References
Ag	2.543 \pm 0.302	0.456 \pm 0.071	0.672 \pm 0.154	-0.422 \pm 0.154	Hodel et al., 2016
Ag	2.260 \pm 0.390	0.325 \pm 0.111	0.276 \pm 0.127	0.181 \pm 0.187	Kennedy et al., 2020
Ag	2.624 \pm 0.751	0.438 \pm 0.171	0.432 \pm 0.306	0.218 \pm 0.244	Mori et al., 2015
Ag	2.551 \pm 0.658	0.239 \pm 0.108	0.224 \pm 0.302	0.132 \pm 0.158	O-Zavala et al., 2019
As	2.416 \pm 0.563	0.236 \pm 0.086	0.171 \pm 0.063	0.248 \pm 0.104	Mori et al., 2015
Lr	1.551 \pm 0.398	0.204 \pm 0.160	0.106 \pm 0.059	0.475 \pm 0.165	S-Lima et al., 2021
Rm	2.416 \pm 0.299	0.428 \pm 0.067	0.314 \pm 0.070	0.221 \pm 0.137	de la Cruz et al., 2018
Rm	1.798 \pm 0.724	0.196 \pm 0.156	0.190 \pm 0.105	-0.015 \pm 0.324	Francisco et al., 2018
Rm	2.452 \pm 0.247	0.425 \pm 0.062	0.489 \pm 0.105	-0.089 \pm 0.127	Hodel et al., 2016
Rm	2.350 \pm 0.257	0.419 \pm 0.071	0.493 \pm 0.112	-0.089 \pm 0.123	Hodel et al., 2017
Rm	2.532 \pm 0.500	0.350 \pm 0.122	0.310 \pm 0.106	0.093 \pm 0.125	Kennedy et al., 2016
Rm	2.407 \pm 0.395	0.317 \pm 0.096	0.265 \pm 0.129	0.211 \pm 0.222	Kennedy et al., 2017
Rm	1.729 \pm 0.237	0.370 \pm 0.113	0.366 \pm 0.129	0.026 \pm 0.221	Takayama et al., 2013

TABLE 2.2 Estimates of genetic diversity and total inbreeding for each dataset that used **SNP** as marker. Species: *A. germinans* (Ag), *A. schaueriana* (As), *L. racemosa* (Lr), and *R. mangle* (Rm); Allelic richness (Ar); Expected heterozygosity (H_E); Observed heterozygosity (H_O); Inbreeding coefficient (F_{IS}); and the reference for each dataset. Results are shown as mean calculated across loci of all populations of each dataset \pm standard deviation.

Species	Ar	H_E	H_O	F_{IS}	References
Ag	1.242 \pm 0.210	0.246 \pm 0.132	0.271 \pm 0.232	-0.101 \pm 0.378	Cruz et al., 2020
As	1.567 \pm 0.354	0.304 \pm 0.160	0.439 \pm 0.318	-0.442 \pm 0.397	Cruz et al., 2019
Lr	1.104 \pm 0.231	0.118 \pm 0.103	0.098 \pm 0.177	0.171 \pm 0.507	Hodel et al., 2018
Rm	1.317 \pm 0.224	0.311 \pm 0.153	0.399 \pm 0.359	-0.283 \pm 0.580	Hodel et al., 2017
Rm	1.063 \pm 0.180	0.063 \pm 0.085	0.058 \pm 0.092	0.081 \pm 0.441	Hodel et al., 2018
Rm	1.910 \pm 0.247	0.286 \pm 0.149	0.084 \pm 0.108	0.707 \pm 0.334	Madeira et al., 2023

References

- Cisneros-de la Cruz, D. J., Martínez-Castillo, J., Herrera-Silveira, J., Yáñez-Espinosa, L., Ortiz-García, M., Us-Santamaria, R., & Andrade, J. L. (2018). Short-distance barriers affect genetic variability of *Rhizophora mangle* L. in the Yucatan Peninsula. *Ecology and Evolution*, 8(22), 11083–11099. <https://doi.org/10.1002/ece3.4575>
- Cruz, M. V., Mori, G. M., Oh, D. H., Dassanayake, M., Zucchi, M. I., Oliveira, R. S., & Souza, A. P. de. (2020). Molecular responses to freshwater limitation in the mangrove tree *Avicennia germinans* (Acanthaceae). *Molecular Ecology*, 29(2), 344–362. <https://doi.org/10.1111/mec.15330>
- Cruz, M. V., Mori, G. M., Signori-Müller, C., da Silva, C. C., Oh, D. H., Dassanayake, M., Zucchi, M. I., Oliveira, R. S., & de Souza, A. P. (2019). Local adaptation of a dominant coastal tree to freshwater availability and solar radiation suggested by genomic and ecophysiological approaches. *Scientific Reports*, 9(1), 1–15. <https://doi.org/10.1038/s41598-019-56469-w>

- Francisco, P. M., Mori, G. M., Alves, F. M., Tambarussi, E. V., & de Souza, A. P. (2018). Population genetic structure, introgression, and hybridization in the genus *Rhizophora* along the Brazilian coast. *Ecology and Evolution*, 8(6), 3491–3504. <https://doi.org/10.1002/ece3.3900>
- Hodel, R. G. J., Chen, S., Payton, A. C., Mcdaniel, S. F., Soltis, P., & Soltis, D. E. (2017). Adding loci improves phylogeographic resolution in red mangroves despite increased missing data : comparing microsatellites and RAD-Seq and investigating loci filtering. *Scientific Reports*, November, 1–14. <https://doi.org/10.1038/s41598-017-16810-7>
- Hodel, R. G. J., De Souza Cortez, M. B., Soltis, P. S., & Soltis, D. E. (2016). Comparative phylogeography of black mangroves (*Avicennia germinans*) and red mangroves (*Rhizophora mangle*) in Florida: Testing the maritime discontinuity in coastal plants. *American Journal of Botany*, 103(4), 730–739. <https://doi.org/10.3732/ajb.1500260>
- Hodel, R. G. J., Knowles, L. L., McDaniel, S. F., Payton, A. C., Dunaway, J. F., Soltis, P. S., & Soltis, D. E. (2018). Terrestrial species adapted to sea dispersal: Differences in propagule dispersal of two Caribbean mangroves. *Molecular Ecology*, 27(22), 4612–4626. <https://doi.org/10.1111/mec.14894>
- Kennedy, J. P., Garavelli, L., Truelove, N. K., Devlin, D. J., Box, S. J., Ch, L. M., & Feller, I. C. (2017). *Contrasting genetic effects of red mangrove (Rhizophora mangle L .) range expansion along West and East Florida*. 335–347. <https://doi.org/10.1111/jbi.12813>
- Kennedy, J. P., Pil, M. W., Proffi, C. E., Boeger, W. A., Stanford, A. M., & Devlin, D. J. (2016). *Postglacial expansion pathways of red mangrove , Rhizophora mangle , in the Caribbean Basin and Florida I*. 103(2), 260–276. <https://doi.org/10.3732/ajb.1500183>
- Kennedy, J. P., Preziosi, R. F., Rowntree, J. K., & Feller, I. C. (2020). Is the central-marginal hypothesis a general rule? Evidence from three distributions of an expanding mangrove species, *Avicennia germinans* (L.) L. *Molecular Ecology*, 29(4), 704–719. <https://doi.org/10.1111/mec.15365>
- Madeira, A. G., Tsuda, Y., Nagano, Y., Iwasaki, T., Zucchi, M. I., Kajita, T., & Mori,

- G. M. (2023). The role of oceanic currents in the dispersal and connectivity of the mangrove *Rhizophora mangle* on the Southwest Atlantic region. *Molecular Ecology Resources*. <https://doi.org/10.1111/1755-0998.13807>
- Mori, G. M., Zucchi, M. I., & Souza, A. P. (2015). Multiple-geographic-scale genetic structure of two mangrove tree species: The roles of mating system, hybridization, limited dispersal and extrinsic factors. *PLoS ONE*, *10*(2), 1–23. <https://doi.org/10.1371/journal.pone.0118710>
- Ochoa-Zavala, M., Jaramillo-Correa, J. P., Piñero, D., Nettel-Hernanz, A., & Núñez-Farfán, J. (2019). Contrasting colonization patterns of black mangrove (*Avicennia germinans* (L.) L.) gene pools along the Mexican coasts. *Journal of Biogeography*, *46*(5), 884–898. <https://doi.org/10.1111/jbi.13536>
- Sereneski-Lima, C., Baggio, R. A., Pil, M. W., Torres Boeger, M. R., & Boeger, W. A. (2021). Historical and contemporary factors affect the genetic diversity and structure of *Laguncularia racemosa* (L.) Gaertn, along the western Atlantic coast. *Estuarine, Coastal and Shelf Science*, *249*(September 2019), 107055. <https://doi.org/10.1016/j.ecss.2020.107055>
- Takayama, K., Tamura, M., Tateishi, Y., Webb, E. L., & Kajita, T. (2013). Strong genetic structure over the American continents and transoceanic dispersal in the mangrove genus *Rhizophora* (Rhizophoraceae) revealed by broad-scale nuclear and chloroplast DNA analysis. *American Journal of Botany*, *100*(6), 1191–1201. <https://doi.org/10.3732/ajb.1200567>