

TABLE S2 Mantel modeling tests performed with symmetric matrices between Slatkin's linearized F_{ST} (D) and shortest distance through water and Euclidean distance and shortest distance through water controlled by Euclidean distance (Partial mantel). Pearson correlation values (r). Significance values (p), showed in bold for $p < 0.05$. Molecular marker: SNP or Msat (microsatellite) and Reference of data source are shown. ¹ Atlantic Ocean Basin; ² Pacific Ocean Basin. The information was not shown when the test could not be performed: the matrices presented dimensions 2x2 ($k=2$).

a) <i>Avicennia germinans</i>				
Mantel D		Partial Mantel D		
Water r (p)	Euclidean r (p)	Water controlled by Euclidean r (p)	Marker	Reference
-0.83 (0.83)	0.03 (0.50)	-1 (1.00)	Msat	Hodel et al., 2016
0.29 (0.50)	0.72 (0.17)	-1 (1.00)	Msat	Kennedy et al., 2020
-0.29 (0.79)	-0.27 (0.79)	-0.29 (0.71)	Msat	O-Zavala et al., 2019 ¹
0.83 (0.17)	0.78 (0.17)	1 (0.66)	SNP	Cruz et al., 2020
b) <i>Laguncularia racemosa</i>				
0.43 (0.16)	0.46 (0.13)	-0.06 (0.43)	SNP	Hodel et al., 2018
c) <i>Rhizophora mangle</i>				
-0.99 (1.00)	-0.62 (0.84)	-1 (1.00)	Msat	Hodel et al., 2016
0.58 (0.11)	0.54 (0.16)	0.31 (0.16)	Msat	Hodel et al., 2017
-0.82 (1.00)	-0.11 (0.67)	-1 (1.00)	Msat	Kennedy et al., 2016
0.28 (0.33)	-0.40 (0.75)	0.21 (0.38)	Msat	Kennedy et al., 2017
-0.18 (0.59)	-0.19 (0.60)	0.01 (0.49)	Msat	Takayama et al., 2013 ²
0.58 (0.11)	0.54 (0.16)	0.31 (0.16)	SNP	Hodel et al., 2017
0.53 (0.04)	0.59 (0.03)	-0.18 (0.67)	SNP	Hodel et al., 2018

References

- Cruz, M. V., Mori, G. M., Oh, D. H., Dassanayake, M., Zucchi, M. I., Oliveira, R. S., & Souza, A. P. de. (2020). Molecular responses to freshwater limitation in the mangrove tree *Avicennia germinans* (Acanthaceae). *Molecular Ecology*, 29(2), 344–362.
<https://doi.org/10.1111/mec.15330>
- Hodel, R. G. J., Chen, S., Payton, A. C., Mcdaniel, S. F., Soltis, P., & Soltis, D. E. (2017). Adding loci improves phylogeographic resolution in red mangroves despite increased missing data : comparing microsatellites and RAD-Seq and investigating loci filtering. *Scientific Reports*, November, 1–14. <https://doi.org/10.1038/s41598-017-16810-7>
- Hodel, R. G. J., De Souza Cortez, M. B., Soltis, P. S., & Soltis, D. E. (2016). Comparative phylogeography of black mangroves (*Avicennia germinans*) and red mangroves (*Rhizophora mangle*) in Florida: Testing the maritime discontinuity in coastal plants. *American Journal of*

Botany, 103(4), 730–739. <https://doi.org/10.3732/ajb.1500260>

- Hodel, R. G. J., Knowles, L. L., McDaniel, S. F., Payton, A. C., Dunaway, J. F., Soltis, P. S., & Soltis, D. E. (2018). Terrestrial species adapted to sea dispersal: Differences in propagule dispersal of two Caribbean mangroves. *Molecular Ecology*, 27(22), 4612–4626. <https://doi.org/10.1111/mec.14894>
- Kennedy, J. P., Garavelli, L., Truelove, N. K., Devlin, D. J., Box, S. J., Ch, L. M., & Feller, I. C. (2017). *Contrasting genetic effects of red mangrove (Rhizophora mangle L .) range expansion along West and East Florida*. 335–347. <https://doi.org/10.1111/jbi.12813>
- Kennedy, J. P., Pil, M. W., Proffi, C. E., Boeger, W. A., Stanford, A. M., & Devlin, D. J. (2016). *Postglacial expansion pathways of red mangrove , Rhizophora mangle , in the Caribbean Basin and Florida I*. 103(2), 260–276. <https://doi.org/10.3732/ajb.1500183>
- Kennedy, J. P., Preziosi, R. F., Rowntree, J. K., & Feller, I. C. (2020). Is the central-marginal hypothesis a general rule? Evidence from three distributions of an expanding mangrove species, *Avicennia germinans* (L.) L. *Molecular Ecology*, 29(4), 704–719. <https://doi.org/10.1111/mec.15365>
- Ochoa-Zavala, M., Jaramillo-Correa, J. P., Piñero, D., Nettel-Hernanz, A., & Núñez-Farfán, J. (2019). Contrasting colonization patterns of black mangrove (*Avicennia germinans* (L.) L.) gene pools along the Mexican coasts. *Journal of Biogeography*, 46(5), 884–898. <https://doi.org/10.1111/jbi.13536>
- Takayama, K., Tamura, M., Tateishi, Y., Webb, E. L., & Kajita, T. (2013). Strong genetic structure over the American continents and transoceanic dispersal in the mangrove genus *Rhizophora* (Rhizophoraceae) revealed by broad- scale nuclear and chloroplast DNA analysis. *American Journal of Botany*, 100(6), 1191–1201. <https://doi.org/10.3732/ajb.1200567>