

References

- Altamura, C., Reinhard, M., Vry, M.-S., Kaller, C. P., Hamzei, F., Vernieri, F., Rossini, P. M., Hetzel, A., Weiller, C., & Saur, D. (2009). The longitudinal changes of BOLD response and cerebral hemodynamics from acute to subacute stroke. A fMRI and TCD study. *BMC Neuroscience*, *10*, 1. <https://doi.org/10.1186/1471-2202-10-151>
- Arichi, T., Fagiolo, G., Varela, M., Melendez-Calderon, A., Allievi, A., Merchant, N., Tusor, N., Counsell, S. J., Burdet, E., Beckmann, C. F., & David, E. A. (2012). Development of BOLD signal hemodynamic responses in the human brain. *NeuroImage*, *63*, 2. <https://doi.org/10.1016/j.neuroimage.2012.06.054>
- Bonakdarpour, B., B, P. T., & K, T. C. (2007). Hemodynamic response function in patients with stroke-induced aphasia: Implications for fMRI data analysis. *NeuroImage*, *36*, 2. <https://doi.org/10.1016/j.neuroimage.2007.02.035>
- Calhoun, V., Adalı, T., Kraut, M., & Pearlson, G. (2000). A weighted least-squares algorithm for estimation and visualization of relative latencies in event-related functional MRI. *Magnetic Resonance in Medicine*, *44*, 6. [https://doi.org/10.1002/1522-2594\(200012\)44:6%3C947::AID-MRM17%3E3.0.CO;2-5](https://doi.org/10.1002/1522-2594(200012)44:6%3C947::AID-MRM17%3E3.0.CO;2-5)
- Cignetti, F., Salvia, E., Anton, J.-L., Grosbras, M.-H., & Assaiante, C. (2016). Pros and Cons of Using the Informed Basis Set to Account for Hemodynamic Response Variability with Developmental Data. *Frontiers in Neuroscience*, *10*. <https://doi.org/10.3389/fnins.2016.00322>
- Duann, J.-R., Jung, T.-P., Kuo, W.-J., Yeh, T.-C., Makeig, S., Hsieh, J.-C., & Sejnowski, T. J. (2002). Single-Trial Variability in Event-Related BOLD Signals. *NeuroImage*, *15*, 4. <https://doi.org/10.1006/nimg.2001.1049>

- Duarte, J. V., Guerra, C., Moreno, C., Gomes, L., & Castelo-Branco, M. (2023). Changes in hemodynamic response function components reveal specific changes in neurovascular coupling in type 2 diabetes. *Frontiers in Physiology*, *13*, 1101470.
<https://doi.org/10.3389/fphys.2022.1101470>
- Grady, C. (2012). The cognitive neuroscience of ageing. *Nature Reviews Neuroscience*, *13*, 7.
<https://doi.org/10.1038/nrn3256>
- J, F. K., Mechelli, A., Turner, R., & J, P. C. (2000). Nonlinear Responses in fMRI: The Balloon Model, Volterra Kernels, and Other Hemodynamics. *NeuroImage*, *12*, 4.
<https://doi.org/10.1006/nimg.2000.0630>
- Marrelec, G., Benali, H., Ciuciu, P., Pélérini-Issac, M., & Poline, J. (2003). Robust Bayesian estimation of the hemodynamic response function in event-related BOLD fMRI using basic physiological information. *Human Brain Mapping*, *19*, 1.
<https://doi.org/10.1002/hbm.10100>
- Ogawa, S., & Sung, Y.-W. (2007). Functional magnetic resonance imaging. *Scholarpedia*, *2*, 10.
<https://doi.org/10.4249/scholarpedia.3105>
- Pedregosa-Izquierdo, F. (2015). *Feature extraction and supervised learning on fMRI : from practice to theory*. <https://theses.hal.science/tel-01100921>
- Richter, W., & Richter, M. (2003). The shape of the fMRI BOLD response in children and adults changes systematically with age. *NeuroImage*, *20*, 2. [https://doi.org/10.1016/S1053-8119\(03\)00347-1](https://doi.org/10.1016/S1053-8119(03)00347-1)