

TOPICS IN ALGEBRAIC LOGIC AND DUALITY THEORY

Lecture 4

Rodrigo N. Almeida, Simon Lemal

June 12, 2025

- Maksimova's characterisation of superintuitionistic logics with the interpolation property

A superintuitionistic logic has the interpolation property iff it is one of

CPC,

$LC_2 = IPC + bd_2 + bw_1$ = logic of the 2-chain,

$LC = IPC + p \rightarrow q \vee q \rightarrow p$ = logic of chains,

$KC = IPC + \neg p \vee \neg\neg p$ = logic of directed frames,

$BD_2 = IPC + bd_2$ = logic of frames of depth at most 2,

$BDW_2 = BD_2 + bw_2$ = logic of the 2-fork,

IPC.

Understanding the statement

Definition

We define a sequence of formulas \mathbf{bd}_n inductively:

$$\begin{aligned}\mathbf{bd}_0 &= \perp, \\ \mathbf{bd}_{n+1} &= p_{n+1} \vee (p_{n+1} \rightarrow \mathbf{bd}_n),\end{aligned}$$

Definition

We define a sequence of formulas \mathbf{bd}_n inductively:

$$\begin{aligned}\mathbf{bd}_0 &= \perp, \\ \mathbf{bd}_{n+1} &= p_{n+1} \vee (p_{n+1} \rightarrow \mathbf{bd}_n),\end{aligned}$$

Proposition

A frame validates \mathbf{bd}_n iff it doesn't contain a chain of $n + 1$ points, i.e. iff its depth is bounded by n .

Definition

We define a sequence of formulas \mathbf{bd}_n inductively:

$$\begin{aligned}\mathbf{bd}_0 &= \perp, \\ \mathbf{bd}_{n+1} &= p_{n+1} \vee (p_{n+1} \rightarrow \mathbf{bd}_n),\end{aligned}$$

Proposition

A frame validates \mathbf{bd}_n iff it doesn't contain a chain of $n + 1$ points, i.e. iff its depth is bounded by n .

Proof.

Exercise. □

Definition

We define formulas bw_n as follows:

$$\text{bw}_n = \bigvee_{i=0}^n \left(p_i \rightarrow \bigvee_{j \neq i} p_j \right).$$

Definition

We define formulas \mathbf{bw}_n as follows:

$$\mathbf{bw}_n = \bigvee_{i=0}^n \left(p_i \rightarrow \bigvee_{j \neq i} p_j \right).$$

Proposition

A frame validates \mathbf{bw}_n iff none of its rooted subframes contain an antichain of $n + 1$ points, i.e. iff its width is bounded by n .

Definition

We define formulas \mathbf{bw}_n as follows:

$$\mathbf{bw}_n = \bigvee_{i=0}^n \left(p_i \rightarrow \bigvee_{j \neq i} p_j \right).$$

Proposition

A frame validates \mathbf{bw}_n iff none of its rooted subframes contain an antichain of $n + 1$ points, i.e. iff its width is bounded by n .

Proof.

Exercise. □

Preliminary tools

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Corollary

Let F be a finite rooted frame and L be a logic. We have $\chi(F) \notin L$ iff $F \models L$.

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Corollary

Let F be a finite rooted frame and L be a logic. We have $\chi(F) \notin L$ iff $F \models L$.

Proof.

Exercise. □

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Corollary

Let F be a finite rooted frame and L be a logic. We have $\chi(F) \notin L$ iff $F \models L$.

Proof.

Exercise. □

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(P) \mid P \not\models L\}.$$

Corollary

The logic of a finite rooted frame is axiomatisable by Jankov formulas.

Definition

We turn the class of finite rooted frames into a poset by setting $F \leqslant G$ iff G is a p-morphic image of F .

Definition

We turn the class of finite rooted frames into a poset by setting $F \leqslant G$ iff G is a p-morphic image of F .

Proposition

Let $\mathcal{F}' \subseteq \mathcal{F}$ be classes of finite rooted frames such that every frame in \mathcal{F} is below a frame in \mathcal{F}' . Then

$$\text{IPC} + \{\chi(F) \mid F \in \mathcal{F}\} = \text{IPC} + \{\chi(F) \mid F \in \mathcal{F}'\}.$$

Lemma

Let F be a frame and $x, y \in F$ be distinct.

- 1. Suppose that y is the only immediate successor of x , i.e. $\uparrow y = \uparrow x \setminus \{x\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an α -reduction.*

Lemma

Let F be a frame and $x, y \in F$ be distinct.

1. Suppose that y is the only immediate successor of x , i.e. $\uparrow y = \uparrow x \setminus \{x\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an α -reduction.
2. Suppose that x and y have the same immediate successors, i.e. $\uparrow x \setminus \{x\} = \uparrow y \setminus \{y\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an β -reduction.

Lemma

Let F be a frame and $x, y \in F$ be distinct.

1. Suppose that y is the only immediate successor of x , i.e. $\uparrow y = \uparrow x \setminus \{x\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an α -reduction.
2. Suppose that x and y have the same immediate successors, i.e. $\uparrow x \setminus \{x\} = \uparrow y \setminus \{y\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an β -reduction.

Lemma

Let F be a frame and $x, y \in F$ be distinct.

1. Suppose that y is the only immediate successor of x , i.e. $\uparrow y = \uparrow x \setminus \{x\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an α -reduction.
2. Suppose that x and y have the same immediate successors, i.e. $\uparrow x \setminus \{x\} = \uparrow y \setminus \{y\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an β -reduction.

Proof.

Routine check. □

Proposition

Let $f: F \rightarrow G$ be an onto p -morphism between finite frames. Then f factors as a sequence of α - and β -reductions.

Lemma

Let F be a frame and $x, y \in F$ be distinct.

1. Suppose that y is the only immediate successor of x , i.e. $\uparrow y = \uparrow x \setminus \{x\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an α -reduction.
2. Suppose that x and y have the same immediate successors, i.e. $\uparrow x \setminus \{x\} = \uparrow y \setminus \{y\}$. Then the equivalence relation identifying x and y is a bisimulation equivalence, and the quotient map an β -reduction.

Proof.

Routine check. □

Proposition

Let $f: F \rightarrow G$ be an onto p -morphism between finite frames. Then f factors as a sequence of α - and β -reductions.

Proof.

Exercise. Hint: induction on the number of points that are identified. □

See board.

Maksimova's characterisation

\Leftarrow : Exercise.

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation.

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \text{CPC}$, we are done.

Proof of Maksimova's characterisation

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \mathbf{CPC}$, we are done. Otherwise, observe that \mathbf{CPC} is the logic of the 1-frame, whose only cover is the 2-chain.



(a) The 1-frame



(b) The 2-chain

Proof of Maksimova's characterisation

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \mathbf{CPC}$, we are done. Otherwise, observe that \mathbf{CPC} is the logic of the 1-frame, whose only cover is the 2-chain.



(a) The 1-frame



(b) The 2-chain

Therefore \mathbf{CPC} is axiomatised by the Jankov formula χ_{2c} of the 2-chain.

Proof of Maksimova's characterisation

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \mathbf{CPC}$, we are done. Otherwise, observe that \mathbf{CPC} is the logic of the 1-frame, whose only cover is the 2-chain.



(a) The 1-frame



(b) The 2-chain

Therefore \mathbf{CPC} is axiomatised by the Jankov formula χ_{2c} of the 2-chain. As $L \subsetneq \mathbf{CPC}$, we have $\chi_{2c} \notin L$, thus the 2-chain is an L -frame.

If L is the logic of the 2-chain, then $L = \mathbf{LC}_2$, and we are done.

Otherwise, observe that the only covers of the 2-chain are the 3-chain and the 2-fork.



(a) 3-chain



(b) 2-fork

Therefore, \mathbf{LC}_2 is axiomatized by the Jankov formulas χ_{3c} of the 3-chain and χ_{2f} of the 2-fork.

Otherwise, observe that the only covers of the 2-chain are the 3-chain and the 2-fork.



(a) 3-chain



(b) 2-fork

Therefore, \mathbf{LC}_2 is axiomatized by the Jankov formulas χ_{3c} of the 3-chain and χ_{2f} of the 2-fork. As $L \subsetneq \mathbf{L}_2$, we have $\chi_{3c} \notin L$ or $\chi_{2f} \notin L$, thus either the 3-chain or the 2-fork are L -frames.

Appendix

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(P) \mid P \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(P) \mid P \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(P) \mid P \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq :



Thank you!
Questions?