

HEYTING ALGEBRAS ONE STEP AT A TIME

Rodrigo Nicolau Almeida – ILLC-UvA – University of Birmingham TCS Seminar

Friday 4, 2025

1. Some open problems of intuitionistic logic.

1. Some open problems of intuitionistic logic.
2. A particular technical problem I have worked on related to free Heyting algebras.

1. Some open problems of intuitionistic logic.
2. A particular technical problem I have worked on related to **free Heyting algebras**.
3. Some hopes for these, and related, techniques in furthering the theory concerning those open problems.

Intuitionistic Logic and its Extensions

Intuitionistic Logic – IPC – is an attempt to capture the “constructive” fragment of logic. This makes it suitable for many computational applications.

Intuitionistic Logic – IPC – is an attempt to capture the “constructive” fragment of logic. This makes it suitable for many computational applications.

Its algebraic models are also mathematically quite natural:

Definition

An algebra $(H, \wedge, \vee, \rightarrow, 0, 1)$ is called a *Heyting algebra* if:

1. $(H, \wedge, \vee, 0, 1)$ is a (distributive) lattice.
2. The following law holds for all $a, b, c \in H$:

$$a \wedge c \leq b \iff c \leq a \rightarrow b.$$

Intuitionistic Logic – IPC – is an attempt to capture the “constructive” fragment of logic. This makes it suitable for many computational applications.

Its algebraic models are also mathematically quite natural:

Definition

An algebra $(H, \wedge, \vee, \rightarrow, 0, 1)$ is called a *Heyting algebra* if:

1. $(H, \wedge, \vee, 0, 1)$ is a (distributive) lattice.
2. The following law holds for all $a, b, c \in H$:

$$a \wedge c \leq b \iff c \leq a \rightarrow b.$$

For example, given a topological space (X, τ) , the topology defines a Heyting algebra with Heyting implication given by $\text{int}(X - U \cup V)$ for U, V open sets. Such algebras are called **topological Heyting algebras**.

Intuitionistic Logic – IPC – is an attempt to capture the “constructive” fragment of logic. This makes it suitable for many computational applications.

Its algebraic models are also mathematically quite natural:

Definition

An algebra $(H, \wedge, \vee, \rightarrow, 0, 1)$ is called a *Heyting algebra* if:

1. $(H, \wedge, \vee, 0, 1)$ is a (distributive) lattice.
2. The following law holds for all $a, b, c \in H$:

$$a \wedge c \leq b \iff c \leq a \rightarrow b.$$

For example, given a topological space (X, τ) , the topology defines a Heyting algebra with Heyting implication given by $\text{int}(X - U \cup V)$ for U, V open sets. Such algebras are called **topological Heyting algebras**. As a special example, if (P, \leq) is a poset, $\text{Up}(P)$ the upwards closed sets of P , are a Heyting algebra. Also every **locale** is a (complete) Heyting algebra.

Often we are interested not just in a single logic, but in the class of its extensions:

Definition

Let L be a set of formulas in the language \mathcal{L}_{IPC} of intuitionistic logic. We say that L is a **superintuitionistic logic** if $IPC \subseteq L$ and L is closed under **Modus Ponens** and **Uniform Substitution**.

We denote by $\Lambda(IPC)$ the lattice of extensions of IPC .

On the algebraic side we dually have $\Lambda(HA)$, the lattice of subvarieties of Heyting algebras.

Often we are interested not just in a single logic, but in the class of its extensions:

Definition

Let L be a set of formulas in the language \mathcal{L}_{IPC} of intuitionistic logic. We say that L is a **superintuitionistic logic** if $IPC \subseteq L$ and L is closed under **Modus Ponens** and **Uniform Substitution**.

We denote by $\Lambda(IPC)$ the lattice of extensions of IPC .

On the algebraic side we dually have $\Lambda(\mathbf{HA})$, the lattice of subvarieties of Heyting algebras.

Studying extensions as a whole gives much deeper insight into the logical system and into the uniqueness of certain extensions and of the properties.

Often we are interested not just in a single logic, but in the class of its extensions:

Definition

Let L be a set of formulas in the language \mathcal{L}_{IPC} of intuitionistic logic. We say that L is a **superintuitionistic logic** if $IPC \subseteq L$ and L is closed under **Modus Ponens** and **Uniform Substitution**.

We denote by $\Lambda(IPC)$ the lattice of extensions of IPC .

On the algebraic side we dually have $\Lambda(HA)$, the lattice of subvarieties of Heyting algebras.

Studying extensions as a whole gives much deeper insight into the logical system and into the uniqueness of certain extensions and of the properties.

An example:

Theorem

There are exactly seven superintuitionistic logics with the interpolation property; equivalently seven varieties of Heyting algebras with the amalgamation property.

Definition

An ordered topological space (X, \leq, τ) is called a **Priestley space** if it is compact and whenever $x \not\leq y$ there is a clopen upwards closed set (upset) U such that $x \in U$ and $y \notin U$. It is called **Esakia** if whenever U is clopen, then $\downarrow U = \{x \in X : \exists y \in U, x \leq y\}$ is clopen.




Figure 1: Hilary Priestley (info missing); Leo Esakia (1934-2010)

Definition

A given order-preserving map $p : X \rightarrow Y$ between partial orders is called a *p-morphism* if whenever $p(x) \leq y$ there is some $x' \geq x$ such that $p(x') = y$.

Theorem

*There is a dual equivalence between **DL** distributive lattices and **Pries** the category of Priestley spaces with order-preserving continuous functions. This restricts to a dual equivalence between **HA**, Heyting algebras and their homomorphisms, and **Esa**, Esakia spaces with continuous p -morphisms.*

Theorem

*There is a dual equivalence between **DL** distributive lattices and **Pries** the category of Priestley spaces with order-preserving continuous functions. This restricts to a dual equivalence between **HA**, Heyting algebras and their homomorphisms, and **Esa**, Esakia spaces with continuous p -morphisms.*

All finite posets are Esakia spaces. Moreover, given a class \mathcal{K} of Esakia spaces we can look at their **logic**: it is the logic of the dual Heyting algebras.

The extensions of IPC have been extensively studied for more than 60 years. Nevertheless some longstanding open problems have remained open.

The extensions of IPC have been extensively studied for more than 60 years. Nevertheless some longstanding open problems have remained open.

I will briefly mention three such problems which are old and have revealed themselves quite sticky.

1. Medvedev's Logic

Consider the family of finite posets of **topless Boolean algebras**:




Figure 2: Medvedev frames

Let ML be the logic of these infinitely many posets.

1. Medvedev's Logic

Consider the family of finite posets of **topless Boolean algebras**:




Figure 2: Medvedev frames

Let ML be the logic of these infinitely many posets.

Question: can you recursively axiomatize this logic?

1. Medvedev's Logic

Consider the family of finite posets of **topless Boolean algebras**:




Figure 2: Medvedev frames

Let ML be the logic of these infinitely many posets.

Question: can you recursively axiomatize this logic?

Other question: is this the only intermediate logic which is **structurally complete** and **has the disjunction property**?

1. Medvedev's Logic

Consider the family of finite posets of **topless Boolean algebras**:




Figure 2: Medvedev frames

Let ML be the logic of these infinitely many posets.

Question: can you recursively axiomatize this logic?

Other question: is this the only intermediate logic which is **structurally complete** and **has the disjunction property**?

This logic appears very naturally elsewhere: it can be formulated as a logic of **finite problems**; it is the natural model of inquisitive logic; etc.

2. Kuznetsov's Completeness Problem

Given an intermediate logic L , sometimes we do not need all (or most) algebras of L :

1. L is tabular if $L = \text{Log}(A)$ for a single finite algebra;
2. L has the FMP if L is the logic of a class of finite algebras;
3. L is *Kripke complete* if its class of algebras $\text{Alg}(L)$ is generated by the algebras of the form $\text{Up}(L)$;
4. L is *topologically complete* if it is generated by its topological Heyting algebras;
5. L is *locally complete* if it is generated by its locales.

2. Kuznetsov's Completeness Problem

Given an intermediate logic L , sometimes we do not need all (or most) algebras of L :

1. L is tabular if $L = \text{Log}(A)$ for a single finite algebra;
2. L has the FMP if L is the logic of a class of finite algebras;
3. L is *Kripke complete* if its class of algebras $\text{Alg}(L)$ is generated by the algebras of the form $\text{Up}(L)$;
4. L is *topologically complete* if it is generated by its topological Heyting algebras;
5. L is *locally complete* if it is generated by its locales.

Question: Is every logic L topologically/locally complete?

2. Kuznetsov's Completeness Problem

Given an intermediate logic L , sometimes we do not need all (or most) algebras of L :

1. L is tabular if $L = \text{Log}(A)$ for a single finite algebra;
2. L has the FMP if L is the logic of a class of finite algebras;
3. L is *Kripke complete* if its class of algebras $\text{Alg}(L)$ is generated by the algebras of the form $\text{Up}(L)$;
4. L is *topologically complete* if it is generated by its topological Heyting algebras;
5. L is *locally complete* if it is generated by its locales.

Question: Is every logic L topologically/locally complete?

This establishes a link between propositional and **First-order intuitionistic logic**.

3. Pitts-Pataaraia Problem

Given an elementary topos \mathcal{E} , the subobjects of the terminal object $\text{Sub}(1)$ form a Heyting algebra.

Question: Can every Heyting algebra arise in this way?

3. Pitts-Pataaraia Problem

Given an elementary topos \mathcal{E} , the subobjects of the terminal object $\text{Sub}(1)$ form a Heyting algebra.

Question: Can every Heyting algebra arise in this way?

This establishes a link with **Propositional Higher-order logic**, i.e., with particular models of the lambda calculus.

Free Heyting Algebras

Mantra

Every logic L is generated by its free algebras. Hence understanding the free algebras gives us crucial information about the structure of L .

A Tale of Nice Algebras

Mantra

Every logic L is generated by its free algebras. Hence understanding the free algebras gives us crucial information about the structure of L .

Example

1. Free **Boolean algebras** are very well-understood: for a set X they can be described as the duals of

$$2^X$$

understood as the X -fold product of the discrete space with two elements.

2. Similarly, for **Distributive Lattices**, the description of the free algebras is similarly straightforward: for a set X ,

$$2_{\bullet}^X$$

understood as the product of X -many copies of 2_{\bullet} , the two element poset $0 < 1$, with the pointwise order and topology.

A Tale of Nice Algebras

Mantra

Every logic L is generated by its free algebras. Hence understanding the free algebras gives us crucial information about the structure of L .

Example

1. Free **Boolean algebras** are very well-understood: for a set X they can be described as the duals of

$$2^X$$

understood as the X -fold product of the discrete space with two elements.

2. Similarly, for **Distributive Lattices**, the description of the free algebras is similarly straightforward: for a set X ,

$$2^X_\bullet$$

understood as the product of X -many copies of 2_\bullet , the two element poset $0 < 1$, with the pointwise order and topology.

Having a good grasp of free algebras is intimately related to having a good grasp of **colimits** of these algebras. **Logically**, this has several natural desirable applications: interpolation, conservativity, etc.




Figure 3: Rieger-Nishimura Lattice

Free n -generated Heyting algebras for $n > 1$ are very complex, in all the ways this can happen.

Free n -generated Heyting algebras for $n > 1$ are very complex, in all the ways this can happen.

The consequence of this complexity is that a number of natural related constructions which would otherwise be at our disposal have not been considered within the reasonable toolkit:

1. Coproducts of Heyting algebras (except for some locally finite subvarieties);
2. Pushouts of Heyting algebras;
3. Heyting subalgebras generated by a given subset;
4. Etc.

Describing the freely generated algebras amounts to studying the adjunction:

$$\mathbf{HA} \begin{array}{c} \xrightarrow{\text{Free}} \\ \xleftarrow{\text{Forget}} \end{array} \mathbf{Set}$$

Figure 4: Free-Forgetful Adjunction

Heyting Algebras from Distributive Lattices

Describing the freely generated algebras amounts to studying the adjunction:

$$\mathbf{HA} \begin{array}{c} \xrightarrow{\text{Free}} \\ \xleftarrow{\text{Forget}} \end{array} \mathbf{Set}$$

Figure 4: Free-Forgetful Adjunction

Instead of studying it directly, we “split” this adjunction:



Figure 5: Split Free Forgetful Adjunction

Key Intuition: We think of generating Heyting algebras as adding infinitely many layers of implications to a distributive lattice step-by-step.

Heyting Algebras from Distributive Lattices, algebraically

Let \mathcal{D} be a distributive lattice, and X a set of generators. Let $F_{DL}(X)$ be the free distributive lattice on X .

Heyting Algebras from Distributive Lattices, algebraically

Let \mathcal{D} be a distributive lattice, and X a set of generators. Let $F_{DL}(X)$ be the free distributive lattice on X .

Let $\mathcal{D}_0 = \mathcal{D}$. Then set

$$\mathcal{D}_1 = F_{DL}(\{a \rightarrow b : a, b \in D\})/\Theta$$

where Θ contains:

1. Axioms of Heyting algebras;
2. Axioms enforcing elements of the form $1 \rightarrow a$, for $a \in D$, to behave like the elements from D .

This gives us **one-step**.

Heyting Algebras from Distributive Lattices, algebraically

Let \mathcal{D} be a distributive lattice, and X a set of generators. Let $F_{DL}(X)$ be the free distributive lattice on X .

Let $\mathcal{D}_0 = \mathcal{D}$. Then set

$$\mathcal{D}_1 = F_{DL}(\{a \rightarrow b : a, b \in D\})/\Theta$$

where Θ contains:

1. Axioms of Heyting algebras;
2. Axioms enforcing elements of the form $1 \rightarrow a$, for $a \in D$, to behave like the elements from D .

This gives us **one-step**.

We set the map $i_1 : \mathcal{D}_0 \rightarrow \mathcal{D}_1$ sending a to $[1 \rightarrow a]$, which is a homomorphism by force. We then iterate; but for \mathcal{D}_2 we need to also add one more rule to Θ :

3. Axioms ensuring that if $a, b \in D$, then:

$$1 \rightarrow_{D_2} (a \rightarrow_{D_1} b) \equiv (1 \rightarrow_{D_1} a) \rightarrow_{D_2} (1 \rightarrow_{D_2} b).$$

This is how we define \mathcal{D}_2 , and i_2 is defined similarly.

We then iterate infinitely:

1. $\mathcal{D}_{n+1} = F_{DL}(\{a \rightarrow b : a, b \in \mathcal{D}_n\})/\Theta$, where Θ is defined as above;
2. $i_n : \mathcal{D}_n \rightarrow \mathcal{D}_{n+1}$ sends a to $[1 \rightarrow a]$.

Problem: We would like to say that the free Heyting algebra generated by \mathcal{D} is the union of all of these. But we do not know whether this is a chain of embeddings, or really anything about this construction.

We then iterate infinitely:

1. $\mathcal{D}_{n+1} = F_{DL}(\{a \rightarrow b : a, b \in \mathcal{D}_n\})/\Theta$, where Θ is defined as above;
2. $i_n : \mathcal{D}_n \rightarrow \mathcal{D}_{n+1}$ sends a to $[1 \rightarrow a]$.

Problem: We would like to say that the free Heyting algebra generated by \mathcal{D} is the union of all of these. But we do not know whether this is a chain of embeddings, or really anything about this construction.

For this we turn to duality.

The idea of dual one-step constructions is not-new, and well-studied in Modal Logic. In the case where we generate from **finite Heyting algebras**, this was studied by Ghilardi (1992), generalizing previous work of Urquhart (1973); Bezhanishvili and Gehrke (2009) gave a detailed outline of this method for various classes.

Let (X, \leq, τ) be a Priestley space (compact, totally order-disconnected space). We denote by $(V(X), \supseteq)$ the **Vietoris Hyperspace** of X with reverse inclusion.

Let (X, \leq, τ) be a Priestley space (compact, totally order-disconnected space). We denote by $(V(X), \supseteq)$ the **Vietoris Hyperspace** of X with reverse inclusion.

$V(X)$ is the set of closed subsets of X ; it has a basis consisting of sets:

$$[U] = \{C \in V(X) : C \subseteq U\} \text{ and } \langle V \rangle = \{C \in V(X) : C \cap V \neq \emptyset\}$$

where $U, V \in \text{Clop}(X)$. It is always a Priestley space.

Definition

Let X, Y, Z be Priestley spaces, and $g : X \rightarrow Y$ and $f : Z \rightarrow X$ be Priestley morphisms. We say that f is **open relative to g** (g -open for short) if it satisfies the following:

$$\forall a \in Z, \forall b \in X, (f(a) \leq b \implies \exists a' \in Z, (a \leq a' \ \& \ g(f(a')) = g(b))). \quad (*)$$

Given $S \subseteq X$ a closed subset, we say that S is **g -open** (understood as a poset with the restricted partial order relation) if the inclusion is itself g -open.

Definition

Let X, Y be a Priestley spaces, $g : X \rightarrow Y$ Priestley morphism. We denote by $V_r(X)$ the set of **closed and rooted** subsets of X . We denote by $V_g(X)$ the set of **closed, rooted and g -open** subsets of X .

This is a Priestley space as well.

We can now proceed with the main construction:

Definition

Let $g : X \rightarrow Y$ be a Priestley morphism. The **g-Vietoris complex** over X $(V_{\bullet}^g(X), \leq_{\bullet})$, is a sequence

$$(V_0(X), V_1(X), \dots, V_n(X))$$

connected by morphisms $r_i : V_{i+1}(X) \rightarrow V_i(X)$ such that:

1. $V_0(X) = X$;
2. $r_0 = g$
3. For $i \geq 0$, $V_{i+1}(X) := V_{r_i}(V_i(X))$;
4. $r_{i+1} = r_{r_i} : V_{i+1}(X) \rightarrow V_i(X)$ is the root map.

We can now proceed with the main construction:

Definition

Let $g : X \rightarrow Y$ be a Priestley morphism. The **g -Vietoris complex** over X $(V_{\bullet}^g(X), \leq, \bullet)$, is a sequence

$$(V_0(X), V_1(X), \dots, V_n(X))$$

connected by morphisms $r_i : V_{i+1}(X) \rightarrow V_i(X)$ such that:

1. $V_0(X) = X$;
2. $r_0 = g$
3. For $i \geq 0$, $V_{i+1}(X) := V_{r_i}(V_i(X))$;
4. $r_{i+1} = r_{r_i} : V_{i+1}(X) \rightarrow V_i(X)$ is the root map.

We denote the projective limit of this family by $V_G^g(X)$, and omit g when this is the terminal map to **1**.

Then one can show that $V_G^g(X)$ is always an **Esakia space**.

By considering maps appropriately, one can use this to prove:

Corollary

*The assignment V_G is a functor mapping **Pries** of Priestley spaces and Priestley morphisms, to the category **Esa** of Esakia spaces and p -morphisms. It is the right adjoint of the inclusion.*

By considering maps appropriately, one can use this to prove:

Corollary

*The assignment V_G is a functor mapping **Pries** of Priestley spaces and Priestley morphisms, to the category **Esa** of Esakia spaces and p -morphisms. It is the right adjoint of the inclusion.*

In the short term we could capitalize on this construction and associated ideas for several results:

1. In (A., 2025, preprint available), I presented this theory as well as a similar adjunction for image-finite posets and profinite Heyting algebras;
2. In (A., Bezhanishvili, 2024 and A.,Dukic, 2025), we applied this to coalgebraic representations of intuitionistic modal logic, resolving an open problem in the field;
3. In (A. 2025, forthcoming), we obtained a proof of uniform interpolation using a dual representation of coproducts of Heyting algebras.

Step by Step as a technique

Consider the functor:

$$\text{Reg} : \mathbf{HA} \rightarrow \mathbf{BA};$$

Consider the functor:

$$\text{Reg} : \mathbf{HA} \rightarrow \mathbf{BA};$$

this functor has a left adjoint (by categorical magic). What does it look like?

Consider the functor:

$$\text{Reg} : \mathbf{HA} \rightarrow \mathbf{BA};$$

this functor has a left adjoint (by categorical magic). What does it look like?

One considers a specific Vietoris complex:

Let $M_\infty(Y) = V_G^r(V_{\max}(Y))$. Explicitly, this is the inverse limit of the sequence:

$$V(Y) \xleftarrow{r_1} V_{\max}(Y) \xleftarrow{r_2} V_2(Y) \xleftarrow{r_3} \dots$$

where $V_{n+1}(Y) = V_{r_n}(V_n(Y))$.

Proposition

The functor $\text{FreeM} : \mathbf{Stone} \rightarrow \mathbf{Esa}$ assigning each Stone space X to $M_\infty(X)$ is right adjoint to $\max : \mathbf{Esa} \rightarrow \mathbf{Stone}$.

A Heyting algebra is said to be **regular** if it is generated by its regularization.

Theorem

Given a Stone space X , $M_\infty(X)$ is always a regular Esakia space, and moreover, regular Esakia spaces are the coalgebras for the comonad induced by this functor.

Theorem

The logic ML is precisely the logic of all the spaces $V(X)$ for X a Stone space.

Hope: Understanding the role of Medvedev spaces in relation to general Esakia spaces might allow us to eventually find a way to mechanically axiomatize Medvedev logic, as well as understand how the disjunction property and admissibility of it work.

Definition

Let (H, \Box) be a Heyting algebra with an operator. We say that this is a **KM**-algebra if \Box satisfies for every $a, b \in H$:

1. (K axiom) $\Box(a \rightarrow b) \leq \Box a \rightarrow \Box b$;
2. (Lax) $a \leq \Box a$;
3. (Strictness) $\Box a \leq b \vee b \rightarrow a$;
4. (WF) $\Box a \rightarrow a \leq a$.

Heyting algebras admit a **KM**-algebra structure **uniquely**. But not every HA admits such a structure (e.g., the 2-generated free Heyting algebra).

Definition

Let (H, \Box) be a Heyting algebra with an operator. We say that this is a **KM**-algebra if \Box satisfies for every $a, b \in H$:

1. (K axiom) $\Box(a \rightarrow b) \leq \Box a \rightarrow \Box b$;
2. (Lax) $a \leq \Box a$;
3. (Strictness) $\Box a \leq b \vee b \rightarrow a$;
4. (WF) $\Box a \rightarrow a \leq a$.

Heyting algebras admit a **KM**-algebra structure **uniquely**. But not every HA admits such a structure (e.g., the 2-generated free Heyting algebra).

Question: Which logics are generated by their **KM**-algebras?

Kuznetsov, in an inaccessible paper, proved that:

Theorem

*Every superintuitionistic logic L is generated by its **KM**-algebras.*

A recent (2025) algebraic proof was obtained by Jibladze and Kuznetsov (*no relation*).

'Little' Kuznetsov

Kuznetsov, in an inaccessible paper, proved that:

Theorem

Every superintuitionistic logic L is generated by its **KM**-algebras.

A recent (2025) algebraic proof was obtained by Jibladze and Kuznetsov (*no relation*).

We have been able to dualize this proof; a scary picture:




Figure 6:

The interest of this is that to solve Kuznetsov's localic problem in the **positive** the minimal requirements go through generalizing the ideas of Jibladze and Kuznetsov:

1. Turning a Heyting algebra into a **KM**-algebra in the same variety, one adds for each $a \in H$ a given "fronton", which is represented by the **filter of dense elements over a** ;
2. To "complete" a Heyting algebra in the same variety, one should at least be able to add for each $S \subseteq H$ a given join to this subset, which is represented by the **filter of upper bounds of S** .

Look upon the board.

Thank you!
Questions?