

REGULAR HEYTING ALGEBRAS AND FREE HEYTING EXTENSIONS OF BOOLEAN ALGEBRAS

RODRIGO NICOLAU ALMEIDA

1. FREE HEYTING EXTENSIONS

The¹ categories **HA** and **BA**, respectively of Heyting algebras with Heyting algebra homomorphisms and Boolean algebras with Boolean algebra homomorphisms, are related by a chain of adjunctions:

$$\mathbf{FreeM} \dashv \mathbf{Reg} \dashv I \dashv \mathbf{Center},$$

where

- (1) $I : \mathbf{BA} \rightarrow \mathbf{HA}$ is the inclusion;
- (2) $\mathbf{Center} : \mathbf{HA} \rightarrow \mathbf{BA}$ takes the *center* of a Heyting algebra H , namely $\mathbf{Center}(H) = \{a \in H : a \vee \neg a = 1\}$;
- (3) $\mathbf{Reg} : \mathbf{HA} \rightarrow \mathbf{BA}$ takes the *regular elements* of a Heyting algebra H namely $\mathbf{Reg}(H) = \{a \in H : \neg\neg a = a\}$.

The functor $\mathbf{FreeM} : \mathbf{BA} \rightarrow \mathbf{HA}$ is guaranteed to exist by the Adjoint Functor theorem; a description using a word construction can be derived from the work of Moraschini [5] (see also [6, Example 4.18]). It was likewise studied in [7], where it was shown to be fully faithful.

In this paper we give a more concrete presentation of the functor \mathbf{FreeM} , by studying the dual problem, employing Esakia duality (see [3] for any unexplained notions such as Priestley spaces, Esakia spaces and p-morphisms): the dual to \mathbf{Reg} is the functor $\mathbf{Max} : \mathbf{Esa} \rightarrow \mathbf{Stone}$, mapping an Esakia space (X, \leq) to its maximum. Here we present the right adjoint to this functor, also denoted by \mathbf{FreeM} , by borrowing key ideas from [1]. Essentially, this amounts to finding some free Heyting extension of a Boolean algebra which still has the same regular elements.

Given X a Stone space, let $(V(X), \supseteq)$ be the Priestley space of its closed subsets. The following is presumably folklore:

Proposition 1. *For each Stone space X , $(V(X), \supseteq)$ is an Esakia space.*

Our goal will be, given X an Esakia space, and Y a Stone space, to extend a function $f : \max X \rightarrow Y$ to a unique p-morphism $\tilde{f} : X \rightarrow \mathbf{FreeM}(Y)$. The first step is to note that $V(X)$ enjoys a universal property related to this:

Proposition 2. *Assume that X is an Esakia space, Y is a Stone space and $f : \max X \rightarrow Y$ is a continuous map. Then there exists a unique order-preserving map $\tilde{f} : X \rightarrow V(Y)$, such that $e_Y \circ \tilde{f} \upharpoonright_{\max} = f$ and \tilde{f} is a p-morphism on maximal elements (i.e., if $\tilde{f}(x) \leq y$ and $y \in V(Y)$ is maximal, then there is some $w \geq x$ such that $\tilde{f}(w) = y$).*

Date: January 10, 2026.

¹The present (2026) version corrects some imprecise statements made in the original abstract submitted to TBILLC. We thank Davide Quadrellaro and Matías Menni for calling our attention them.

Relatedly, one can consider a variation of the rooted Vietoris construction from [1]:

Definition 3. Given two Priestley spaces X, Y and a continuous and order-preserving map $g : X \rightarrow Y$ between them, we say that a subset $S \subseteq X$ is *g-open* if it satisfies:

$$\forall x \in S, y \in X (x \leq y \rightarrow \exists z \in S (x \leq z \wedge g(z) = g(y))).$$

We denote by $V_g(X)$ the set of closed, rooted and *g-open* subsets of X .

Note that if $Y = \{\bullet\}$, and g is the terminal map, $V_g(X)$ is the set of all closed and rooted subsets, which we denote by $V_r(X)$. Recall that there is a map called the *root map* $r : V_g(X) \rightarrow X$ which is a surjective order preserving map.

Proposition 4. *Let Y be a Stone space and let $V_{\max}(Y) = \{C \in V_r(V(Y)) : \forall D \in C, \forall x \in D, \{x\} \in C\}$. Then $V_{\max}(Y)$ is a Priestley space, and the restriction $r : V_{\max}(Y) \rightarrow V(Y)$ is such that for any map $f : \max X \rightarrow Y$, and its unique lifting $\tilde{f} : X \rightarrow V(Y)$ from Proposition 2, there is a unique r -open $g_f : X \rightarrow V_{\max}(Y)$ making the diagram commute.*

Let $M_\infty(Y) = V_G^r(V_{\max}(Y))$. The latter is constructed as follows: we consider the following sequence:

$$V(Y) \xleftarrow{r_1} V_{\max}(Y) \xleftarrow{r_2} V_2(Y) \xleftarrow{r_3} \dots$$

where $V_{n+1}(Y) = V_{r_n}(V_n(Y))$, as in [1], and $r_{n+1} : V_{n+1}(Y) \rightarrow V_n(Y)$ is the root map. Then $V_G^r(V_{\max}(Y))$ is the inverse limit of this sequence. $M_\infty(Y)$ is then an Esakia space, with the property that $\max(M_\infty(Y)) \cong Y$ through a natural isomorphism; moreover this assignment is functorial by using the functoriality of $V(-)$, $V_{\max}(-)$ and $V_G^r(-)$.

Our main result then shows the following adjunction:

Proposition 5. *The functor $\mathbf{FreeM} : \mathbf{Stone} \rightarrow \mathbf{Esa}$ assigning each Stone space X to $M_\infty(X)$ is right adjoint to $\max : \mathbf{Esa} \rightarrow \mathbf{Stone}$.*

2. REGULAR HEYTING ALGEBRAS

We use the description of \mathbf{FreeM} to study *regular Heyting algebras*, and their dual, regular Esakia spaces, in the sense of [4].

Definition 6. Let H be a Heyting algebra. We say that H is *regular* if $H = \langle \mathbf{Reg}(H) \rangle$. We say that an Esakia space X is *regular*, if its dual Heyting algebra is regular.

These are known to provide algebraic (respectively, order-topological) semantics for inquisitive logic [2]; indeed, given B a Boolean algebra, the Heyting algebra $\mathbf{ClopUp}(V(X_B))$ is studied there as the *inquisitive extension* of B . Using our construction, we show:

Theorem 7. *Given a Stone space X , $M_\infty(X)$ is always a regular Esakia space, and moreover, regular Esakia spaces are those spaces for which the unit of the adjunction is injective.*

This result can be seen as a dual algebraic semantics for inquisitive logic, and provides a categorical explanation for the role played by regular Heyting algebras in the study of such a logic. This furthermore answers a question of Grilleti and Quadrellaro[4], providing a categorical description of regular Esakia spaces. As an elaboration on the main ideas used to prove this theorem, we provide two theorems concerning regular Heyting algebras.

Definition 8. Given $n \in \omega$ the *n-universal regular model* if the (unique) poset (\mathcal{R}_n, \leq) satisfying the following:

- (1) $\max(P)$ contains 2^n points.
- (2) For each antichain $S \subseteq R_n$ where $|S| \geq 1$, there is a unique point $x \in P$ which covers S .

Compared with the usual n -universal model, the regular such model can be obtained by identifying all points with the same color, and hence it is a p-morphic image of the n -universal model. Then we can show:

Theorem 9. *Inquisitive logic InqL is sound and complete with respect to the class $\{\mathcal{R}_n : n \in \omega\}$.*

We conclude by tracing some connections and further avenues of research connected with the intermediate logic ML, Medvedev’s logic. This is well known to be the logic of the posets $V(X)$ for X a finite Stone space, i.e., a finite set, since $V(X)$ is exactly the powerset without the empty set. It likewise appears naturally in the study of inquisitive logic, as the *schematic fragment* of InqL. Here we prove:

Theorem 10. *The logic ML is precisely the logic of all the spaces $V(X)$ for X .*

We can phrase this result in more properly logical terms:

Definition 11. We define $R_n := \text{Log}(\{M_n(B) : B \text{ is a (finite) Boolean algebra}\})$,

where $M_n(B)$ is the n -th-step in the step-by-step construction outlined. We then have that $R_0 = \text{Med}$, and by [4], $\bigcap_{n \in \omega} R_n = \text{IPC}$. We leave it as an interesting avenue of research to obtain a further analysis of R_n for other n .

BIBLIOGRAPHY

- [1] Rodrigo Nicolau Almeida. *Colimits of Heyting Algebras through Esakia Duality*. 2024. DOI: 10.48550/ARXIV.2402.08058. URL: <https://arxiv.org/abs/2402.08058>.
- [2] Nick Bezhanishvili, Gianluca Grilletti, and Wesley H. Holliday. “Algebraic and Topological Semantics for Inquisitive Logic via Choice-Free Duality”. In: *Logic, Language, Information, and Computation*. Springer Berlin Heidelberg, 2019, pp. 35–52. ISBN: 9783662595336. DOI: 10.1007/978-3-662-59533-6_3. URL: http://dx.doi.org/10.1007/978-3-662-59533-6_3.
- [3] Leo Esakia. *Heyting Algebras: Duality Theory*. Ed. by Guram Bezhanishvili and Wesley Holliday. English translation of the original 1985 book. Springer, 2019.
- [4] Gianluca Grilletti and Davide Emilio Quadrellaro. “Esakia duals of regular Heyting algebras”. In: *Algebra universalis* 85.1 (Nov. 2023). ISSN: 1420-8911. DOI: 10.1007/s00012-023-00833-5. URL: <http://dx.doi.org/10.1007/s00012-023-00833-5>.
- [5] Tommaso Moraschini. “A Logical and Algebraic Characterisation of Adjunctions between Generalised Quasi-Varieties”. In: *The Journal of Symbolic Logic* 83.3 (2018), pp. 899–919.
- [6] Rodrigo Nicolau Almeida. “Polyatomic logics and generalized Blok–Esakia theory”. In: *Journal of Logic and Computation* 34.5 (June 2023), pp. 887–935. ISSN: 1465-363X. DOI: 10.1093/logcom/exad043. URL: <http://dx.doi.org/10.1093/logcom/exad043>.
- [7] Juan Soliveres Tur and Juan Climent Vidal. “Functors of Lindenbaum-Tarski, Schematic Interpretations, and Adjoint Cylinders between Sentential Logics”. In: *Notre Dame Journal of Formal Logic* 49.2 (2008).