

TOPICS IN ALGEBRAIC LOGIC AND DUALITY THEORY

Lecture 2

Rodrigo N. Almeida, Simon Lemal
June 5, 2025

- Beth definability property.
- Epimorphism surjectivity.

Beth definability

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, r)$ *implicitly defines* r if

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, r)$ *implicitly defines* r if

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, r)$ if

$$\Gamma(\bar{p}, r) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, r)$ *implicitly defines* r if

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, r)$ if

$$\Gamma(\bar{p}, r) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

The logic L has the *Beth property* if for any set of formulas $\Gamma(\bar{p}, r)$, if Γ implicitly defines r , then there is an explicit definition of r relative to Γ .

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, r)$ *implicitly defines* r if

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, r)$ if

$$\Gamma(\bar{p}, r) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

The logic L has the *Beth property* if for any set of formulas $\Gamma(\bar{p}, r)$, if Γ implicitly defines r , then there is an explicit definition of r relative to Γ .

Example

In CPC, consider $\Gamma(p_1, p_2, r) = \{r \rightarrow p_1, r \rightarrow p_2, p_1 \rightarrow (p_2 \rightarrow r)\}$.

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, r)$ *implicitly defines* r if

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, r)$ if

$$\Gamma(\bar{p}, r) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

The logic L has the *Beth property* if for any set of formulas $\Gamma(\bar{p}, r)$, if Γ implicitly defines r , then there is an explicit definition of r relative to Γ .

Example

In **CPC**, consider $\Gamma(p_1, p_2, r) = \{r \rightarrow p_1, r \rightarrow p_2, p_1 \rightarrow (p_2 \rightarrow r)\}$.
 Γ implicitly defines r .

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, r)$ *implicitly defines* r if

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, r)$ if

$$\Gamma(\bar{p}, r) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

The logic L has the *Beth property* if for any set of formulas $\Gamma(\bar{p}, r)$, if Γ implicitly defines r , then there is an explicit definition of r relative to Γ .

Example

In CPC, consider $\Gamma(p_1, p_2, r) = \{r \rightarrow p_1, r \rightarrow p_2, p_1 \rightarrow (p_2 \rightarrow r)\}$.

Γ implicitly defines r . $\phi_r = p_1 \wedge p_2$ is an explicit definition relative to Γ .

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, r)$ *implicitly defines* r if

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, r)$ if

$$\Gamma(\bar{p}, r) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

The logic L has the *Beth property* if for any set of formulas $\Gamma(\bar{p}, r)$, if Γ implicitly defines r , then there is an explicit definition of r relative to Γ .

Example

In **CPC**, consider $\Gamma(p_1, p_2, r) = \{r \rightarrow p_1, r \rightarrow p_2, p_1 \rightarrow (p_2 \rightarrow r)\}$.

Γ implicitly defines r . $\phi_r = p_1 \wedge p_2$ is an explicit definition relative to Γ .

However, in the implicative fragment of **CPC**, such an explicit is not possible.

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Proof.

Assume that Γ implicitly define r , that is,

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Proof.

Assume that Γ implicitly define r , that is,

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

By compactness, we may assume Γ to be finite, and taking conjunctions, we may assume that Γ is a single formula.

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Proof.

Assume that Γ implicitly define r , that is,

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

By compactness, we may assume Γ to be finite, and taking conjunctions, we may assume that Γ is a single formula.

Substituting r for r_1 and \top for r_2 , we obtain $\Gamma(\bar{p}, r), \Gamma(\bar{p}, \top) \vdash_L r$.

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Proof.

Assume that Γ implicitly define r , that is,

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

By compactness, we may assume Γ to be finite, and taking conjunctions, we may assume that Γ is a single formula.

Substituting r for r_1 and \top for r_2 , we obtain $\Gamma(\bar{p}, r), \Gamma(\bar{p}, \top) \vdash_L r$. By the deduction theorem,

$$\Gamma(\bar{p}, r) \vdash_L \Gamma(\bar{p}, \top) \rightarrow r.$$

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Proof.

Assume that Γ implicitly define r , that is,

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

By compactness, we may assume Γ to be finite, and taking conjunctions, we may assume that Γ is a single formula.

Substituting r for r_1 and \top for r_2 , we obtain $\Gamma(\bar{p}, r), \Gamma(\bar{p}, \top) \vdash_L r$. By the deduction theorem,

$$\Gamma(\bar{p}, r) \vdash_L \Gamma(\bar{p}, \top) \rightarrow r.$$

We also have $r \leftrightarrow \top \vdash_L \Gamma(\bar{p}, r) \leftrightarrow \Gamma(\bar{p}, \top)$.

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Proof.

Assume that Γ implicitly define r , that is,

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

By compactness, we may assume Γ to be finite, and taking conjunctions, we may assume that Γ is a single formula.

Substituting r for r_1 and \top for r_2 , we obtain $\Gamma(\bar{p}, r), \Gamma(\bar{p}, \top) \vdash_L r$. By the deduction theorem,

$$\Gamma(\bar{p}, r) \vdash_L \Gamma(\bar{p}, \top) \rightarrow r.$$

We also have $r \leftrightarrow \top \vdash_L \Gamma(\bar{p}, r) \leftrightarrow \Gamma(\bar{p}, \top)$. In particular,
 $r \vdash_L \Gamma(\bar{p}, r) \rightarrow \Gamma(\bar{p}, \top)$.

Beth property of intermediate logic

Theorem (Kreisel, 1960)

Any intermediate logic L has the Beth property.

Proof.

Assume that Γ implicitly define r , that is,

$$\Gamma(\bar{p}, r_1), \Gamma(\bar{p}, r_2) \vdash_L r_1 \leftrightarrow r_2.$$

By compactness, we may assume Γ to be finite, and taking conjunctions, we may assume that Γ is a single formula.

Substituting r for r_1 and \top for r_2 , we obtain $\Gamma(\bar{p}, r), \Gamma(\bar{p}, \top) \vdash_L r$. By the deduction theorem,

$$\Gamma(\bar{p}, r) \vdash_L \Gamma(\bar{p}, \top) \rightarrow r.$$

We also have $r \leftrightarrow \top \vdash_L \Gamma(\bar{p}, r) \leftrightarrow \Gamma(\bar{p}, \top)$. In particular, $r \vdash_L \Gamma(\bar{p}, r) \rightarrow \Gamma(\bar{p}, \top)$. By the deduction theorem,

$$\Gamma(\bar{p}, r) \vdash_L r \rightarrow \Gamma(\bar{p}, \top).$$



Theorem (Maksimova, 1993)


Any normal extension of K4 has the Beth property.

Theorem (Maksimova, 1993)

Any normal extension of K4 has the Beth property.

Example

Let L be the logic of the frame below. Note that it is weakly transitive.




Theorem (Maksimova, 1993)

Any normal extension of K4 has the Beth property.

Example

Let L be the logic of the frame below. Note that it is weakly transitive.




Consider $\Gamma(r) = r \leftrightarrow \Box \neg r$.

Theorem (Maksimova, 1993)

Any normal extension of K4 has the Beth property.

Example

Let L be the logic of the frame below. Note that it is weakly transitive.




Consider $\Gamma(r) = r \leftrightarrow \Box \neg r$. The only valuation validating it is $r \mapsto \{b\}$, thus Γ implicitly defines r .

Theorem (Maksimova, 1993)

Any normal extension of K4 has the Beth property.

Example

Let L be the logic of the frame below. Note that it is weakly transitive.



Consider $\Gamma(r) = r \leftrightarrow \Box \neg r$. The only valuation validating it is $r \mapsto \{b\}$, thus Γ implicitly defines r .

If r were explicitly definable, it would be by a variable free formula. However, every variable free formula is equivalent to \top or \perp (by induction).

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, \bar{r})$ *implicitly defines* \bar{r} if for every corresponding pair $r_1 \in \bar{r}_1, r_2 \in \bar{r}_2$,

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, \bar{r})$ *implicitly defines* \bar{r} if for every corresponding pair $r_1 \in \bar{r}_1, r_2 \in \bar{r}_2$,

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, \bar{r})$ if

$$\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, \bar{r})$ *implicitly defines* \bar{r} if for every corresponding pair $r_1 \in \bar{r}_1, r_2 \in \bar{r}_2$,

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, \bar{r})$ if

$$\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

The logic L has the *infinitary Beth property* if for any set of formulas $\Gamma(\bar{p}, \bar{r})$ that implicitly defines \bar{r} , every $r \in \bar{r}$ has an explicit definition relative to Γ .

Infinitary Beth property

Definition

Let L be a logic.

We say that a set of formulas $\Gamma(\bar{p}, \bar{r})$ *implicitly defines* \bar{r} if for every corresponding pair $r_1 \in \bar{r}_1, r_2 \in \bar{r}_2$,

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

We say that a formula $\phi_r(\bar{p})$ is an *explicit definition* of r relative to $\Gamma(\bar{p}, \bar{r})$ if

$$\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r(\bar{p}).$$

The logic L has the *infinitary Beth property* if for any set of formulas $\Gamma(\bar{p}, \bar{r})$ that implicitly defines \bar{r} , every $r \in \bar{r}$ has an explicit definition relative to Γ .

Proposition

If a logic L has the infinitary Beth property, then it has the Beth property.

Definition

A logic L has the *local deduction property* if for each pair of formulas ϕ, ψ , there is a formula δ_ϕ in the language of ϕ such that for each formula ξ ,

1. $\xi, \phi \vdash_L \psi$ iff $\xi \vdash_L \delta_\phi \rightarrow \psi$,
2. $\phi, \delta_\phi \rightarrow \xi \vdash_L \xi$.

Definition

A logic L has the *local deduction property* if for each pair of formulas ϕ, ψ , there is a formula δ_ϕ in the language of ϕ such that for each formula ξ ,

1. $\xi, \phi \vdash_L \psi$ iff $\xi \vdash_L \delta_\phi \rightarrow \psi$,
2. $\phi, \delta_\phi \rightarrow \xi \vdash_L \xi$.

Example

Any intermediate logic L has a local deduction property, with δ_ϕ given by ϕ .

Local deduction property

Definition

A logic L has the *local deduction property* if for each pair of formulas ϕ, ψ , there is a formula δ_ϕ in the language of ϕ such that for each formula ξ ,

1. $\xi, \phi \vdash_L \psi$ iff $\xi \vdash_L \delta_\phi \rightarrow \psi$,
2. $\phi, \delta_\phi \rightarrow \xi \vdash_L \xi$.

Example

Any intermediate logic L has a local deduction property, with δ_ϕ given by ϕ .

Example

Any normal modal logic L has a local deduction property, with $\delta_\phi = \phi \wedge \Box\phi \wedge \cdots \wedge \Box^n\phi$, where n is the modal depth of ψ .

Local deduction property

Definition

A logic L has the *local deduction property* if for each pair of formulas ϕ, ψ , there is a formula δ_ϕ in the language of ϕ such that for each formula ξ ,

1. $\xi, \phi \vdash_L \psi$ iff $\xi \vdash_L \delta_\phi \rightarrow \psi$,
2. $\phi, \delta_\phi \rightarrow \xi \vdash_L \xi$.

Example

Any intermediate logic L has a local deduction property, with δ_ϕ given by ϕ .

Example

Any normal modal logic L has a local deduction property, with $\delta_\phi = \phi \wedge \Box\phi \wedge \cdots \wedge \Box^n\phi$, where n is the modal depth of ψ .

Proposition (Exercise)

Let L be a logic with the local deduction property. If L has the Craig interpolation property, then L has the deductive interpolation property.

Theorem

Let L be a (compact, conjunctive) logic with the local deduction property. If L has the Craig interpolation property, then L has the (infinitary) Beth property.

Theorem

Let L be a (compact, conjunctive) logic with the local deduction property. If L has the Craig interpolation property, then L has the (infinitary) Beth property.

Proof.

See seminar 2.



Epimorphism surjectivity

Definition

A map $h: A \rightarrow B$ is *epic* (or an *epimorphism*) if for all maps $f_1, f_2: B \rightarrow C$, we have $f_1 \circ h = f_2 \circ h$ implies $f_1 = f_2$.

Definition

A map $h: A \rightarrow B$ is *epic* (or an *epimorphism*) if for all maps $f_1, f_2: B \rightarrow C$, we have $f_1 \circ h = f_2 \circ h$ implies $f_1 = f_2$.

Intuitively, h covers enough of B so that any two maps agreeing on the range of h agree on B . In other words, every $b \in B$ is defined implicitly from A .

Epimorphisms and surjections

Definition

A map $h: A \rightarrow B$ is *epic* (or an *epimorphism*) if for all maps $f_1, f_2: B \rightarrow C$, we have $f_1 \circ h = f_2 \circ h$ implies $f_1 = f_2$.

Intuitively, h covers enough of B so that any two maps agreeing on the range of h agree on B . In other words, every $b \in B$ is defined implicitly from A .

Definition

A map $h: A \rightarrow B$ is *onto* (or a *surjection*) if for each $b \in B$, there is $a \in A$ such that $h(a) = b$.

Epimorphisms and surjections

Definition

A map $h: A \rightarrow B$ is *epic* (or an *epimorphism*) if for all maps $f_1, f_2: B \rightarrow C$, we have $f_1 \circ h = f_2 \circ h$ implies $f_1 = f_2$.

Intuitively, h covers enough of B so that any two maps agreeing on the range of h agree on B . In other words, every $b \in B$ is defined implicitly from A .

Definition

A map $h: A \rightarrow B$ is *onto* (or a *surjection*) if for each $b \in B$, there is $a \in A$ such that $h(a) = b$.

In other words, every $b \in B$ is defined explicitly from A .

Epimorphisms and surjections

Definition

A map $h: A \rightarrow B$ is *epic* (or an *epimorphism*) if for all maps $f_1, f_2: B \rightarrow C$, we have $f_1 \circ h = f_2 \circ h$ implies $f_1 = f_2$.

Intuitively, h covers enough of B so that any two maps agreeing on the range of h agree on B . In other words, every $b \in B$ is defined implicitly from A .

Definition

A map $h: A \rightarrow B$ is *onto* (or a *surjection*) if for each $b \in B$, there is $a \in A$ such that $h(a) = b$.

In other words, every $b \in B$ is defined explicitly from A .

Definition

In general, onto maps are epic. A variety of algebras is said to have the *epimorphism surjectivity property* if the converse is true (for its morphisms).

Epimorphisms and surjections

Definition

A map $h: A \rightarrow B$ is *epic* (or an *epimorphism*) if for all maps $f_1, f_2: B \rightarrow C$, we have $f_1 \circ h = f_2 \circ h$ implies $f_1 = f_2$.

Intuitively, h covers enough of B so that any two maps agreeing on the range of h agree on B . In other words, every $b \in B$ is defined implicitly from A .

Definition

A map $h: A \rightarrow B$ is *onto* (or a *surjection*) if for each $b \in B$, there is $a \in A$ such that $h(a) = b$.

In other words, every $b \in B$ is defined explicitly from A .

Definition

In general, onto maps are epic. A variety of algebras is said to have the *epimorphism surjectivity property* if the converse is true (for its morphisms).

Example

The varieties **BA**, **HA**, **MSL** and **Lat** have epimorphism surjectivity.

The variety **DL** doesn't, as is witnessed by the map $\mathbf{3} \rightarrow \mathbf{2} \times \mathbf{2}$.

Theorem (Hoogland, 2000)

Let L be a logic and K the variety of its algebras. Then L has the Beth property iff K has epimorphism surjectivity.

Beth definability and epimorphism surjectivity

Theorem (Hoogland, 2000)

Let L be a logic and K the variety of its algebras. Then L has the Beth property iff K has epimorphism surjectivity.

Proof.

First assume that K has epimorphism surjectivity, and let $\Gamma(\bar{p}, \bar{r})$ such that for every corresponding pair $r_1 \in \bar{r}_1, r_2 \in \bar{r}_2$,

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

Beth definability and epimorphism surjectivity

Theorem (Hoogland, 2000)

Let L be a logic and K the variety of its algebras. Then L has the Beth property iff K has epimorphism surjectivity.

Proof.

First assume that K has epimorphism surjectivity, and let $\Gamma(\bar{p}, \bar{r})$ such that for every corresponding pair $r_1 \in \bar{r}_1, r_2 \in \bar{r}_2$,

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

$$\begin{array}{ccc} F(\bar{p}) & & \\ \downarrow \iota & \searrow \pi \circ \iota & \\ F(\bar{p}, \bar{r}) & \xrightarrow{\pi} & F(\bar{p}, \bar{r})/\Gamma \end{array}$$

The map $\pi \circ \iota$ is epic.

Beth definability and epimorphism surjectivity

Theorem (Hoogland, 2000)

Let L be a logic and K the variety of its algebras. Then L has the Beth property iff K has epimorphism surjectivity.

Proof.

First assume that K has epimorphism surjectivity, and let $\Gamma(\bar{p}, \bar{r})$ such that for every corresponding pair $r_1 \in \bar{r}_1, r_2 \in \bar{r}_2$,

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

$$\begin{array}{ccc} F(\bar{p}) & & \\ \downarrow \iota & \searrow \pi \circ \iota & \\ F(\bar{p}, \bar{r}) & \xrightarrow{\pi} & F(\bar{p}, \bar{r})/\Gamma \end{array}$$

The map $\pi \circ \iota$ is epic. Take $h_1, h_2: F(\bar{p}, r)/\Gamma \rightarrow A$ such that $h_1 \circ \pi \circ \iota = h_2 \circ \pi \circ \iota$. Clearly h_1 and h_2 agree on \bar{p} , and it is sufficient to show $h_1(r) = h_2(r)$ for each $r \in \bar{r}$.



Proof.

We define a valuation

$$v: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow A$$

$$p \mapsto h_1(p) = h_2(p) \quad \text{for } p \in \bar{p}$$

$$r_1 \mapsto h_1(r) \quad \text{for } r_1 \in \bar{r}_1 \text{ corresponding to } r \in \bar{r}$$

$$r_2 \mapsto h_2(r) \quad \text{for } r_2 \in \bar{r}_2 \text{ corresponding to } r \in \bar{r}.$$

Proof.

We define a valuation

$$v: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow A$$

$$p \mapsto h_1(p) = h_2(p) \quad \text{for } p \in \bar{p}$$

$$r_1 \mapsto h_1(r) \quad \text{for } r_1 \in \bar{r}_1 \text{ corresponding to } r \in \bar{r}$$

$$r_2 \mapsto h_2(r) \quad \text{for } r_2 \in \bar{r}_2 \text{ corresponding to } r \in \bar{r}.$$

Observe that for $\gamma \in \Gamma$,

$$v(\gamma(\bar{p}, \bar{r}_1)) = \gamma(v(\bar{p}), v(\bar{r}_1)) = \gamma(h_1(\bar{p}), h_1(\bar{r})) = h_1(\gamma(\bar{p}, \bar{r})) = h_1(1) = 1.$$

Proof.

We define a valuation

$$v: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow A$$

$$p \mapsto h_1(p) = h_2(p) \quad \text{for } p \in \bar{p}$$

$$r_1 \mapsto h_1(r) \quad \text{for } r_1 \in \bar{r}_1 \text{ corresponding to } r \in \bar{r}$$

$$r_2 \mapsto h_2(r) \quad \text{for } r_2 \in \bar{r}_2 \text{ corresponding to } r \in \bar{r}.$$

Observe that for $\gamma \in \Gamma$,

$$v(\gamma(\bar{p}, \bar{r}_1)) = \gamma(v(\bar{p}), v(\bar{r}_1)) = \gamma(h_1(\bar{p}), h_1(\bar{r})) = h_1(\gamma(\bar{p}, \bar{r})) = h_1(1) = 1.$$

Therefore v sends $\Gamma(\bar{p}, \bar{r}_1)$ to 1.

Proof.

We define a valuation

$$v: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow A$$

$$p \mapsto h_1(p) = h_2(p) \quad \text{for } p \in \bar{p}$$

$$r_1 \mapsto h_1(r) \quad \text{for } r_1 \in \bar{r}_1 \text{ corresponding to } r \in \bar{r}$$

$$r_2 \mapsto h_2(r) \quad \text{for } r_2 \in \bar{r}_2 \text{ corresponding to } r \in \bar{r}.$$

Observe that for $\gamma \in \Gamma$,

$$v(\gamma(\bar{p}, \bar{r}_1)) = \gamma(v(\bar{p}), v(\bar{r}_1)) = \gamma(h_1(\bar{p}), h_1(\bar{r})) = h_1(\gamma(\bar{p}, \bar{r})) = h_1(1) = 1.$$

Therefore v sends $\Gamma(\bar{p}, \bar{r}_1)$ to 1. Similarly for $\Gamma(\bar{p}, \bar{r}_2)$.

Proof.

We define a valuation

$$v: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow A$$

$$p \mapsto h_1(p) = h_2(p) \quad \text{for } p \in \bar{p}$$

$$r_1 \mapsto h_1(r) \quad \text{for } r_1 \in \bar{r}_1 \text{ corresponding to } r \in \bar{r}$$

$$r_2 \mapsto h_2(r) \quad \text{for } r_2 \in \bar{r}_2 \text{ corresponding to } r \in \bar{r}.$$

Observe that for $\gamma \in \Gamma$,

$$v(\gamma(\bar{p}, \bar{r}_1)) = \gamma(v(\bar{p}), v(\bar{r}_1)) = \gamma(h_1(\bar{p}), h_1(\bar{r})) = h_1(\gamma(\bar{p}, \bar{r})) = h_1(1) = 1.$$

Therefore v sends $\Gamma(\bar{p}, \bar{r}_1)$ to 1. Similarly for $\Gamma(\bar{p}, \bar{r}_2)$. Therefore, as Γ implicitly defines \bar{r} , every $r_1 \leftrightarrow r_2$ for corresponding r_1, r_2 is sent to 1, thus

$$h_1(r) = v(r_1) = v(r_2) = h_2(r).$$

Proof.

We define a valuation

$$v: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow A$$

$$p \mapsto h_1(p) = h_2(p) \quad \text{for } p \in \bar{p}$$

$$r_1 \mapsto h_1(r) \quad \text{for } r_1 \in \bar{r}_1 \text{ corresponding to } r \in \bar{r}$$

$$r_2 \mapsto h_2(r) \quad \text{for } r_2 \in \bar{r}_2 \text{ corresponding to } r \in \bar{r}.$$

Observe that for $\gamma \in \Gamma$,

$$v(\gamma(\bar{p}, \bar{r}_1)) = \gamma(v(\bar{p}), v(\bar{r}_1)) = \gamma(h_1(\bar{p}), h_1(\bar{r})) = h_1(\gamma(\bar{p}, \bar{r})) = h_1(1) = 1.$$

Therefore v sends $\Gamma(\bar{p}, \bar{r}_1)$ to 1. Similarly for $\Gamma(\bar{p}, \bar{r}_2)$. Therefore, as Γ implicitly defines \bar{r} , every $r_1 \leftrightarrow r_2$ for corresponding r_1, r_2 is sent to 1, thus

$$h_1(r) = v(r_1) = v(r_2) = h_2(r).$$

This proves that $h_1 = h_2$, thus $\pi \circ \iota$ is epic. □

Proof.

As $\pi \circ \iota$ is epic, it is surjective.

Proof.

As $\pi \circ \iota$ is epic, it is surjective. Thus there is $\phi_r \in F(\bar{p})$ such that $\pi \circ \iota(\phi_r) = r$ for all $r \in \bar{r}$.

Proof.

As $\pi \circ \iota$ is epic, it is surjective. Thus there is $\phi_r \in F(\bar{p})$ such that $\pi \circ \iota(\phi_r) = r$ for all $r \in \bar{r}$. We have $\phi_r = r$ in $F(\bar{p}, \bar{r})/\Gamma$ iff $\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r$.

Proof.

As $\pi \circ \iota$ is epic, it is surjective. Thus there is $\phi_r \in F(\bar{p})$ such that $\pi \circ \iota(\phi_r) = r$ for all $r \in \bar{r}$. We have $\phi_r = r$ in $F(\bar{p}, \bar{r})/\Gamma$ iff $\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r$. Thus L has the Beth property.

Proof.

As $\pi \circ \iota$ is epic, it is surjective. Thus there is $\phi_r \in F(\bar{p})$ such that $\pi \circ \iota(\phi_r) = r$ for all $r \in \bar{r}$. We have $\phi_r = r$ in $F(\bar{p}, \bar{r})/\Gamma$ iff $\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r$. Thus L has the Beth property.

Now assume that L has the Beth property, and let us show that K has epimorphism surjectivity.

Proof.

As $\pi \circ \iota$ is epic, it is surjective. Thus there is $\phi_r \in F(\bar{p})$ such that $\pi \circ \iota(\phi_r) = r$ for all $r \in \bar{r}$. We have $\phi_r = r$ in $F(\bar{p}, \bar{r})/\Gamma$ iff $\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r$. Thus L has the Beth property.

Now assume that L has the Beth property, and let us show that K has epimorphism surjectivity. Take an epimorphism $h: A \rightarrow B$.

Proof.

As $\pi \circ \iota$ is epic, it is surjective. Thus there is $\phi_r \in F(\bar{p})$ such that $\pi \circ \iota(\phi_r) = r$ for all $r \in \bar{r}$. We have $\phi_r = r$ in $F(\bar{p}, \bar{r})/\Gamma$ iff $\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r$. Thus L has the Beth property.

Now assume that L has the Beth property, and let us show that K has epimorphism surjectivity. Take an epimorphism $h: A \rightarrow B$. By the first isomorphism theorem, we have.

$$\begin{array}{ccc} A & & \\ \pi \downarrow & \searrow h & \\ A/\ker h \cong h(A) & \xrightarrow{\iota} & B \end{array}$$

h is epic iff ι is, and h is onto iff ι is.

Proof.

As $\pi \circ \iota$ is epic, it is surjective. Thus there is $\phi_r \in F(\bar{p})$ such that $\pi \circ \iota(\phi_r) = r$ for all $r \in \bar{r}$. We have $\phi_r = r$ in $F(\bar{p}, \bar{r})/\Gamma$ iff $\Gamma(\bar{p}, \bar{r}) \vdash_L r \leftrightarrow \phi_r$. Thus L has the Beth property.

Now assume that L has the Beth property, and let us show that K has epimorphism surjectivity. Take an epimorphism $h: A \rightarrow B$. By the first isomorphism theorem, we have.

$$\begin{array}{ccc} A & & \\ \pi \downarrow & \searrow h & \\ A/\ker h \cong h(A) & \xrightarrow{\iota} & B \end{array}$$

h is epic iff ι is, and h is onto iff ι is. Therefore, we can restrict our investigation of epimorphisms to epic inclusions. □

Proof.

Let A be a subalgebra of B such that the inclusion is epic.

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$.

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$. We claim that Γ implicitly defines \bar{r} .

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$. We claim that Γ implicitly defines \bar{r} .

Indeed, consider a map $h: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow C$ such that $\Gamma(\bar{p}, \bar{r}_1)$ and $\Gamma(\bar{p}, \bar{r}_2)$ are sent to 1.

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$. We claim that Γ implicitly defines \bar{r} .

Indeed, consider a map $h: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow C$ such that $\Gamma(\bar{p}, \bar{r}_1)$ and $\Gamma(\bar{p}, \bar{r}_2)$ are sent to 1. This defines two maps $h_1, h_2: F(\bar{p}, \bar{r}) \rightarrow C$ (with $h_1(p) = h_2(p) = h(p)$, $h_1(r) = h(r_1)$ and $h_2(r) = h(r_2)$).

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$. We claim that Γ implicitly defines \bar{r} .

Indeed, consider a map $h: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow C$ such that $\Gamma(\bar{p}, \bar{r}_1)$ and $\Gamma(\bar{p}, \bar{r}_2)$ are sent to 1. This defines two maps $h_1, h_2: F(\bar{p}, \bar{r}) \rightarrow C$ (with $h_1(p) = h_2(p) = h(p)$, $h_1(r) = h(r_1)$ and $h_2(r) = h(r_2)$).

Those two maps identify Γ with 1, hence they factor through π by definition of Γ .

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$. We claim that Γ implicitly defines \bar{r} .

Indeed, consider a map $h: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow C$ such that $\Gamma(\bar{p}, \bar{r}_1)$ and $\Gamma(\bar{p}, \bar{r}_2)$ are sent to 1. This defines two maps $h_1, h_2: F(\bar{p}, \bar{r}) \rightarrow C$ (with $h_1(p) = h_2(p) = h(p)$, $h_1(r) = h(r_1)$ and $h_2(r) = h(r_2)$).

Those two maps identify Γ with 1, hence they factor through π by definition of Γ . As they agree on A , and the inclusion $A \rightarrow B$ is epic, we must have $h_1 = h_2$.

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$. We claim that Γ implicitly defines \bar{r} .

Indeed, consider a map $h: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow C$ such that $\Gamma(\bar{p}, \bar{r}_1)$ and $\Gamma(\bar{p}, \bar{r}_2)$ are sent to 1. This defines two maps $h_1, h_2: F(\bar{p}, \bar{r}) \rightarrow C$ (with $h_1(p) = h_2(p) = h(p)$, $h_1(r) = h(r_1)$ and $h_2(r) = h(r_2)$).

Those two maps identify Γ with 1, hence they factor through π by definition of Γ . As they agree on A , and the inclusion $A \rightarrow B$ is epic, we must have $h_1 = h_2$. Therefore $h(r_1) = h_1(r) = h_2(r) = h(r_2)$ for any corresponding r, r_1, r_2 , and

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

Proof.

Let A be a subalgebra of B such that the inclusion is epic. Let $\bar{p} = A$, $\bar{r} = B \setminus A$ and consider the diagram below.

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \uparrow & & \uparrow \pi \\ F(\bar{p}) & \hookrightarrow & F(\bar{p}, \bar{r}) \end{array}$$

Let $\Gamma = \pi^{-1}(\{1\})$. We claim that Γ implicitly defines \bar{r} .

Indeed, consider a map $h: F(\bar{p}, \bar{r}_1, \bar{r}_2) \rightarrow C$ such that $\Gamma(\bar{p}, \bar{r}_1)$ and $\Gamma(\bar{p}, \bar{r}_2)$ are sent to 1. This defines two maps $h_1, h_2: F(\bar{p}, \bar{r}) \rightarrow C$ (with $h_1(p) = h_2(p) = h(p)$, $h_1(r) = h(r_1)$ and $h_2(r) = h(r_2)$).

Those two maps identify Γ with 1, hence they factor through π by definition of Γ . As they agree on A , and the inclusion $A \rightarrow B$ is epic, we must have $h_1 = h_2$. Therefore $h(r_1) = h_1(r) = h_2(r) = h(r_2)$ for any corresponding r, r_1, r_2 , and

$$\Gamma(\bar{p}, \bar{r}_1), \Gamma(\bar{p}, \bar{r}_2) \vdash_L r_1 \leftrightarrow r_2.$$

This prove that Γ implicitly defines \bar{r} . □

Proof.

As Γ implicitly define \bar{r} , it must also explicitly define it.

Proof.

As Γ implicitly define \bar{r} , it must also explicitly define it. Therefore, for any $r \in \bar{r}$ (or any $b \in B \setminus A$), there is $\phi_r \in F(\bar{p})$ that explicitly defines r relative to Γ . From

$$\Gamma(\bar{p}, \bar{r}) \vdash r \leftrightarrow \phi_r,$$

we deduce that $r = \phi_r$ in B (since B validates Γ).

Proof.

As Γ implicitly define \bar{r} , it must also explicitly define it. Therefore, for any $r \in \bar{r}$ (or any $b \in B \setminus A$), there is $\phi_r \in F(\bar{p})$ that explicitly defines r relative to Γ . From

$$\Gamma(\bar{p}, \bar{r}) \vdash r \leftrightarrow \phi_r,$$

we deduce that $r = \phi_r$ in B (since B validates Γ). Now remember that $\bar{r} = B \setminus A$, so for any $b \in B \setminus A$, there is $\phi_b \in F(\bar{p})$ such that $\phi_b = b$ in B .

Proof.

As Γ implicitly define \bar{r} , it must also explicitly define it. Therefore, for any $r \in \bar{r}$ (or any $b \in B \setminus A$), there is $\phi_r \in F(\bar{p})$ that explicitly defines r relative to Γ . From

$$\Gamma(\bar{p}, \bar{r}) \vdash r \leftrightarrow \phi_r,$$

we deduce that $r = \phi_r$ in B (since B validates Γ). Now remember that $\bar{r} = B \setminus A$, so for any $b \in B \setminus A$, there is $\phi_b \in F(\bar{p})$ such that $\phi_b = b$ in B . As $\bar{p} = A$, this ϕ_b is an element of A , and we are done. \square

Thank you!
Questions?