

TOPICS IN ALGEBRAIC LOGIC AND DUALITY THEORY

Lecture 4

Rodrigo N. Almeida, Simon Lemal

June 13, 2025

- Maksimova's characterisation of superintuitionistic logics with the interpolation property

A superintuitionistic logic has the interpolation property iff it is one of

CPC,

$LC_2 = IPC + bd_2 + bw_1$ = logic of the 2-chain,

$LC = IPC + p \rightarrow q \vee q \rightarrow p$ = logic of chains,

$KC = IPC + \neg p \vee \neg\neg p$ = logic of directed frames,

$BD_2 = IPC + bd_2$ = logic of frames of depth at most 2,

$BD_2W_2 = BD_2 + bw_2$ = logic of the 2-fork,

IPC.

Understanding the statement

Definition

We define a sequence of formulas \mathbf{bd}_n inductively:

$$\begin{aligned}\mathbf{bd}_0 &= \perp, \\ \mathbf{bd}_{n+1} &= p_{n+1} \vee (p_{n+1} \rightarrow \mathbf{bd}_n),\end{aligned}$$

Definition

We define a sequence of formulas \mathbf{bd}_n inductively:

$$\begin{aligned}\mathbf{bd}_0 &= \perp, \\ \mathbf{bd}_{n+1} &= p_{n+1} \vee (p_{n+1} \rightarrow \mathbf{bd}_n),\end{aligned}$$

Proposition

A frame validates \mathbf{bd}_n iff it doesn't contain a chain of $n + 1$ points, i.e. iff its depth is bounded by n .

Definition

We define a sequence of formulas \mathbf{bd}_n inductively:

$$\begin{aligned}\mathbf{bd}_0 &= \perp, \\ \mathbf{bd}_{n+1} &= p_{n+1} \vee (p_{n+1} \rightarrow \mathbf{bd}_n),\end{aligned}$$

Proposition

A frame validates \mathbf{bd}_n iff it doesn't contain a chain of $n + 1$ points, i.e. iff its depth is bounded by n .

Proof.

Exercise. □

Definition

We define formulas \mathbf{bw}_n as follows:

$$\mathbf{bw}_n = \bigvee_{i=0}^n \left(p_i \rightarrow \bigvee_{j \neq i} p_j \right).$$

Definition

We define formulas \mathbf{bw}_n as follows:

$$\mathbf{bw}_n = \bigvee_{i=0}^n \left(p_i \rightarrow \bigvee_{j \neq i} p_j \right).$$

Proposition

A frame validates \mathbf{bw}_n iff none of its rooted subframes contain an antichain of $n + 1$ points, i.e. iff its width is bounded by n .

Definition

We define formulas \mathbf{bw}_n as follows:

$$\mathbf{bw}_n = \bigvee_{i=0}^n \left(p_i \rightarrow \bigvee_{j \neq i} p_j \right).$$

Proposition

A frame validates \mathbf{bw}_n iff none of its rooted subframes contain an antichain of $n + 1$ points, i.e. iff its width is bounded by n .

Proof.

Exercise. □

Preliminary tools

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Corollary

Let F be a finite rooted frame and L be a logic. We have $\chi(F) \notin L$ iff $F \models L$.

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Corollary

Let F be a finite rooted frame and L be a logic. We have $\chi(F) \notin L$ iff $F \models L$.

Proof.

Exercise. □

Theorem (Jankov)

Let F be a finite rooted frame (i.e. the dual of a finite SI algebra). Then there is a formula $\chi(F)$ such that for any frame G , we have $G \not\models \chi(F)$ iff F is a p -morphic image of a subframe of G .

Corollary

Let F be a finite rooted frame and L be a logic. We have $\chi(F) \notin L$ iff $F \models L$.

Proof.

Exercise. □

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Definition

We turn the class of finite rooted frames into a poset by setting $F \leq G$ iff G is a p-morphic image of a generated subframe of F .

Definition

We turn the class of finite rooted frames into a poset by setting $F \leqslant G$ iff G is a p-morphic image of a generated subframe of F .

Definition

A frame F covers a frame G iff $F \leqslant G$ and there is no frame H such that $F < H < G$.

Definition

We turn the class of finite rooted frames into a poset by setting $F \leq G$ iff G is a p-morphic image of a generated subframe of F .

Definition

A frame F covers a frame G iff $F \leq G$ and there is no frame H such that $F < H < G$.

Proposition

Let F be a finite rooted frame. Then

$$\text{Log}(F) = \text{IPC} + \{\chi(G) \mid G \not\geq F\}.$$

See board.

Maksimova's characterisation

\Leftarrow : Exercise.

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation.

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \text{CPC}$, we are done.

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \mathbf{CPC}$, we are done. Otherwise, observe that \mathbf{CPC} is the logic of the 1-frame, whose only cover is the 2-chain.



(a) The 1-frame



(b) The 2-chain

Proof of Maksimova's characterisation

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \mathbf{CPC}$, we are done. Otherwise, observe that \mathbf{CPC} is the logic of the 1-frame, whose only cover is the 2-chain.



(a) The 1-frame



(b) The 2-chain

Therefore \mathbf{CPC} is axiomatised by the Jankov formula χ_{2c} of the 2-chain.

Proof of Maksimova's characterisation

\Leftarrow : Exercise.

\Rightarrow : Assume that L is a superintuitionistic with interpolation. If $L = \mathbf{CPC}$, we are done. Otherwise, observe that \mathbf{CPC} is the logic of the 1-frame, whose only cover is the 2-chain.



(a) The 1-frame



(b) The 2-chain

Therefore \mathbf{CPC} is axiomatised by the Jankov formula χ_{2c} of the 2-chain. As $L \subsetneq \mathbf{CPC}$, we have $\chi_{2c} \notin L$, thus the 2-chain is an L -frame.

If L is the logic of the 2-chain, then $L = \mathbf{LC}_2$, and we are done.

Otherwise, observe that the only covers of the 2-chain are the 3-chain and the 2-fork.



(a) 3-chain



(b) 2-fork

Otherwise, observe that the only covers of the 2-chain are the 3-chain and the 2-fork.



(a) 3-chain



(b) 2-fork

Therefore, \mathbf{LC}_2 is axiomatized by the Jankov formulas χ_{3c} of the 3-chain and χ_{2f} of the 2-fork.

Otherwise, observe that the only covers of the 2-chain are the 3-chain and the 2-fork.



(a) 3-chain



(b) 2-fork

Therefore, \mathbf{LC}_2 is axiomatized by the Jankov formulas χ_{3c} of the 3-chain and χ_{2f} of the 2-fork. As $L \subsetneq \mathbf{L}_2$, we have $\chi_{3c} \notin L$ or $\chi_{2f} \notin L$, thus either the 3-chain or the 2-fork are L -frames.

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame.

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame. Then $L \supseteq \text{BD}_2 = \text{IPC} + \chi_{3c}$.

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame. Then $L \supseteq \text{BD}_2 = \text{IPC} + \chi_{3c}$. What's more, the 2-fork is an L -frame, thus $L \subseteq \text{BD}_2\text{W}_2$.

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame. Then $L \supseteq \text{BD}_2 = \text{IPC} + \chi_{3c}$. What's more, the 2-fork is an L -frame, thus $L \subseteq \text{BD}_2\text{W}_2$. If $L = \text{BD}_2\text{W}_2$, we are done.

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame. Then $L \supseteq \text{BD}_2 = \text{IPC} + \chi_{3c}$. What's more, the 2-fork is an L -frame, thus $L \subseteq \text{BD}_2\text{W}_2$. If $L = \text{BD}_2\text{W}_2$, we are done. Otherwise, notice that the only cover of the 2-fork which does not cover the 3-chain is the 3-fork.



Figure 3: 3-fork

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame. Then $L \supseteq \mathbf{BD}_2 = \mathbf{IPC} + \chi_{3c}$. What's more, the 2-fork is an L -frame, thus $L \subseteq \mathbf{BD}_2\mathbf{W}_2$. If $L = \mathbf{BD}_2\mathbf{W}_2$, we are done. Otherwise, notice that the only cover of the 2-fork which does not cover the 3-chain is the 3-fork.



Figure 3: 3-fork

Therefore, \mathbf{BD}_2 is axiomatised relative to \mathbf{BD}_2 by the Jankov formula χ_{3f} of the 3-fork.

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame. Then $L \supseteq \text{BD}_2 = \text{IPC} + \chi_{3c}$. What's more, the 2-fork is an L -frame, thus $L \subseteq \text{BD}_2\text{W}_2$. If $L = \text{BD}_2\text{W}_2$, we are done. Otherwise, notice that the only cover of the 2-fork which does not cover the 3-chain is the 3-fork.



Figure 3: 3-fork

Therefore, BD_2 is axiomatised relative to BD_2 by the Jankov formula χ_{3f} of the 3-fork. As $\text{BD}_2 \subseteq L \subsetneq \text{BD}_2\text{W}_2$, we have $\chi_{3f} \notin L$, thus the 3-fork is an L -frame.

Assume that $\chi_{3c} \in L$, i.e. the 3-chain is not an L -frame. Then $L \supseteq \text{BD}_2 = \text{IPC} + \chi_{3c}$. What's more, the 2-fork is an L -frame, thus $L \subseteq \text{BD}_2\text{W}_2$. If $L = \text{BD}_2\text{W}_2$, we are done. Otherwise, notice that the only cover of the 2-fork which does not cover the 3-chain is the 3-fork.



Figure 3: 3-fork

Therefore, BD_2 is axiomatised relative to BD_2 by the Jankov formula χ_{3f} of the 3-fork. As $\text{BD}_2 \subseteq L \subsetneq \text{BD}_2\text{W}_2$, we have $\chi_{3f} \notin L$, thus the 3-fork is an L -frame. An amalgamation argument shows that every n -fork is an L -frame, thus $L = \text{BD}_2$.

Assume that both the 2-fork and the 3-chain are L -frames.

Assume that both the 2-fork and the 3-chain are L -frames. A non-trivial amalgamation trick shows that the 3-fork is an L -frame, thus all n -forks are L -frames.

Assume that both the 2-fork and the 3-chain are L -frames. A non-trivial amalgamation trick shows that the 3-fork is an L -frame, thus all n -forks are L -frames. An amalgamation argument show that every n -chain is an L -frame.

Assume that both the 2-fork and the 3-chain are L -frames. A non-trivial amalgamation trick shows that the 3-fork is an L -frame, thus all n -forks are L -frames. An amalgamation argument shows that every n -chain is an L -frame. Another non-trivial amalgamation trick shows that every n -ary tree is an L -frame, thus every finite tree is an L -frame.

Assume that both the 2-fork and the 3-chain are L -frames. A non-trivial amalgamation trick shows that the 3-fork is an L -frame, thus all n -forks are L -frames. An amalgamation argument shows that every n -chain is an L -frame. Another non-trivial amalgamation trick shows that every n -ary tree is an L -frame, thus every finite tree is an L -frame. As IPC is the logic of finite trees, we have $L = \text{IPC}$.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame. Then $L \supseteq \text{KC} = \text{IPC} + \chi_{2f}$.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame. Then $L \supseteq \mathbf{KC} = \mathbf{IPC} + \chi_{2f}$. What's more, the 3-chain is an L -frame, and an amalgamation argument shows that every n -chain is an L -frame.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame. Then $L \supseteq \mathbf{KC} = \mathbf{IPC} + \chi_{2f}$. What's more, the 3-chain is an L -frame, and an amalgamation argument shows that every n -chain is an L -frame. Thus $L \subseteq \mathbf{LC}$.

If $L = \mathbf{LC}$, we are done.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame. Then $L \supseteq \text{KC} = \text{IPC} + \chi_{2f}$. What's more, the 3-chain is an L -frame, and an amalgamation argument shows that every n -chain is an L -frame. Thus $L \subseteq \text{LC}$.

If $L = \text{LC}$, we are done. Otherwise, observe that LC is axiomatised relative to KC by the Jankov formula χ_d of the diamond, i.e. $\text{LC} = \text{KC} + \chi_d$.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame. Then $L \supseteq \text{KC} = \text{IPC} + \chi_{2f}$. What's more, the 3-chain is an L -frame, and an amalgamation argument shows that every n -chain is an L -frame. Thus $L \subseteq \text{LC}$.

If $L = \text{LC}$, we are done. Otherwise, observe that LC is axiomatised relative to KC by the Jankov formula χ_d of the diamond, i.e. $\text{LC} = \text{KC} + \chi_d$. As $\text{KC} \subseteq L \subsetneq \text{LC}$, we have $\chi_d \notin L$, so the diamond is an L -frame.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame. Then $L \supseteq \text{KC} = \text{IPC} + \chi_{2f}$. What's more, the 3-chain is an L -frame, and an amalgamation argument shows that every n -chain is an L -frame. Thus $L \subseteq \text{LC}$.

If $L = \text{LC}$, we are done. Otherwise, observe that LC is axiomatised relative to KC by the Jankov formula χ_d of the diamond, i.e. $\text{LC} = \text{KC} + \chi_d$. As $\text{KC} \subseteq L \subsetneq \text{LC}$, we have $\chi_d \notin L$, so the diamond is an L -frame.

The 3-chain and diamond are L -frames, we proceed as in the case of the 3-chain and the 2-fork, showing that every tree with a point on top is an L -frame.

First assume that $\chi_{2f} \in L$, i.e. the 2-fork is not an L -frame. Then $L \supseteq \text{KC} = \text{IPC} + \chi_{2f}$. What's more, the 3-chain is an L -frame, and an amalgamation argument shows that every n -chain is an L -frame. Thus $L \subseteq \text{LC}$.

If $L = \text{LC}$, we are done. Otherwise, observe that LC is axiomatised relative to KC by the Jankov formula χ_d of the diamond, i.e. $\text{LC} = \text{KC} + \chi_d$. As $\text{KC} \subseteq L \subsetneq \text{LC}$, we have $\chi_d \notin L$, so the diamond is an L -frame.

The 3-chain and diamond are L -frames, we proceed as in the case of the 3-chain and the 2-fork, showing that every tree with a point on top is an L -frame. This way we obtain every directed finite poset, so L must be KC .

Appendix

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite. If $X \models L$, then $\phi \notin L$.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite. If $X \models L$, then $\phi \notin L$. If $X \not\models L$, then $\chi(X) \in L'$, so $X \models \chi(X)$, which is absurd.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite. If $X \models L$, then $\phi \notin L$. If $X \not\models L$, then $\chi(X) \in L'$, so $X \models \chi(X)$, which is absurd.

If X is infinite, m -generated, it has infinite depth^a.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite. If $X \models L$, then $\phi \notin L$. If $X \not\models L$, then $\chi(X) \in L'$, so $X \models \chi(X)$, which is absurd.

If X is infinite, m -generated, it has infinite depth^a. Let k be the largest depth of an m -generated L model.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite. If $X \models L$, then $\phi \notin L$. If $X \not\models L$, then $\chi(X) \in L'$, so $X \models \chi(X)$, which is absurd.

If X is infinite, m -generated, it has infinite depth^a. Let k be the largest depth of an m -generated L model. Take some $x \in X$ at depth $k + 1$ and let Y be the subframe generated by x .

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite. If $X \models L$, then $\phi \notin L$. If $X \not\models L$, then $\chi(X) \in L'$, so $X \models \chi(X)$, which is absurd.

If X is infinite, m -generated, it has infinite depth^a. Let k be the largest depth of an m -generated L model. Take some $x \in X$ at depth $k + 1$ and let Y be the subframe generated by x . Then $Y \not\models L$, so $\chi(Y) \in L'$.

Theorem

A locally finite logic L can be axiomatised by the Jankov formulas

$$\{\chi(F) \mid F \not\models L\}.$$

Proof.

Let $L' = \text{IPC} + \{\chi(P) \mid P \not\models L\}$. We claim that $L = L'$.

\supseteq : Exercise.

\subseteq : Suppose $\phi \notin L'$. Let A be a subdirectly irreducible finitely generated model of L' which refutes ϕ and let X be its dual.

Suppose that X is finite. If $X \models L$, then $\phi \notin L$. If $X \not\models L$, then $\chi(X) \in L'$, so $X \models \chi(X)$, which is absurd.

If X is infinite, m -generated, it has infinite depth^a. Let k be the largest depth of an m -generated L model. Take some $x \in X$ at depth $k + 1$ and let Y be the subframe generated by x . Then $Y \not\models L$, so $\chi(Y) \in L'$. As Y is an L' -frame, we have $Y \models \chi(Y)$, which is a contradiction. □

^aThe curious student should refer to Nick's thesis, sections 3.1 & 3.2

Thank you!
Questions?