

The Intracellular Life Cycle of *Brucella* spp.

JEAN CELLI

Paul G. Allen School for Global Animal Health, Washington State University, Pullman, WA 99164

ABSTRACT Bacteria of the genus *Brucella* colonize a wide variety of mammalian hosts, in which their infectious cycle and ability to cause disease predominantly rely on an intracellular lifestyle within phagocytes. Upon entry into host cells, *Brucella* organisms undergo a complex, multistage intracellular cycle in which they sequentially traffic through, and exploit functions of, the endocytic, secretory, and autophagic compartments via type IV secretion system (T4SS)-mediated delivery of bacterial effectors. These effectors modulate an array of host functions and machineries to first promote conversion of the initial endosome-like *Brucella*-containing vacuole (eBCV) into a replication-permissive organelle derived from the host endoplasmic reticulum (rBCV) and then to an autophagy-related vacuole (aBCV) that mediates bacterial egress. Here we detail and discuss our current knowledge of cellular and molecular events of the *Brucella* intracellular cycle. We discuss the importance of the endosomal stage in determining T4SS competency, the roles of autophagy in rBCV biogenesis and aBCV formation, and T4SS-driven mechanisms of modulation of host secretory traffic in rBCV biogenesis and bacterial egress.

THE *BRUCELLA* INTRACELLULAR CYCLE

Brucella Pathogenesis

Bacteria of the genus *Brucella* belong to the alpha-2 subgroup of *Alphaproteobacteria*, a phylogenetic subgroup which includes a variety of bacteria that are either animal or plant pathogens or symbionts. As such, these bacteria have experienced a long-standing coevolution with eukaryotic hosts that has likely shaped their biology. The genus *Brucella* is composed of an increasing number of species that infect a wide variety of mammals as primary hosts, such as bovines (*Brucella abortus*), goats (*B. melitensis*), swine (*B. suis*), ovines (*B. ovis*), camels, elk, bison (*B. abortus*), canines (*B. canis*), rodents (*B. neotomae* and *B. microti*), and monkeys (*B. papionis*), as well as marine mammals such as seals, porpoises, dol-

phins, and whales (*B. pinnipedialis* and *B. ceti*) and also amphibians (*B. inopinata*) (1). Most species cause in their hosts a disease named brucellosis, which manifests as abortion, sterility, and lameness in animals and which can also be transmitted to humans via inhalation of aerosolized bacteria or via ingestion of, or contact with, contaminated tissues or derived products, classically by the most pathogenic species, *B. melitensis*, *B. suis*, and *B. abortus*, with additional cases due to *B. canis* and *B. neotomae* (2–4). Human brucellosis is characterized by nonspecific flu-like symptoms during an early acute phase, which is followed by a chronic infection with debilitating consequences, including recurrent fever, osteomyelitis, arthritis, neurological symptoms, and endocarditis, if not treated with antibiotic therapy in a timely manner (4). Animal and human brucellosis share common pathophysiological features at the cellular level, where bacteria undergo an intracellular cycle that ensures their survival, proliferation, and persistence within phagocytic cells of various tissues, including macrophages and dendritic cells (4, 5). Initially described in placental tissues of infected animals (6), the ability of *B. abortus* to extensively proliferate in mammalian cells was reproduced in a variety of tissue culture models of epithelial and phagocytic cells that have been instrumental in defining the main features of the bacterium's intracellular cycle (7–12).

Received: 27 March 2018, **Accepted:** 10 January 2019,
Published: 8 March 2019

Editors: Pascale Cossart, Institut Pasteur, Paris, France; Craig R. Roy, Yale University School of Medicine, New Haven, Connecticut; and Philippe Sansonetti, Institut Pasteur, Paris, France

Citation: Celli J. 2019. The intracellular life cycle of *Brucella* spp. *Microbiol Spectrum* 7(2):BAI-0006-2019. doi:10.1128/microbiolspec.BAI-0006-2019.

Correspondence: Jean Celli, jean.celli@wsu.edu

© 2019 American Society for Microbiology. All rights reserved.


A Multistage Intracellular Cycle

Upon entry into either phagocytic or nonphagocytic cells, *Brucella* bacteria are enclosed within a membrane-bound compartment that has been named the *Brucella*-containing vacuole (BCV) (8, 13) (Fig. 1), based on the concept that the original phagosome is functionally modified by the bacterium into an idiosyncratic membrane-bound vacuole. Early microscopy-based trafficking studies established that the initial BCV normally traffics along the endocytic pathway, sequentially acquiring early and

then late endosomal membrane markers (7, 8), and becomes acidified, reaching a pH of 4.5 (14–16), consistent with a complete phagosomal maturation process. Based on the endosomal nature of the bacterial vacuole at this stage of the intracellular cycle (between 0 and 8 h postinfection), it was named the endosomal BCV (eBCV) (Fig. 1 and 2A) (17).

Subsequent to eBCV interactions with the endocytic compartment, the vacuole progressively loses endosomal markers (between 8 and 12 h postinfection), concomitant

FIGURE 1 Model of the *Brucella* intracellular cycle in macrophages. Following phagocytic uptake by macrophages, *Brucella* spp. reside during the first 8 to 12 h postinfection within a membrane-bound vacuole that undergoes endosomal maturation via sequential interactions with early (EE) and late (LE) endosomes and lysosomes (LYS) to become an acidified eBCV. The host small GTPase Rab7 contributes to eBCV maturation, which provides physicochemical cues promoting expression of the VirB T4SS, which translocates effector proteins (red) that mediate eBCV interactions with the ER exit site and acquisition of ER and Golgi apparatus-derived membranes. These events lead to the biogenesis of replication permissive, ER-derived BCVs, called rBCVs. The host proteins Sar1, IRE1 α , Yip1A, Atg9, and WIP1 and the COG complex contribute to rBCV biogenesis. Bacteria then undergo extensive replication in rBCVs between 12 and 48 h postinfection, after which rBCVs are captured within autophagosome-like structures in a VirB T4SS-dependent manner to become aBCVs. aBCV formation requires the host autophagy proteins beclin1, ULK1, and Atg14. aBCVs harbor features of autolysosomes and are required for bacterial egress and new cycles of intracellular infections.






FIGURE 2 Structure and membrane composition of BCVs during the *Brucella* intracellular cycle. (A) Confocal fluorescence micrograph of HeLa cells expressing green fluorescent protein (GFP)-Rab7 and infected with DsRed_m-expressing *B. abortus* strain 2308 for 6 h. The inset shows an eBCV with the typical accumulation of the late endosomal/lysosomal markers Rab7 and LAMP1. Bars, 10 and 2 μ m. (B) Confocal fluorescence micrograph of a HeLa cell infected with DsRed_m-expressing *B. abortus* strain 2308 for 24 h and stained for the ER marker calreticulin. The inset shows a cluster of calreticulin-positive rBCVs containing replicating bacteria and associated with the ER network. Bars, 10 and 1 μ m. (C) Confocal fluorescence micrograph of a primary murine bone marrow-derived macrophage expressing the autophagy marker GFP-LC3 and infected with DsRed_m-expressing *B. abortus* strain 2308 for 72 h. The inset shows a group of aBCVs with the typical accumulation of the late endosomal/lysosomal LAMP1 but not LC3. Bars, 10 and 2 μ m. (D) Transmission electron micrographs of bone marrow-derived macrophages infected with *B. abortus* strain 2308 for 72 h and showing the ultrastructures of rBCVs (left, single-membrane-bound vacuoles [inset a]), of forming aBCVs (inset b and arrows), and of completed double-membrane-bound aBCVs (right [insets c and d and arrows]). Bars, 500 and 200 nm. Reprinted from reference 17 with permission.

with sustained interactions with endoplasmic reticulum (ER) structures, and eventually acquires ER membrane-associated markers such as calreticulin, calnexin, and Sec61 β (7, 8, 13), indicating a turnover of eBCV membranes and accretion of ER-derived membranes. These vacuoles also gain structural and functional features of the ER (7, 8), further indicating a conversion of the eBCV into an ER-derived organelle. These structural and functional changes correlate with the onset of bacterial replication, suggesting that these organelles provide intracellular conditions that promote bacterial growth and replication. Based on these considerations, these vacuoles

were named replicative BCVs (rBCVs) (Fig. 1 and 2B) (17). Original ultrastructural characterizations of rBCVs suggested that these vacuoles were individual and rarely connected with ER cisternae in either macrophage or HeLa cells (7, 8) (Fig. 2D), while other studies observed bacteria within the ER lumen of trophoblasts (6), leaving the actual organization of rBCVs uncertain. Recent electron tomography three-dimensional reconstruction of rBCVs clearly determined that these vacuoles are intricately connected with the ER and also seem to interact with vesicular traffic between the ER and the Golgi apparatus (18), demonstrating intimate interactions with

the host secretory pathway. The rBCV stage is tightly associated with bacterial proliferation (between 12 and 48 h postinfection) and causes a dramatic reorganization of the host cell ER into the rBCV network (8).

Following extensive bacterial replication, an additional stage in the *Brucella* intracellular cycle consists of the progressive capture of rBCVs by crescent-like membrane structures, leading to the formation of multimembrane vacuoles that are structurally reminiscent of autophagosomes that contain small to large groups of bacteria and acquire late endosomal/lysosomal features (17). These remodeled rBCVs, which are called autophagic BCVs (aBCVs) (Fig. 1 and 2C and D) form between 48 and 72 h postinfection and are associated with bacterial release from infected cells and new infection events, indicating that they facilitate completion of the *Brucella* intracellular cycle (17).

By their functional and spatial diversity, the sequential eBCV, rBCV, and aBCV stages of the *Brucella* intracellular cycle point towards their necessity to the bacterium's infectious cycle. Hence, significant efforts have been made in the last 2 decades to understand the underlying mechanisms that define these stages and how the bacterium exploits the corresponding cellular pathways to complete its intracellular cycle.

The VirB Type IV Secretion System

A hallmark of *Brucella* virulence is the expression of a VirB family type IVA secretion system (T4SS), which was identified by homology to the VirB T4SS of the plant pathogen *Agrobacterium tumefaciens* and shown to be required for virulence in a murine model of chronic brucellosis and intracellular replication of *B. abortus*, *B. melitensis*, and *B. suis* in various host cell models (19–21). Indeed, deletions of various genes within the *virB* operon, or transposon insertions in different *virB* genes, rendered *Brucella* unable to convert eBCVs into rBCVs and replicate intracellularly (8, 13, 22) and unable to establish a chronic infection in mice (23, 24), emphasizing major roles of this T4SS in rBCV biogenesis and *Brucella* intracellular replication. Based on the ability of various T4SSs to deliver effector proteins or nucleoprotein complexes across biological membranes, and by analogy to those that contribute to bacterial virulence (25, 26), the VirB T4SS of *Brucella* was presumed to deliver effector proteins into host cells across the eBCV membrane to modulate specific cellular functions and mediate rBCV biogenesis. This assumption was confirmed by the discoveries of an array of *Brucella* effector proteins translocated into host cells in a T4SS-dependent manner (27–32), which opened avenues to understand at

the molecular level the underlying mechanisms of the *Brucella* intracellular cycle.

THE eBCV

Early BCV Maturation

Upon uptake by phagocytes or entry into nonphagocytic cells, *Brucella* bacteria initially reside in a phagosome that rapidly acquires markers of the early endocytic compartment, including the small GTPase Rab5, which controls early endosomal maturation, and the early endosomal antigen EEA-1 (8, 13, 33). Since these markers are acquired with kinetics similar to those of classical phagosomes, it is likely that the early BCV undergoes an unaltered maturation process. Consistently, the vacuole then acquires markers of late endosomes, such as the lysosome-associated membrane proteins LAMP1 (Fig. 2A) and LAMP2, CD63, and the small GTPase Rab7 (7, 8, 13, 16), which controls fusion with late endocytic compartments and lysosomes, further indicating a normal vacuolar maturation process within the first hours postinfection. Importantly, the BCV also rapidly becomes acidified to a pH in the range of 4.5 to 5.0 (14), further supporting a maturation process into phagolysosomes. However, early attempts at detecting delivery of lysosomal luminal contents to the BCV as evidence for fusion with lysosomes failed to demonstrate accumulation of cathepsin D by immunostaining (7, 13, 34). This led to the proposal that the BCV avoids fusion with terminal degradative lysosomes, despite advanced maturation events, thus preventing bacterial killing and ensuring intracellular survival.

The eBCV: A Necessary Evil?

Several studies monitoring bacterial viability within eBCVs have revealed that a large proportion of intracellular bacteria (up to 90% in primary bone marrow-derived macrophages) are killed (8, 16), arguing for bactericidal conditions within eBCVs. Using live-cell imaging of fluorescent fluid-phase markers that were chased to terminal lysosomes, as a method that precludes any issue with detection of soluble lysosomal antigens, Starr et al. showed that eBCVs in epithelial HeLa cells undergo fusion with terminal lysosomes, although to a restricted extent (16), directly demonstrating that the eBCV matures into a compartment resembling a phagolysosome. While these findings are consistent with the loss of bacterial viability in eBCVs observed in *in vitro* models, they may appear counterintuitive with regard to *Brucella* intracellular survival. This has led to the speculation that eBCVs that undergo full lysosomal maturation are not those permis-

sive for bacterial survival and represent dead-end paths for the bacteria, while a small fraction of BCVs that do not mature along the endocytic pathway are those that generate rBCVs and allow proliferation of the surviving bacteria. However, the following observations are inconsistent with this possibility. First, LAMP1-positive eBCVs interact with ER cisternae and acquire ER markers, arguing that they undergo conversion into rBCVs (8). Second, eBCV acidification to a pH range of terminal lysosomes is necessary for rBCV biogenesis and bacterial replication (14–16), as inhibition of lysosomal acidification prevents eBCV to rBCV conversion and bacterial replication (14–16). Third, inhibition of Rab7 activity, which controls fusion with lysosomes, also prevents eBCV conversion into rBCVs and bacterial replication (16), demonstrating that a functional late endocytic pathway and lysosomal fusion are required for *Brucella* to undergo its intracellular cycle. Fourth, eBCV acidification is essential for the induction of the *virB* operon (15, 16), which rapidly occurs postuptake in maturing eBCVs, peaking at 4 h postinfection (16, 35). Taken together, these findings rather support a model in which the eBCV stage is a necessary step in the *Brucella* intracellular cycle, which provides intravacuolar cues, including a lysosomal pH, that are necessary for the induction of the VirB T4SS and the resulting conversion of eBCVs to rBCVs.

In addition to its role in promoting *Brucella* T4SS competency, the eBCV may provide cues that initiate intracellular bacterial growth prior to completion of the replication-permissive rBCV. Deghelt et al. established using fluorescence microscopy methods that monitor chromosomal replication in intracellular *Brucella* that bacteria within LAMP1-positive eBCVs initiate chromosomal replication by 8 h postinfection, i.e., during the eBCV-to-rBCV conversion stage, while the infectious forms are in G₁-arrested phase prior to 6 h postinfection (36). Whether this cell cycle change is triggered by the endosomal conditions of the eBCV or by intravacuolar alterations during conversion to rBCV remains to be determined. However, these findings indicate that *Brucella* extensively responds to its intravacuolar environment and initiates growth at the eBCV stage, further emphasizing the importance of the endosomal stage in the *Brucella* intracellular cycle.

THE rBCV

Role of the ER in rBCV Biogenesis

Biogenesis of the rBCV is a hallmark of the *Brucella* intracellular cycle, as it consists of the conversion of the original endosomal bacterial vacuole into a specialized

organelle derived from the ER that promotes bacterial proliferation. The underlying mechanisms, host functions, and bacterial effectors that mediate this process have been the subject of extensive research. The nature of the rBCV was initially discovered via its ultrastructural characterization, showing fusion with ER cisternae and studding with ribosomes, and was further confirmed by the detection of ER-associated proteins on its membrane by immunofluorescence microscopy, intravacuolar detection of glucose-6 phosphatase activity, an ER luminal enzyme, and sensitivity to the ER-vacuolating toxin aerolysin (7, 8). The first clues about how a bacterial vacuole interacting with late endocytic and lysosomal compartments could convert into an ER-derived organelle came from the demonstration that ER exit sites (ERES), an ER subcompartment where secretory transport is initiated through the formation of COPII-coated cargo vesicles (37), are functionally important for rBCV biogenesis (22). eBCVs undergo T4SS-dependent, sustained interactions with COPII coat-positive ERES structures during the eBCV-to-rBCV conversion stage (Fig. 3), and inhibition of the small GTPase Sar1, which controls COPII coat assembly and ERES formation and function, inhibits rBCV biogenesis and bacterial replication (22, 38). Consistent with these observations, *Brucella* infection upregulates production of Sar1 and the COPII components Sec23 and Sec24D by an unknown mechanism (39), suggesting that the bacterium upregulates Sar1- and COPII-mediated vesicular trafficking to promote rBCV biogenesis. Hence, *Brucella* may modulate ERES functions to promote rBCV biogenesis, by enhancing production of secretory vesicles that might fuse with the eBCV to initiate its conversion into a vacuole with ER fusogenic properties (Fig. 3), possibly creating a direct port of entry to the ER for the bacteria, while bypassing classical endosome-to-Golgi-to-ER retrograde trafficking processes.

Role of the UPR in the *Brucella* Intracellular Cycle

Another ER-mediated function associated with the *Brucella* intracellular cycle is the unfolded-protein response (UPR). Upon ER stress triggered by accumulation of misfolded proteins within the ER, signaling pathways controlled by the ER-localized UPR receptors IRE1 α , PERK, and ATF6 are triggered and lead to a substantial reorganization of ER functions towards increased ER folding capacity, inhibition of protein synthesis, and activation of autophagy, all aimed at resolving ER stress (40). Although there have been disagreements about its extent, the consensus is that *Brucella* infection induces the UPR in macrophages or HeLa cells via at least the IRE1 α pathway




FIGURE 3 Model of VirB T4SS-dependent biogenesis of the rBCV. Bacteria in eBCVs induce expression of the VirB T4SS, which delivers effector proteins into the host cell. Among these, BspB traffics to Golgi membranes via the ER-to-Golgi intermediate compartment (ERGIC) and binds to the COG complex to promote redirection of Golgi apparatus-derived vesicular traffic to BCVs. RicA binds the small host secretory GTPase Rab2, which contributes to its recruitment on maturing eBCVs and role in rBCV biogenesis. Additionally, eBCV interaction with ERES is accompanied by the upregulation of COPII coat components, induction of IRE1 α , and Yip1A-dependent formation of ER-derived vesicles, which are also thought to contribute to rBCV biogenesis. T4SS-dependent acquisition of ER- and Golgi apparatus-derived secretory membranes by BCVs is thought to mediate eBCV-to-rBCV conversion.

(39, 41, 42). Moreover, IRE1 α is required for *Brucella* replication (43), and pharmacological compensation of protein misfolding using tauroursodeoxycholic acid impairs replication of *B. melitensis* (41), which supports the idea that the UPR is beneficial to the bacterium's intracellular cycle. However, whether and how this stress response promotes rBCV biogenesis and bacterial replication remain unclear. Taguchi et al. revealed a link between UPR induction, ERES, and rBCV biogenesis by showing that IRE1 α is activated via the ERES-localized protein Yip1A, which controls formation of large vacuoles in a manner dependent on the autophagy-associated proteins Atg9 and WIPI1, and is also required for rBCV biogenesis (39). Hence, these findings support a model in which *Brucella* actively causes UPR induction via acti-

vation of the IRE1 α pathway to promote an activation cascade leading to formation at ERES of vacuoles of a possible autophagic nature (43), which ultimately contributes to rBCV biogenesis (Fig. 3).

Is Autophagy Involved in rBCV Biogenesis?

Autophagy is a membrane-based process that normally captures intracellular contents, whether cytosolic contents, damaged organelles, or microorganisms, into double-membrane vesicles called autophagosomes, to deliver them for degradation to the lysosomal compartment. While it can act as an innate immune antibacterial mechanism, it can also be beneficial to some pathogens (44, 45). Based on the roles of IRE1 α and Yip1A in rBCV biogenesis and *Brucella* replication, autophagosome for-

mation at ERES may provide ER-derived membranes that may contribute to eBCV-to-rBCV conversion; however, whether the canonical autophagic cascade is involved in rBCV biogenesis remains controversial. While a classical autophagic process was originally proposed as a mechanism for rBCV biogenesis in HeLa cells, based on the observation of multimembrane structures on BCVs and accumulation of the lysosomotropic compound monodansylcadaverine (7, 34), such structures are not seen on eBCVs in macrophages (8), nor does the autophagic marker LC3 accumulate on eBCVs, arguing against a canonical autophagic process. Furthermore, inhibition of autophagy via small-interfering-RNA (siRNA)-mediated depletion of specific autophagy components involved in the autophagosome nucleation and elongation steps, namely, ULK1, beclin1, Atg5, Atg7, ATG16L, and LC3B, does not prevent rBCV biogenesis and bacterial replication (17). Hence, there is no substantial evidence to support a role of the canonical autophagic cascade in rBCV biogenesis. However, the demonstrated roles of the autophagosome nucleation protein WIPI1 and of the autophagy protein Atg9 in rBCV biogenesis (39) clearly invoke specific autophagy-associated machineries in this process, suggesting that rBCV biogenesis requires a subset of cellular machineries associated with autophagosome formation at ERES and Atg9-mediated delivery of membranes. Further studies are needed to clarify how these autophagy-related events actually contribute to rBCV biogenesis.

Role of the Secretory Pathway in rBCV Biogenesis

Based on the ER nature of the rBCV, the host secretory pathway certainly plays an essential role in the *Brucella* intracellular cycle. Whether secretory compartments other than the ER, such as the ER-to-Golgi intermediate compartment, the Golgi apparatus, and the *trans*-Golgi network, are important to the bacterium is a long-standing question, warranted by examples of other intracellular pathogens that exploit these compartments (46). These include *Legionella pneumophila*, which intercepts Arf1- and Rab1-dependent secretory traffic to acquire ER-derived membranes (47, 48), and *Chlamydia trachomatis*, which redirects Golgi apparatus-derived lipid trafficking pathways for acquisition of sphingolipids (49, 50). Cellular approaches based on altering specific functions along the secretory pathway using either pharmacological inhibition of host secretion, expression of dominant inactive alleles of small GTPases, or siRNA-mediated depletions have revealed that ER-Golgi secretory traffic is important for rBCV biogenesis

and bacterial replication (22, 38). Inhibition of Arf1-dependent secretory traffic using brefeldin A or via siRNA treatment impairs rBCV biogenesis to levels similar to those seen with depletion of Sar1 (22, 38). Additionally, depletion of the small GTPases Rab1a and Rab2, which control anterograde and retrograde vesicular traffic between the ER and Golgi apparatus (51), negatively affects replication of *B. abortus* in macrophages (38), further indicating a complex role of vesicular traffic between these compartments in promoting *Brucella* proliferation, possibly through acquisition by the eBCV of secretory membranes during their conversion process into the rBCV. Consistent with *Brucella* exploiting these pathways, host secretory traffic is altered in *Brucella*-infected cells in a T4SS-dependent manner (30), which argues that the bacterium actively modulates secretory functions via delivery of effectors to promote its intracellular cycle.

Bacterial Effectors of rBCV Biogenesis

The recent discoveries of several VirB T4SS effectors (27–32) and the characterization of the modes of action of some of them (Table 1) have provided opportunities to start deciphering at the molecular level the bacterial mechanisms of rBCV biogenesis and bacterial replication. VceC, identified as being coregulated with the VirB T4SS, binds the ER chaperone Grp78/BiP and induces the UPR, which triggers an inflammatory response (42), yet whether it plays a role in rBCV biogenesis has not been elucidated. TcpB/BtpA, a TIR-domain-containing effector that down-modulates proinflammatory responses (52–58), also induces the UPR and promotes bacterial proliferation once rBCVs are formed (41), suggesting that it may not contribute to rBCV biogenesis. SepA, in contrast, is possibly involved in rBCV biogenesis, as a Δ sepA mutant is retained in eBCVs (32), yet its mode of action remains unknown.

The T4SS effectors BspA, BspB, and BspF, which inhibit host protein secretion and promote bacterial replication (30, 38), support the hypothesis that *Brucella* specifically targets various host secretory functions, possibly for rBCV biogenesis purposes. While the modes of action of BspA and BspF are unknown, that of BspB has been elucidated. BspB is required for rBCV biogenesis and optimal bacterial replication in macrophages (38). This effector is delivered into host cells and traffics to the Golgi apparatus, where it interacts with the conserved oligomeric Golgi (COG) complex (38) (Fig. 3), a CATCHR-family multisubunit tethering complex that serves as an interaction hub on Golgi membranes for secretory Rab GTPases, Golgi tethers, and SNAREs and that regulates intra-Golgi and retrograde vesicular traffic

TABLE 1 *Brucella* T4SS effectors

Name	ORF in <i>B. abortus</i>	Host target	Function	Reference(s)
RicA	BAB1_1279	Rab2	Modulates rBCV biogenesis	28
VceA	BAB1_1652	Unknown	Unknown	27
VceC	BAB1_1058	Grp78/BiP	UPR activation	27 , 42
BspA	BAB1_0678	Unknown	Intracellular replication	30
BspB	BAB1_0712	COG	Redirects Golgi vesicular traffic; rBCV biogenesis; intracellular replication	30 , 38
BspC	BAB1_0847	Unknown	Unknown	30
BspE	BAB1_1675	Unknown	Unknown	30
BspF	BAB1_1948	Unknown	Intracellular replication	30
BtpA/TcpB	BAB1_0279	MAL	Inhibition of TLR signaling; UPR induction	52
BtpB	BAB1_0756	Unknown	Unknown	31
SepA	BAB1_1492	Unknown	eBCV trafficking	32
BPE005	BAB1_2005	Unknown	Unknown	29
BPE043	BAB1_1043	Unknown	Unknown	29
BPE275	BAB1_1275	Unknown	Unknown	29
BPE123	BAB1_0123	Unknown	Unknown	29

along the secretory pathway ([59](#)). COG functions are important for rBCV biogenesis and bacterial replication ([38](#)), implicating Golgi apparatus-associated functions in the *Brucella* intracellular cycle. By mechanisms that remain to be defined, BspB-COG interactions alter COG functions and lead to redirection of COG-dependent Golgi retrograde vesicular traffic to BCVs and acquisition of Golgi apparatus-derived membranes ([38](#)), demonstrating that *Brucella* likely recruits membranes from this secretory compartment during rBCV biogenesis, in addition to ER-derived membranes ([Fig. 3](#)). Interestingly, inhibition of Rab2-dependent Golgi-to-ER retrograde traffic via depletion of Rab2 suppresses the replication defect of a $\Delta bspB$ mutant ([38](#)), suggesting that BspB may affect retrograde secretory traffic to redirect COG-dependent Golgi vesicular traffic to the BCV. Interestingly, the T4SS effector RicA is involved in controlling rBCV biogenesis and binds the GDP-bound form of Rab2 ([28](#)) ([Fig. 3](#)). Although its mode of action is unknown, as it does not show any GEF (guanine nucleotide exchange factor) activity ([28](#)), this suggests that *Brucella* delivers several effectors that may coordinately act to modulate Rab2-dependent vesicular trafficking and promote rBCV biogenesis.

Altogether, the knowledge gained by studying VirB T4SS effectors that target the host secretory pathway has revealed yet another facet of *Brucella* interactions with this intracellular compartment and emphasizes how the identification and characterization of these effectors are key to a comprehensive understanding of how this bacterium subverts host secretory functions.

THE aBCV

While replication in ER-associated rBCVs is a key step in the pathogenesis of *Brucella*, how the bacterium com-

pletes its intracellular cycle following this proliferation stage has remained unknown for many decades. Unlike many pathogens, which cause cell death to exit the cells in which they have replicated, *Brucella* prevents cell death programs from being carried out ([60](#), [61](#)), thus preserving its intracellular niche. Starr et al. observed that instead, following proliferation in rBCVs, by 48 h postinfection and afterwards, the *Brucella* intracellular niche was converted from an ER-derived organelle to large vacuoles harboring features of late endosomal and lysosomal compartments, such as accumulation of LAMP1 and acidification ([17](#)). From an ultrastructural standpoint, these vacuoles are surrounded by multiple membranes and originate from the capture of rBCVs by crescent-shaped membrane structures reminiscent of autophagosomes, despite the lack of accumulation of the canonical autophagosome marker LC3 ([17](#)) ([Fig. 2](#)). The formation of these vacuoles, named aBCVs, requires functions of the canonical autophagy nucleation but not elongation complexes, as depletion or deletion of beclin1, ULK1, and Atg14L, but not that of Atg5, Atg7, Atg4, or Atg16L, blocked their formation ([17](#)). aBCV formation therefore seems to require a subset of autophagy-associated molecular machineries, which may typify an alternate autophagic process or indicate that the bacterium actively exploits discrete functions of the canonical autophagy pathway to generate aBCVs.

Importantly, aBCV formation is tightly linked to bacterial egress, as new infection events of adjacent cells occur during the aBCV formation stage and are reduced upon inhibition of aBCV formation ([17](#)). Whether bacteria are released in a free state or contained within a membrane-bound vacuole remains to be established, but the maintenance of the originally infected cells argues that bacterial release is a nonlytic event, suggesting an exocytic

process (Fig. 1). Based on the autophagic nature of the aBCV, a tempting hypothesis to explain aBCV-dependent bacterial release is that the bacterium takes advantage of the secretory functions of the autophagy pathway (62) to deliver aBCV-enclosed bacteria to the extracellular milieu.

By mediating the final step of the *Brucella* intracellular cycle, aBCV formation is likely controlled by the bacterium, potentially via the VirB T4SS. Testing this hypothesis is challenging, as the preceding step of rBCV formation is T4SS dependent, precluding the classical use of reverse genetics and VirB T4SS-deficient mutants. Transient intracellular production of the VirB11 ATPase via a tightly controlled promoter, which energizes the T4SS, allows rBCV biogenesis and bacterial replication prior to T4SS intracellular inactivation and showed a requirement for a functional VirB apparatus in aBCV formation and bacterial egress, indicating that this final step of the *Brucella* intracellular cycle is likely controlled by T4SS effectors (63). Their future identification and characterization will likely shed light on the molecular mechanisms of aBCV formation and bacterial egress.

CONCLUDING REMARKS

Bacteria of the genus *Brucella* belong to a phylogenetic group closely associated with eukaryotic hosts. In this context, *Brucella*'s long-standing coevolution with mammalian hosts has shaped its complex intracellular cycle into the sequential exploitation of intracellular compartments of the endocytic, secretory, and autophagy pathways. *Brucella* uses an array of T4SS-delivered effectors and other virulence factors to modulate discrete functions of these compartments, modifying its original phagosome into an ER-associated replication-permissive organelle that subsequently coopts autophagic functions for completion of the bacterium's intracellular cycle, thus ensuring its survival, proliferation, and egress. While a few T4SS effectors have been identified, much remains to be understood about their modes of action and contributions to *Brucella*'s intracellular cycle. Future characterization of these proteins not only will reveal unsuspected aspects of *Brucella* intracellular strategies but also will teach us a great deal about host cell functions and their roles in many aspects of bacterial pathogenesis.

ACKNOWLEDGMENTS

We are most grateful to Andrew Rader for graphical assistance with figures. This work was supported by NIH grants AI129992 and AI127830.

REFERENCES

- Moreno E. 2014. Retrospective and prospective perspectives on zoonotic brucellosis. *Front Microbiol* 5:213 <http://dx.doi.org/10.3389/fmicb.2014.00213>.
- Pappas G, Akritidis N, Bosilkovski M, Tsianos E. 2005. Brucellosis. *N Engl J Med* 352:2325–2336 <http://dx.doi.org/10.1056/NEJMra050570>.
- Pappas G, Papadimitriou P, Akritidis N, Christou L, Tsianos EV. 2006. The new global map of human brucellosis. *Lancet Infect Dis* 6:91–99 [http://dx.doi.org/10.1016/S1473-3099\(06\)70382-6](http://dx.doi.org/10.1016/S1473-3099(06)70382-6).
- Atluri VL, Xavier MN, de Jong MF, den Hartigh AB, Tsois RM. 2011. Interactions of the human pathogenic *Brucella* species with their hosts. *Annu Rev Microbiol* 65:523–541 <http://dx.doi.org/10.1146/annurev-micro-090110-102905>.
- Celli J. 2015. The changing nature of the *Brucella*-containing vacuole. *Cell Microbiol* 17:951–958 <http://dx.doi.org/10.1111/cmi.12452>.
- Anderson TD, Cheville NF, Meador VP. 1986. Pathogenesis of placentitis in the goat inoculated with *Brucella abortus*. II. Ultrastructural studies. *Vet Pathol* 23:227–239 <http://dx.doi.org/10.1177/030098588602300302>.
- Pizarro-Cerdá J, Méresse S, Parton RG, van der Goot G, Sola-Landa A, Lopez-Goni I, Moreno E, Gorvel JP. 1998. *Brucella abortus* transits through the autophagic pathway and replicates in the endoplasmic reticulum of nonprofessional phagocytes. *Infect Immun* 66:5711–5724.
- Celli J, de Chastellier C, Franchini D-M, Pizarro-Cerdá J, Moreno E, Gorvel J-P. 2003. *Brucella* evades macrophage killing via VirB-dependent sustained interactions with the endoplasmic reticulum. *J Exp Med* 198:545–556 <http://dx.doi.org/10.1084/jem.20030088>.
- Salcedo SP, Chevrier N, Lacerda TLS, Ben Amara A, Gerart S, Gorvel VA, de Chastellier C, Blasco JM, Mege J-L, Gorvel J-P. 2013. Pathogenic brucellae replicate in human trophoblasts. *J Infect Dis* 207:1075–1083 <http://dx.doi.org/10.1093/infdis/jit007>.
- Detilleux PG, Deyoe BL, Cheville NF. 1990. Entry and intracellular localization of *Brucella* spp. in Vero cells: fluorescence and electron microscopy. *Vet Pathol* 27:317–328 <http://dx.doi.org/10.1177/030098589002700503>.
- Detilleux PG, Deyoe BL, Cheville NF. 1990. Penetration and intracellular growth of *Brucella abortus* in nonphagocytic cells in vitro. *Infect Immun* 58:2320–2328.
- Arenas GN, Staskevich AS, Aballay A, Mayorga LS. 2000. Intracellular trafficking of *Brucella abortus* in J774 macrophages. *Infect Immun* 68:4255–4263 <http://dx.doi.org/10.1128/IAI.68.7.4255-4263.2000>.
- Comerci DJ, Martínez-Lorenzo MJ, Sicra R, Gorvel J-P, Ugalde RA. 2001. Essential role of the VirB machinery in the maturation of the *Brucella abortus*-containing vacuole. *Cell Microbiol* 3:159–168 <http://dx.doi.org/10.1046/j.1462-5822.2001.00102.x>.
- Porte F, Liautard JP, Köhler S. 1999. Early acidification of phagosomes containing *Brucella suis* is essential for intracellular survival in murine macrophages. *Infect Immun* 67:4041–4047.
- Boschiroli ML, Ouahrani-Bettache S, Foulongne V, Michaux-Charachon S, Bourg G, Allardet-Servent A, Cazevielle C, Liautard JP, Ramuz M, O'Callaghan D. 2002. The *Brucella suis* virB operon is induced intracellularly in macrophages. *Proc Natl Acad Sci USA* 99:1544–1549 <http://dx.doi.org/10.1073/pnas.032514299>.
- Starr T, Ng TW, Wehrly TD, Knodler LA, Celli J. 2008. *Brucella* intracellular replication requires trafficking through the late endosomal/lysosomal compartment. *Traffic* 9:678–694 <http://dx.doi.org/10.1111/j.1600-0854.2008.00718.x>.
- Starr T, Child R, Wehrly TD, Hansen B, Hwang S, López-Otin C, Virgin HW, Celli J. 2012. Selective subversion of autophagy complexes facilitates completion of the *Brucella* intracellular cycle. *Cell Host Microbe* 11:33–45 <http://dx.doi.org/10.1016/j.chom.2011.12.002>.
- Sedzicki J, Tschon T, Low SH, Willemart K, Goldie KN, Letesson J-J, Stahlberg H, Dehio C. 2018. 3D correlative electron microscopy reveals

continuity of *Brucella*-containing vacuoles with the endoplasmic reticulum. *J Cell Sci* 131:jcs210799 <http://dx.doi.org/10.1242/jcs.210799>.

19. Sieira R, Comerci DJ, Sánchez DO, Ugalde RA. 2000. A homologue of an operon required for DNA transfer in *Agrobacterium* is required in *Brucella abortus* for virulence and intracellular multiplication. *J Bacteriol* 182:4849–4855 <http://dx.doi.org/10.1128/JB.182.17.4849-4855.2000>.

20. Delrue RM, Martínez-Lorenzo M, Lestrade P, Danese I, Bielarsz V, Mertens P, De Bolle X, Tibor A, Gorvel JP, Letesson JJ. 2001. Identification of *Brucella* spp. genes involved in intracellular trafficking. *Cell Microbiol* 3:487–497 <http://dx.doi.org/10.1046/j.1462-5822.2001.00131.x>.

21. O'Callaghan D, Cazevielle C, Allardet-Servent A, Boschiroli ML, Bourg G, Foulongne V, Frutos P, Kulakov Y, Ramuz M. 1999. A homologue of the *Agrobacterium tumefaciens* VirB and *Bordetella pertussis* Ptl type IV secretion systems is essential for intracellular survival of *Brucella suis*. *Mol Microbiol* 33:1210–1220 <http://dx.doi.org/10.1046/j.1365-2958.1999.01569.x>.

22. Celli J, Salcedo SP, Gorvel J-P. 2005. *Brucella* coopts the small GTPase Sar1 for intracellular replication. *Proc Natl Acad Sci USA* 102:1673–1678 <http://dx.doi.org/10.1073/pnas.0406873102>.

23. Hartigh den AB, Rolán HG, de Jong MF, Tsois RM. 2008. VirB3 to VirB6 and VirB8 to VirB11, but not VirB7, are essential for mediating persistence of *Brucella* in the reticuloendothelial system. *J Bacteriol* 190:4427–4436.

24. Lestrade P, Delrue RM, Danese I, Didembourg C, Taminiau B, Mertens P, De Bolle X, Tibor A, Tang CM, Letesson JJ. 2000. Identification and characterization of in vivo attenuated mutants of *Brucella melitensis*. *Mol Microbiol* 38:543–551 <http://dx.doi.org/10.1046/j.1365-2958.2000.02150.x>.

25. Green ER, Mecsas J. 2016. Bacterial secretion systems: an overview. *Microbiol Spectr* 4(1):VMBF-0012-2015.

26. Juhas M, Crook DW, Hood DW. 2008. Type IV secretion systems: tools of bacterial horizontal gene transfer and virulence. *Cell Microbiol* 10:2377–2386 <http://dx.doi.org/10.1111/j.1462-5822.2008.01187.x>.

27. de Jong MF, Sun Y-H, den Hartigh AB, van Dijk JM, Tsois RM. 2008. Identification of VceA and VceC, two members of the VjbR regulon that are translocated into macrophages by the *Brucella* type IV secretion system. *Mol Microbiol* 70:1378–1396 <http://dx.doi.org/10.1111/j.1365-2958.2008.06487.x>.

28. de Barsy M, Jamet A, Filopon D, Nicolas C, Laloux G, Rual J-F, Muller A, Twizere J-C, Nkengfac B, Vandenhaute J, Hill DE, Salcedo SP, Gorvel J-P, Letesson J-J, De Bolle X. 2011. Identification of a *Brucella* spp. secreted effector specifically interacting with human small GTPase Rab2. *Cell Microbiol* 13:1044–1058 <http://dx.doi.org/10.1111/j.1462-5822.2011.01601.x>.

29. Marchesini MI, Herrmann CK, Salcedo SP, Gorvel J-P, Comerci DJ. 2011. In search of *Brucella abortus* type IV secretion substrates: screening and identification of four proteins translocated into host cells through VirB system. *Cell Microbiol* 13:1261–1274 <http://dx.doi.org/10.1111/j.1462-5822.2011.01618.x>.

30. Myeni S, Child R, Ng TW, Kupko JJ III, Wehrly TD, Porcella SF, Knodler LA, Celli J. 2013. *Brucella* modulates secretory trafficking via multiple type IV secretion effector proteins. *PLoS Pathog* 9:e1003556 <http://dx.doi.org/10.1371/journal.ppat.1003556>.

31. Salcedo SP, Marchesini MI, Degos C, Terwagne M, Von Bargen K, Lepidi H, Herrmann CK, Santos Lacerda TL, Imbert PR, Pierre P, Alexopoulou L, Letesson JJ, Comerci DJ, Gorvel JP. 2013. BtpB, a novel *Brucella* TIR-containing effector protein with immune modulatory functions. *Front Cell Infect Microbiol* 3:28 <http://dx.doi.org/10.3389/fcimb.2013.00028>.

32. Döhmer PH, Valguarnera E, Czibener C, Ugalde JE. 2014. Identification of a type IV secretion substrate of *Brucella abortus* that participates in the early stages of intracellular survival. *Cell Microbiol* 16:396–410 <http://dx.doi.org/10.1111/cmi.12224>.

33. Chaves-Olarte E, Guzmán-Verri C, Méresse S, Desjardins M, Pizarro-Cerdá J, Badilla J, Gorvel J-P, Moreno E. 2002. Activation of Rho and

Rab GTPases dissociates *Brucella abortus* internalization from intracellular trafficking. *Cell Microbiol* 4:663–676 <http://dx.doi.org/10.1046/j.1462-5822.2002.00221.x>.

34. Pizarro-Cerdá J, Moreno E, Sanguedolce V, Mege JL, Gorvel JP. 1998. Virulent *Brucella abortus* prevents lysosome fusion and is distributed within autophagosome-like compartments. *Infect Immun* 66:2387–2392.

35. Sieira R, Comerci DJ, Pietrasanta LI, Ugalde RA. 2004. Integration host factor is involved in transcriptional regulation of the *Brucella abortus* virB operon. *Mol Microbiol* 54:808–822 <http://dx.doi.org/10.1111/j.1365-2958.2004.04316.x>.

36. Deghelt M, Mullier C, Sternon J-F, Francis N, Laloux G, Dotreppe D, Van der Henst C, Jacobs-Wagner C, Letesson J-J, De Bolle X. 2014. G1-arrested newborn cells are the predominant infectious form of the pathogen *Brucella abortus*. *Nat Commun* 5:4366 <http://dx.doi.org/10.1038/ncomms5366>.

37. Budnik A, Stephens DJ. 2009. ER exit sites—localization and control of COPII vesicle formation. *FEBS Lett* 583:3796–3803 <http://dx.doi.org/10.1016/j.febslet.2009.10.038>.

38. Miller CN, Smith EP, Cundiff JA, Knodler LA, Bailey Blackburn J, Lupashin V, Celli J. 2017. A *Brucella* type IV effector targets the COG tethering complex to remodel host secretory traffic and promote intracellular replication. *Cell Host Microbe* 22:317–329.e7 <http://dx.doi.org/10.1016/j.chom.2017.07.017>.

39. Taguchi Y, Imaoka K, Kataoka M, Uda A, Nakatsu D, Horii-Okazaki S, Kunishige R, Kano F, Murata M. 2015. Yip1A, a novel host factor for the activation of the IRE1 pathway of the unfolded protein response during *Brucella* infection. *PLoS Pathog* 11:e1004747 <http://dx.doi.org/10.1371/journal.ppat.1004747>.

40. Walter P, Ron D. 2011. The unfolded protein response: from stress pathway to homeostatic regulation. *Science* 334:1081–1086 <http://dx.doi.org/10.1126/science.1209038>.

41. Smith JA, Khan M, Magnani DD, Harms JS, Durward M, Radhakrishnan GK, Liu Y-P, Splitter GA. 2013. *Brucella* induces an unfolded protein response via TcpB that supports intracellular replication in macrophages. *PLoS Pathog* 9:e1003785 <http://dx.doi.org/10.1371/journal.ppat.1003785>.

42. de Jong MF, Starr T, Winter MG, den Hartigh AB, Child R, Knodler LA, van Dijk JM, Celli J, Tsois RM. 2013. Sensing of bacterial type IV secretion via the unfolded protein response. *mBio* 4:e00418-12 <http://dx.doi.org/10.1128/mBio.00418-12>.

43. Qin Q-M, Pei J, Ancona V, Shaw BD, Ficht TA, de Figueiredo P. 2008. RNAi screen of endoplasmic reticulum-associated host factors reveals a role for IRE1α in supporting *Brucella* replication. *PLoS Pathog* 4:e1000110–e1000116 <http://dx.doi.org/10.1371/journal.ppat.1000110>.

44. Levine B, Mizushima N, Virgin HW. 2011. Autophagy in immunity and inflammation. *Nature* 469:323–335 <http://dx.doi.org/10.1038/nature09782>.

45. Huang J, Brumell JH. 2014. Bacteria-autophagy interplay: a battle for survival. *Nat Rev Microbiol* 12:101–114 <http://dx.doi.org/10.1038/nrmicro3160>.

46. Asrat S, de Jesús DA, Hempstead AD, Ramabhadran V, Isberg RR. 2014. Bacterial pathogen manipulation of host membrane trafficking. *Annu Rev Cell Dev Biol* 30:79–109 <http://dx.doi.org/10.1146/annurev-cellbio-100913-013439>.

47. Kagan JC, Roy CR. 2002. *Legionella* phagosomes intercept vesicular traffic from endoplasmic reticulum exit sites. *Nat Cell Biol* 4:945–954 <http://dx.doi.org/10.1038/ncb883>.

48. Kagan JC, Stein M-P, Pypaert M, Roy CR. 2004. *Legionella* subvert the functions of Rab1 and Sec22b to create a replicative organelle. *J Exp Med* 199:1201–1211 <http://dx.doi.org/10.1084/jem.20031706>.

49. Hackstadt T, Rockey DD, Heinzen RA, Scidmore MA. 1996. *Chlamydia trachomatis* interrupts an exocytic pathway to acquire endogenously synthesized sphingomyelin in transit from the Golgi apparatus to

- the plasma membrane. *EMBO J* 15:964–977 <http://dx.doi.org/10.1002/j.1460-2075.1996.tb00433.x>.
50. Scidmore MA, Fischer ER, Hackstadt T. 1996. Sphingolipids and glycoproteins are differentially trafficked to the *Chlamydia trachomatis* inclusion. *J Cell Biol* 134:363–374 <http://dx.doi.org/10.1083/jcb.134.2.363>.
51. Bhui T, Roy JK. 2014. Rab proteins: the key regulators of intracellular vesicle transport. *Exp Cell Res* 328:1–19 <http://dx.doi.org/10.1016/j.yexcr.2014.07.027>.
52. Salcedo SP, Marchesini MI, Lelouard H, Fugier E, Jolly G, Balor S, Muller A, Lapaque N, Demaria O, Alexopoulou L, Comerci DJ, Ugalde RA, Pierre P, Gorvel J-P. 2008. Brucella control of dendritic cell maturation is dependent on the TIR-containing protein Btp1. *PLoS Pathog* 4:e21 <http://dx.doi.org/10.1371/journal.ppat.0040021>.
53. Jakka P, Namani S, Murugan S, Rai N, Radhakrishnan G. 2017. The *Brucella* effector protein TcpB induces degradation of inflammatory caspases and thereby subverts non-canonical inflammasome activation in macrophages. *J Biol Chem* 292:20613–20627 <http://dx.doi.org/10.1074/jbc.M117.815878>.
54. Alaidarous M, Ve T, Casey LW, Valkov E, Ericsson DJ, Ullah MO, Schembri MA, Mansell A, Sweet MJ, Kobe B. 2014. Mechanism of bacterial interference with TLR4 signaling by *Brucella* Toll/interleukin-1 receptor domain-containing protein TcpB. *J Biol Chem* 289:654–668 <http://dx.doi.org/10.1074/jbc.M113.523274>.
55. Chaudhary A, Ganguly K, Cabantous S, Waldo GS, Micheva-Viteva SN, Nag K, Hlavacek WS, Tung C-S. 2012. The *Brucella* TIR-like protein TcpB interacts with the death domain of MyD88. *Biochem Biophys Res Commun* 417:299–304 <http://dx.doi.org/10.1016/j.bbrc.2011.11.104>.
56. Sengupta D, Koblansky A, Gaines J, Brown T, West AP, Zhang D, Nishikawa T, Park SG, Roop RM II, Ghosh S. 2010. Subversion of innate immune responses by *Brucella* through the targeted degradation of the TLR signaling adapter, MAL. *J Immunol* 184:956–964 <http://dx.doi.org/10.4049/jimmunol.0902008>.
57. Radhakrishnan GK, Yu Q, Harms JS, Splitter GA. 2009. *Brucella* TIR domain-containing protein mimics properties of the Toll-like receptor adaptor protein TIRAP. *J Biol Chem* 284:9892–9898 <http://dx.doi.org/10.1074/jbc.M805458200>.
58. Ciril C, Wieser A, Yadav M, Duerr S, Schubert S, Fischer H, Stappert D, Wantia N, Rodriguez N, Wagner H, Svanborg C, Miethke T. 2008. Subversion of Toll-like receptor signaling by a unique family of bacterial Toll/interleukin-1 receptor domain-containing proteins. *Nat Med* 14:399–406 <http://dx.doi.org/10.1038/nm1734>.
59. Willett R, Ungar D, Lupashin V. 2013. The Golgi puppet master: COG complex at center stage of membrane trafficking interactions. *Histochem Cell Biol* 140:271–283 <http://dx.doi.org/10.1007/s00418-013-1117-6>.
60. Cui G, Wei P, Zhao Y, Guan Z, Yang L, Sun W, Wang S, Peng Q. 2014. *Brucella* infection inhibits macrophages apoptosis via Nedd4-dependent degradation of calpain2. *Vet Microbiol* 174:195–205 <http://dx.doi.org/10.1016/j.vetmic.2014.08.033>.
61. Gross A, Terraza A, Ouahrani-Bettache S, Liautard JP, Dornand J. 2000. In vitro *Brucella suis* infection prevents the programmed cell death of human monocytic cells. *Infect Immun* 68:342–351 <http://dx.doi.org/10.1128/IAI.68.1.342-351.2000>.
62. Kimura T, Jia J, Claude-Taupin A, Kumar S, Choi SW, Gu Y, Mudd M, Dupont N, Jiang S, Peters R, Farzam F, Jain A, Lidke KA, Adams CM, Johansen T, Deretic V. 2017. Cellular and molecular mechanism for secretory autophagy. *Autophagy* 13:1084–1085 <http://dx.doi.org/10.1080/15548627.2017.1307486>.
63. Smith EP, Miller CN, Child R, Cundiff JA, Celli J. 2016. Postreplication roles of the *Brucella* VirB type IV secretion system uncovered via conditional expression of the VirB11 ATPase. *mBio* 7:e01730-16.