


## 6.1



## 6.2/6.3

$$H_0 t = \int_0^a \frac{da}{[\Omega_0 / a + (1 - \Omega_0)]^{1/2}} = \int_0^a \frac{ada}{[\Omega_0 a + (1 - \Omega_0) a^2]^{1/2}}$$

$$= \int_0^a \frac{ada}{(1 - \Omega_0)^{1/2} [(a + \frac{\Omega_0}{2(1 - \Omega_0)})^2 - (\frac{\Omega_0}{2(1 - \Omega_0)})^2]^{1/2}}$$

$$\Omega_0 > 1$$

$$H_0 t = \int_0^a \frac{ada}{(\Omega_0 - 1)^{1/2} [-(a - \frac{\Omega_0}{2(\Omega_0 - 1)})^2 + (\frac{\Omega_0}{2(1 - \Omega_0)})^2]^{1/2}}$$

$$a = \frac{\Omega_0}{2(\Omega_0 - 1)}(1 - \cos \theta)$$

$$\Rightarrow H_0 t = \int_0^\theta \frac{\frac{\Omega_0}{2(\Omega_0 - 1)}(1 - \cos \theta) \cdot \frac{\Omega_0}{2(\Omega_0 - 1)} \sin \theta d\theta}{(\Omega_0 - 1)^{1/2} [-(\frac{\Omega_0}{2(\Omega_0 - 1)})^2 \cos^2 \theta + (\frac{\Omega_0}{2(1 - \Omega_0)})^2]^{1/2}}$$

$$= \frac{\Omega_0}{2(\Omega_0 - 1)^{3/2}} \int_0^\theta (1 - \cos \theta) d\theta = \frac{\Omega_0}{2(\Omega_0 - 1)^{3/2}} (\theta - \sin \theta)$$

$$\Rightarrow t = \frac{\Omega_0}{2H_0(\Omega_0 - 1)^{3/2}} (\theta - \sin \theta)$$

$$\Omega_0 < 1$$

$$H_0 t = \int_0^a \frac{ada}{(1 - \Omega_0)^{1/2} [(a + \frac{\Omega_0}{2(1 - \Omega_0)})^2 - (\frac{\Omega_0}{2(1 - \Omega_0)})^2]^{1/2}}$$

$$a = \frac{\Omega_0}{2(\Omega_0 - 1)}(\cosh \theta - 1)$$

$$\Rightarrow H_0 t = \int_0^\theta \frac{\frac{\Omega_0}{2(1 - \Omega_0)}(\cosh \theta - 1) \cdot \frac{\Omega_0}{2(1 - \Omega_0)} \sinh \theta d\theta}{(1 - \Omega_0)^{1/2} [(\frac{\Omega_0}{2(\Omega_0 - 1)})^2 \cosh^2 \theta - (\frac{\Omega_0}{2(1 - \Omega_0)})^2]^{1/2}}$$

$$= \frac{\Omega_0}{2(1 - \Omega_0)^{3/2}} \int_0^\theta (\cosh \theta - 1) d\theta = \frac{\Omega_0}{2(1 - \Omega_0)^{3/2}} (\sinh \theta - \theta)$$

$$\Rightarrow t = \frac{\Omega_0}{2H_0(\Omega_0 - 1)^{3/2}} (\theta - \sinh \theta)$$

## 6.4a

$$\frac{H^2}{H_0^2} = \frac{\Omega_{m,0}}{a^3} + (1 - \Omega_{m,0}) = \frac{\dot{a}^2}{H_0^2 a^2}$$

$$\Rightarrow \frac{da}{H_0 a dt} = \left[ \frac{\Omega_{m,0}}{a^3} + (1 - \Omega_{m,0}) \right]^{1/2}$$

$$\Rightarrow H_0 dt = \frac{da}{a \left[ \frac{\Omega_{m,0}}{a^3} - (\Omega_{m,0} - 1) \right]^{1/2}} = \frac{da}{a \left( \frac{\Omega_{m,0}}{(\Omega_{m,0} - 1) a^3} - 1 \right)^{1/2} (\Omega_{m,0} - 1)^{1/2}}$$

$$a^3 = \frac{\Omega_{m,0}}{(\Omega_{m,0} - 1) \csc^2 \theta} = \frac{a_{\max}^3}{\csc^2 \theta} \Rightarrow a = \frac{a_{\max}}{\csc^{2/3} \theta} \Rightarrow \theta = \sin^{-1} \left( \frac{a}{a_{\max}} \right)^{3/2}$$

$$H_0 dt = \frac{a_{\max} \left( -\frac{2}{3} \csc^{-5/3} \theta \right) (-\csc \theta \cot \theta) d\theta}{\frac{a_{\max}}{\csc^{2/3} \theta} \cot \theta (\Omega_{m,0} - 1)^{1/2}} = \frac{2 d\theta}{3 (\Omega_{m,0} - 1)^{1/2}}$$

$$\Rightarrow H_0 t = \frac{2\theta}{3 (\Omega_{m,0} - 1)^{1/2}} = \frac{2}{3 (\Omega_{m,0} - 1)^{1/2}} \sin^{-1} \left( \frac{a}{a_{\max}} \right)^{3/2}$$

## 6.4b

$$\frac{H^2}{H_0^2} = \frac{\Omega_{m,0}}{a^3} + (1 - \Omega_{m,0}) = \frac{\dot{a}^2}{H_0^2 a^2}$$

$$\Rightarrow \frac{da}{H_0 a dt} = \left[ \frac{\Omega_{m,0}}{a^3} + (1 - \Omega_{m,0}) \right]^{1/2}$$

$$\Rightarrow H_0 dt = \frac{da}{a \left[ \frac{\Omega_{m,0}}{a^3} + (1 - \Omega_{m,0}) \right]^{1/2}} = \frac{da}{a \left( \frac{\Omega_{m,0}}{(1 - \Omega_{m,0}) a^3} + 1 \right)^{1/2} (1 - \Omega_{m,0})^{1/2}}$$

$$a^3 = \frac{\Omega_{m,0}}{(1 - \Omega_{m,0}) \cot^2 \theta} = \frac{a_{m\Lambda}^3}{\cot^2 \theta} \Rightarrow a = \frac{a_{m\Lambda}}{\cot^{2/3} \theta} \Rightarrow \tan \theta = \left( \frac{a}{a_{m\Lambda}} \right)^{3/2}$$

$$H_0 dt = \frac{a_{m\Lambda} \left( -\frac{2}{3} \cot^{-5/3} \theta \right) (-\csc^2 \theta) d\theta}{\frac{a_{m\Lambda}}{\cot^{2/3} \theta} \csc \theta (1 - \Omega_{m,0})^{1/2}} = \frac{2 \sec \theta d\theta}{3(1 - \Omega_{m,0})^{1/2}}$$

$$\Rightarrow H_0 t = \frac{2}{3(1 - \Omega_{m,0})^{1/2}} \int_0^\theta \sec \theta d\theta = \frac{2}{3(1 - \Omega_{m,0})^{1/2}} \int_0^\theta \sec \theta \frac{\sec \theta + \tan \theta}{\sec \theta + \tan \theta} d\theta$$

$$H_0 t = \frac{2}{3(1 - \Omega_{m,0})^{1/2}} \int_0^\theta \frac{d(\sec \theta + \tan \theta)}{\sec \theta + \tan \theta} = \frac{2}{3(1 - \Omega_{m,0})^{1/2}} \ln(\sec \theta + \tan \theta)$$

$$H_0 t = \frac{2}{3(1 - \Omega_{m,0})^{1/2}} \ln \left[ \left( \frac{a}{a_{m\Lambda}} \right)^{3/2} + \sqrt{1 + \left( \frac{a}{a_{m\Lambda}} \right)^3} \right]$$

$$\Rightarrow t_0 = \frac{2H_0^{-1}}{3(1 - \Omega_{m,0})^{1/2}} \ln \left[ \left( \frac{1 - \Omega_{m,0}}{\Omega_{m,0}} \right)^{1/2} + \frac{1}{\Omega_{m,0}^{1/2}} \right] = \frac{2H_0^{-1}}{3(1 - \Omega_{m,0})^{1/2}} \ln \left[ \frac{\sqrt{1 - \Omega_{m,0}} + 1}{\Omega_{m,0}^{1/2}} \right]$$

$$\Omega_{m,0} = 0.27, H_0 = 71 \text{ km/s/Mpc}$$

$$\Rightarrow t_0 = \frac{2}{71 \text{ km/s/Mpc} \times 3(1 - 0.27)^{1/2}} \ln \left[ \frac{\sqrt{0.73} + 1}{0.27^{1/2}} \right] = 13.6 \text{ Gyr}$$

## 6.5

$$H_0 t = \frac{ada}{\sqrt{\Omega_{r,0}}} \frac{1}{\sqrt{1 + \frac{a}{a_{rm}}}}$$

$$x = \frac{a}{a_{rm}}$$

$$\Rightarrow H_0 t = \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} \frac{xdx}{\sqrt{1+x}}$$

$$x = \tan^2 \theta$$

$$\Rightarrow H_0 t = \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} \frac{\tan^2 \theta \cdot 2 \tan \theta \sec^2 \theta d\theta}{\sec \theta} = \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} \tan^2 \theta \cdot 2 \tan \theta \sec \theta d\theta$$

$$= \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} 2 \tan^2 \theta d(\sec \theta) = \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} 2(\sec^2 \theta - 1) d(\sec \theta)$$

$$\Rightarrow H_0 t = \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} 2 \left( \frac{1}{3} \sec^3 \theta - \sec \theta \right) \Big|_0^\theta = \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} \left[ \frac{4}{3} + 2 \sec \theta \left( \frac{1}{3} \sec^2 \theta - 1 \right) \right]$$

$$= \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} \left[ \frac{4}{3} + 2\sqrt{1+x} \left( \frac{1}{3} (1+x) - 1 \right) \right] = \frac{a_{rm}^2}{\sqrt{\Omega_{r,0}}} \frac{4}{3} \left[ 1 - \sqrt{1+x} \left( 1 - \frac{x}{2} \right) \right]$$

$$\Rightarrow H_0 t = \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left[ 1 - \left( 1 - \frac{a}{2a_{rm}} \right) \left( 1 + \frac{a}{a_{rm}} \right)^{1/2} \right]$$

$$a \ll a_{rm}$$

$$H_0 t = \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left[ 1 - \left( 1 - \frac{a}{2a_{rm}} \right) \left( 1 + \frac{a}{a_{rm}} \right)^{1/2} \right] \approx \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left[ 1 - \left( 1 - \frac{a}{2a_{rm}} \right) \left( 1 + \frac{a}{2a_{rm}} - \frac{a^2}{8a_{rm}^2} \right) \right]$$

$$= \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left[ 1 - \left( 1 - \frac{3}{8} \left( \frac{a}{a_{rm}} \right)^2 \right) \right] = \frac{a^2}{2\sqrt{\Omega_{r,0}}}$$

$$\Rightarrow a \approx (2\sqrt{\Omega_{r,0}} H_0 t)^{1/2}$$

$$a \gg a_{rm}$$

$$\begin{aligned} H_0 t &= \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left[ 1 - \left( 1 - \frac{a}{2a_{rm}} \right) \left( 1 + \frac{a}{a_{rm}} \right)^{1/2} \right] \approx \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left( \frac{a}{2a_{rm}} \right) \left( \frac{a}{a_{rm}} \right)^{1/2} \\ &= \frac{2a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left( \frac{a}{a_{rm}} \right)^{3/2} = \frac{2}{3\sqrt{\Omega_{r,0}/a_{rm}}} a^{3/2} = \frac{2}{3\sqrt{\Omega_{m,0}}} a^{3/2} \\ \Rightarrow a &\approx \left( \frac{3}{2} \sqrt{\Omega_{m,0}} H_0 t \right)^{2/3} \end{aligned}$$

$$a = a_{rm}$$

$$\begin{aligned} \Rightarrow H_0 t_{rm} &= \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left[ 1 - \left( 1 - \frac{1}{2} \right) (1+1)^{1/2} \right] = \frac{4a_{rm}^2}{3\sqrt{\Omega_{r,0}}} \left[ 1 - \frac{1}{\sqrt{2}} \right] \\ \Rightarrow t_{rm} &\approx 0.39 \frac{\Omega_{r,0}^{3/2}}{\Omega_{m,0}^2} H_0^{-1} \end{aligned}$$

$$\Omega_{r,0} = 8.4 \times 10^{-5}, \Omega_{m,0} = 0.3, H_0^{-1} = 14 \text{ Gyr}$$

$$\Rightarrow t_{rm} \approx 0.39 \frac{\Omega_{r,0}^{3/2}}{\Omega_{m,0}^2} H_0^{-1} = 47,000 \text{ yr}$$