

Rick Sarmento

Look at parametric substitutions for Matter - Curvature only universe

Original diff eq for Matter - Curvature only universe

$$\frac{da}{dt} = H_0 \left(\frac{\Omega_m}{a} + (1 - \Omega_m) \right)^{\frac{1}{2}}$$

Look at parametric solution for $\Omega_m > 1.0$

The following are ' given ' :

$$\text{In[48]:= } a_\theta = \frac{1}{2} \frac{\Omega_m}{(\Omega_m - 1)} (1 - \cos[\theta]);$$

$$\text{In[51]:= } da_\theta = \partial_\theta (a_\theta)$$

$$\text{Out[51]= } \frac{\sin[\theta] \Omega_m}{2 (-1 + \Omega_m)}$$

$$\text{In[54]:= } t_\theta = \frac{1}{2 H_0} \frac{\Omega_m}{(\Omega_m - 1)^{\frac{3}{2}}} (\theta - \sin[\theta]);$$

$$\text{In[55]:= } dt_\theta = \partial_\theta (t_\theta)$$

$$\text{Out[55]= } \frac{(1 - \cos[\theta]) \Omega_m}{2 H_0 (-1 + \Omega_m)^{3/2}}$$

So

$$\text{In[56]:= } da_t = \frac{da_\theta}{dt_\theta}$$

$$\text{Out[56]= } \frac{\sin[\theta] H_0 \sqrt{-1 + \Omega_m}}{1 - \cos[\theta]}$$

Substitute into the Original equation (the “.” operator means substitute)

$$\text{In[59]:= } \frac{da}{dt} == H_0 \left(\frac{\Omega_m}{a} + (1 - \Omega_m) \right)^{\frac{1}{2}} / . a \rightarrow a_\theta / . da \rightarrow da_\theta / . dt \rightarrow dt_\theta$$

$$\text{Out[59]= } \frac{\sin[\theta] H_0 \sqrt{-1 + \Omega_m}}{1 - \cos[\theta]} == H_0 \sqrt{1 + \frac{2 (-1 + \Omega_m)}{1 - \cos[\theta]} - \Omega_m}$$

$$\text{In[61]:= Assuming } [2 \pi \geq \theta \geq 0 \& \Omega_m > 1,$$

$$\text{FullSimplify} \left[\frac{\sin[\theta] H_0 \sqrt{-1 + \Omega_m}}{1 - \cos[\theta]} == H_0 \sqrt{1 + \frac{2 (-1 + \Omega_m)}{1 - \cos[\theta]} - \Omega_m}, \text{Reals} \right]$$

$$\text{Out[61]= } \cot\left[\frac{\theta}{2}\right] H_0 \sqrt{-1 + \Omega_m} == H_0 \sqrt{\cot^2\left[\frac{\theta}{2}\right] (-1 + \Omega_m)}$$

Assuming

$$\cot\left[\frac{\theta}{2}\right] = \sqrt{\cot\left[\frac{\theta}{2}\right]^2}$$

we get

$$\cot\left[\frac{\theta}{2}\right] H_0 \sqrt{-1 + \Omega_m} == \cot\left[\frac{\theta}{2}\right] H_0 \sqrt{-1 + \Omega_m}$$

which is always TRUE.

```
In[63]:= ParametricPlot[
  {H0 ((θ - Sin[θ]) Ωm, 1/2 Ωm (1 - Cos[θ]))},
  {θ, 0, 2 π}, AspectRatio -> .4]
```

