

Hubble's Survey of the Ultraviolet universe: Panchromatic Extragalactic Research (“SUPER”)

Rogier Windhorst (Arizona State University) & the HST SUPER team:

H. Teplitz (co-PI), T. Ashcraft, N. Bond, G. Bruzual, D. Calzetti, S. Casertano, S. Cohen, C. Conselice, D. de Mello, M. Dickinson, S. Driver, H. Ferguson, K. Finkelstein, S. Finkelstein, J. Frogel, J. Gallego, J. Gardner, E. Gawiser, M. Giavalisco, A. Grazian, N. Grogin, N. Hathi, T. Heckman, R. Jansen, C. Kaleida, S. Kassin, S. Kaviraj, H. Kim, A. Koekemoer, D. Koo, J. Lotz, R. Lucas, J. MacKenty, S. Malhotra, P. McCarthy, M. Mechtley, G. Meurer, M. Mutchler, K. Noeske, R. O’Connell, R. Overzier, N. Pirzkal, M. Rafelski, S. Ravindranath, J. Rhoads, M. Rieke, B. Robertson, M. Rutkowski, R. Ryan, C. Scarlata, M. Seibert, S. Salim, B. Siana, J. Silk, A. Straughn, K. Tamura, S. Wilkins, H. Yan, & S. Yi


221st AAS meeting, Session 228 “High Resolution Ultraviolet Imaging with the HST — II [high redshift]”.

Tuesday January 8, 2013, 2:00 pm, Room 202A, Long Beach Convention Center, Long Beach, CA.


Outline and 4 Science Goals

- Deep wide-field HST WFC3 UV imaging is critical NOW to prepare for JWST ($\lambda \gtrsim 0.7 \mu\text{m}$), and to define a $\gtrsim 8$ -meter UV-optical sequel to HST:
- (1) The physics and evolution of SF in low-mass galaxies over the LAST 9 Gyrs: Critical benchmark to understand cosmic reionization at $z \gtrsim 6$;
- (2) Evolution of the star/dust/gas mixture in SF regions, and the influence of supernovae and AGN feedback;
- (3) Evolution of young, star-forming sub-galactic clumps induced by mergers or gas accretion, and the growth of stable galaxy disks;
- (4) Late-epoch SF & structural evolution in massive early-type galaxies.
- (5) Summary and Conclusions.

Role of ACS in HST Post-SM4 Imaging Capability



ACS/WFC superior to WFC3 survey efficiency at visible-red wavelengths



WFC3/UVIS channel unprecedented UV–blue throughput & areal coverage:

$QE \gtrsim 70\%$, $4k \times 4k$ array of $0''.04$ pixel, $FOV \simeq 2'.67 \times 2'.67$.

WFC3/IR channel unprecedented near–IR throughput & areal coverage:


$QE \gtrsim 70\%$, $1k \times 1k$ array of $0''.13$ pixel, $FOV \simeq 2'.25 \times 2'.25$.

- WFC3 filters designed for star-formation and galaxy assembly at $z \simeq 0-8$.
- HST WFC3 and its UVIS channel a critical pathfinder for JWST science.




10 filters with HST/WFC3 & ACS reaching $AB=26.5-27.0$ mag ($10-\sigma$) over 40 arcmin^2 at $0.07-0.15''$ FWHM from $0.2-1.7 \mu\text{m}$ (UVUBVizYJH).

- ERS in GOODS-S v2: using WFC3 for what it was designed to do.
- JWST adds $0.05-0.2''$ FWHM imaging to $AB \simeq 31.5$ mag at $0.7-5 \mu\text{m}$, and $0.2-1.2''$ FWHM at $5-29 \mu\text{m}$, tracing young+old SEDs & dust.




- *Rest-frame* SEDs of objects with **young**, **mixed**, and **old** stellar populations with WFC3 UV, ACS B & CANDELS VIJH filters.
- **Diamonds**: Rest-frame 1300Å and 2600Å to measure UV β -slope.
- For $0.2 \lesssim z \lesssim 2.0$ WFC3 UV, ACS F435W & CANDELS (F606W, F814W) filters sample the β -slope for SF objects in $\gtrsim 2$ filters at $\lesssim 0''.1$ FWHM.

SUPER COSMOS Field





SUPER EGS Field



- Viable APT solutions to cover CANDELS fields (split over 2 ORIENTs ~ 180 days apart to aid scheduling).
- WFC3/UVIS tiles in red, ACS/F435W tiles in green (parallels), or blue (primary); CANDELS WFC3/IR F125W, F160W tiles in grey.

(1) Physics and evolution of SF in low-mass galaxies over the LAST 9 Gyrs:





[LEFT] Evolution of the UV spectral slope β (Finkelstein et al. 2011). Dark pink/dark blue region is WFC3 ERS β -range (Hathi et al. 2012).

- Deep wide-field WFC3 UV traces β -evolution for 16,000 SF clumps at $0.2 \lesssim z \lesssim 2$ (grey strip) \Rightarrow dust A_V in SF knots of sub- L^* galaxies.

[RIGHT] Evolution of the cosmic SF-rate density (“SFRD”; Bouwens et al. 2011). Blue dots are before and red dots after dust-correction.

- Deep wide-field WFC3 UV will yield SFRD in low-mass galaxies at $z \lesssim 2$.
- Essential synergy with Herschel FIR \Rightarrow relation between β and dust attenuation, providing the most robust estimate of the SFRD at $z \lesssim 2$.




[LEFT panels] Rest-frame UV Luminosity Functions (LFs) based on UV-dropouts (Hathi et al. 2010, 2012).

[RIGHT] ● (top) Evolution of the faint-end Schechter slope α and M^* (e.g., Hathi et al. 2010, Oesch et al. 2010b).

● (Bottom) M^* vs. z behavior resembles the cosmic SF history (Madau et al. 1996), and reflects the process of galaxy assembly and downsizing.

● Dark pink indicates current WFC ERS + CANDELS uncertainties.

● Grey wedge shows the significant improvement from deep wide-field WFC3 UV imaging for $0.2 \lesssim z \lesssim 2.5$.





[LEFT panels] WFC3 ERS color images of galaxies at $z \simeq 0.75$ shown in the 7 WFC3 UV+B & CANDELS filters. All show measurable UV flux.

[RIGHT] $z \simeq 1$ galaxy in same filters (upper left). WFC3 UV priors can dissect deep ground-based U_{360} -images (0.9 FWHM; Grazian et al. 2006), recovering fluxes for $\gtrsim 65\%$ of HST's SF-clumps (lower left).

- Right panels show pixel-to-pixel dust (A_V) & SF-age ($\log t/\text{yr}$) maps.
- Deep wide-field WFC3 UV will yield pixel-to-pixel mass, age, A_V and dust-maps for ~ 2000 galaxies at $0.2 \lesssim z \lesssim 2$.

(2) Evolution of star/dust/gas mix in SF regions, and SNe/AGN feedback:




[LEFT] Average IGM transmission vs. wavelength. WFC3 F275W samples Lyman-continuum escape fraction f_{esc} at $z \simeq 2.45$, where average transmission is $e^{-\tau} \simeq 0.250$ (Siana et al. 2012).

- Deep wide-field WFC3 UV will yield f_{esc} for ~ 800 galaxies at $z \simeq 2.45$, and relate it to physical galaxy properties (mass, type, A_V , V_{rot}/σ_g).

[RIGHT] Bolometric luminosity vs. dust attenuation (L_{IR}/L_{UV}) for $z \simeq 2$ compared to local galaxies (Reddy et al. 2010), suggesting evolution of the net extinction in SF galaxies with time.

- Deep wide-field WFC3 UV: +800 dusty LBGs in redshift gap $0.2 \lesssim z \lesssim 2$.

(3) Evolution of SF clumps and the growth of stable galaxy disks:





[LEFT 2 panels] Radial variation of age and mass surface-density of subgalactic clumps at $z \simeq 2$ vs. galacto-centric distance (Guo et al. 2011, 2012).

[MIDDLE 2] Ordered rotation (V_{rot}) & disturbed motions (σ_g) shows strong correlation with rest-frame blue morphology (Kassin⁺ 2007).

[RIGHT] Ratio of ordered/disordered motions (V_{rot}/σ_g) correlated with HST rest-frame B -band morphology: the most disturbed galaxies have the lowest V_{rot}/σ_g ratio (Kassin et al. 2007, 2012).

- Deep wide-Field WFC3 UV will yield $\lesssim 2000$ UV objects, showing how galaxies disks have grown and stabilized for $0.2 \lesssim z \lesssim 2$.

(4) Late-epoch SF and structural evolution in massive early-type galaxies.



- 10-band WFC3 ERS data measured rest-frame UV-light in nearly all early-type galaxies at $0.3 \lesssim z \lesssim 1.5$ (Rutkowski et al. 2012, ApJS, 199, 4).
 \Rightarrow Most ETGs have continued residual star-formation after they form.
- Can determine their $N(z_{form})$, which resembles the cosmic SFH diagram (Madau et al. 1996), directly constraining the process of galaxy assembly & down-sizing (Kaviraj et al. 2012, MNRAS).
- Deep wide-field WFC3 UV increases sample 10-fold, providing critical UV data to delineate the range in z_{form} for ETGs as function of mass.

(5) Conclusions

For as long as we still have HST, deep wide-field WFC3 UV surveys must be done to address the following critical science questions:

These are critical and unique data in preparation for JWST ($\lambda \gtrsim 0.7 \mu\text{m}$), and to define a $\gtrsim 8$ -meter UV-optical sequel to HST:

- (1) The physics and evolution of SF in low-mass galaxies over the LAST 9 Gyrs: critical benchmark to understand cosmic reionization at $z \gtrsim 6$;
- (2) Evolution of the star/dust/gas mixture in SF regions, and the influence of supernovae and AGN feedback;
- (3) Evolution of young, star-forming sub-galactic clumps induced by mergers or gas accretion, and the growth of stable galaxy disks;
- (4) Late-epoch SF and structural evolution in massive early-type galaxies.

SPARE CHARTS

- References and other sources of material shown:

<http://www.asu.edu/clas/hst/www/jwst/> [Talk, Movie, Java-tool]

<http://www.jwst.nasa.gov/> & <http://www.stsci.edu/jwst/>

Bouwens, R. et al. 2010, ApJ, 709, L133

Bouwens, R. et al. 2011, ApJ, 737, 90

Finkelstein, S. et al. 2012, ApJ, 428, 925

Grazian, A., et al. 2006, A&A, 449, 951

Gardner, J. P., et al. 2006, Space Science Reviews, 123, 485

Guo, et al. 2011, ApJ, 735, 18

Hathi, N., et al. 2010, ApJ, 720, 1708 & AAS 221, 228.06

Kassin, S., et al. 2007, ApJ, 660, L35 & AAS 221, 210.04

Kaviraj, et al. 2012, MNRAS, 428, 925


Reddy, N., et al. 2010, ApJ, 712, 1070

Rutkowski, et al. 2012, ApJS, 199, 4



Ryan, R., et al. 2012, ApJ, 749, 53

Siana, B., et al. 2010, ApJ, 723, 241 & AAS 221, 228.05


Windhorst, R., et al., 2011, ApJS, 193, 27



Lyman break galaxies at the peak of cosmic SF ($z \simeq 1-3$; Hathi et al. 2010)





- Limited flux-range as yet, which limits the M^* , ϕ^* , α -accuracy.
- Deep wide-field WFC3 UV: UV LF's in z -range hard to do from ground.
- Deep wide-field WFC3 UV: significantly improves bright-end, M^* & ϕ^* .



Measured faint-end LF slope evolution (α ; top) and characteristic luminosity evolution (M^* ; bottom) from Hathi et al. 2010 (ApJ, 720, 1708).

- Still poorly determined LF parameters at $1 \lesssim z \lesssim 3$, when most stars born:
- Deep wide-field WFC3 UV imaging will vastly improve on this.



WFC3 ERS 10-band redshift estimates accurate to $\lesssim 4\%$ with small systematic errors (Hathi et al. 2010, 2012), resulting in a reliable $N(z)$:

- Measure masses of faint galaxies to $AB=26.5 \text{ mag}$, tracing the process of galaxy assembly: downsizing, merging, (& weak AGN growth?).

ERS shows WFC3's new panchromatic capabilities on galaxies at $z \simeq 0-8$:

- Deep wide-field WFC3 UV will significantly improve SED-fits & photo- z 's, both in rms and catastrophic failures (Cohen et al. 2012).