

The Infrared Universe Beyond Hubble: The James Webb Space Telescope in 2022!


Rogier Windhorst (ASU) — JWST Interdisciplinary Scientist

S. Cohen, R. Jansen, T. Carleton, & R. O'Brien (ASU), C. Conselice (UK), S. Driver (OZ), & H. Yan (U-MO)

+ the PEARLS team of 85 scientists world-wide



1973~2030⁺(?);



1996~2042(?);

Talk at the Earth and Space Exploration Open House, Arizona State University, Tempe, AZ

Friday November 4, 2022. All presented materials are ITAR-cleared.

Outline

- (1) Update on the James Webb Space Telescope (JWST), 2022.
- (2) Webb's first images: the “Cosmic Circle of Life”
- (3) Summary and Conclusions
- (4) What Hubble has done: Galaxy Assembly & SMBH Growth
- (5) How can JWST measure Star-formation & Earth-like exoplanets?

Thank you, Europe & ESA, for your very significant work on JWST!



Sponsored by NASA/HST & JWST

Talk is on: http://www.asu.edu/clas/hst/www/jwst/jwsttalks/sese_OpenHouse_jwst_nov22.pdf



Edwin P. Hubble (1889–1953) — Carnegie astronomer



James E. Webb (1906–1992) — Second NASA Administrator

Hubble: Concept in 1970's; Made in 1980's; Operational 1990– \gtrsim 2022?.

JWST: The infrared sequel to Hubble from 2021–2026 (–2031?).


**JWST primary
mirror**



**Hubble primary
mirror**

JWST $\simeq 2.5 \times$ larger than Hubble, so at $\sim 2.5 \times$ larger wavelengths:
JWST has the same resolution in the near-IR as Hubble in the optical.

(1) Update of the James Webb Space Telescope as of 2022



- A fully deployable 6.5 meter (25 m^2) segmented IR telescope for imaging and spectroscopy at $0.6\text{--}28 \mu\text{m}$ wavelength, launched Dec. 25, 2021.
- Nested array of sun-shields to keep its ambient temperature at 40 K, allowing faint imaging (31.5 mag \sim 1 FF from Moon), and spectroscopy.

THE JAMES WEBB SPACE TELESCOPE

JWST LAUNCH


- LAUNCH VEHICLE IS AN ARIANE 5 ROCKET, SUPPLIED BY ESA
- SITE WILL BE THE ARIANESPACE'S ELA-3 LAUNCH COMPLEX NEAR KOUROU, FRENCH GUIANA



ARIANESPACE - ESA - NASA

- The JWST launch weight is $\lesssim 6500$ kg, and it was launched to L2 with an ESA Ariane-V launch vehicle from Kourou in French Guiana.


(1a) How did JWST travel to its L2 orbit?



- After launch on Dec. 25, 2022 with an ESA Ariane-V, JWST will orbit around the Earth–Sun Lagrange point L2, 1.5 million km from Earth.
- JWST can cover the whole sky in segments that move along with the Earth, observe $\gtrsim 70\%$ of the time, and send data back to Earth every day.


- (1b) How was JWST automatically deployed?

OTE in folded configuration



- During its two month journey to L2, JWST will be automatically deployed, its instruments will be cooled, and be inserted into an L2 orbit.
- The entire JWST deployment sequence is being tested several times on the ground — but only in 1-G: component and system tests in 2014–2019 at GSFC (MD), Northrop (CA), and JSC (Houston).
- Component fabrication, testing, & system integration: 18 out of 18 flight mirrors done in 2015, and meet the 40K specifications (2017).

Actuators for 6 degrees of freedom rigid body motion



Actuator
development
unit

Lightweighted
Beryllium Mirror


Actuator for radius
of curvature adjustment

Active mirror segment support through “hexapods”, similar to Keck.

Redundant & doubly-redundant mechanisms, quite forgiving against failures.




JWST Hardware Status



2021: 100% of launch mass designed and built (100% weighed).³



TELESCOPE ARCHITECTURE





3/31/11

2014–2021: Complete system integration at GSFC and Northrop.



JWST Hardware Progress




JWST remains on track for an October 2018 launch within its replan budget guidelines


29

July 2014: ● Secondary Mirror Support deployment successfully tested.
2015: ● Engineering sunshield successfully deployed at Northrop (CA).

Much progress has been made in OTE integration



Where we were at last month's call



Current: all 18 PMSAs installed, liquid-shim-cured, & metrologized. Alignments meet specifications, and actuator motions verified

Big milestone!



8 February 2016 JWST Monthly Telecon 8

JWST lifetime: Requirement: 5 yrs; Goal: 10 yrs; Propellant: 20+ yrs!




NASA team-work to take JWST mirror covers off!




JWST being tilted into the right position



Webb mirrors finally mounted and ready!




JWST stowed for further instrument mounting



All Instruments Integrated



Fine Guidance Sensor



Mid-Infrared Instrument





Near Infrared Camera



Near Infrared Spectrometer


JWST: A Product of the Nation






- JWST hardware made in 27 US States: 100% of launch-mass finished.
- Ariane V Launch & NIRSpec provided by ESA; & MIRI by ESA & JPL.
- JWST Fine Guider Sensor + NIRISS provided by Canadian Space Agency.
- JWST NIRCam made by UofA and Lockheed.

Thank you, Europe & ESA, for your very significant work on JWST!

Micro Shutters



Metal Mask/Fixed Slit






April 2017: Last portrait of JWST at Goddard Space Flight Center (MD).



SMSS Deployment Sequence (1)




190812 JWST Monthly Telecon 8

July 2019: Full 1-G deployment of JWST secondary mirror (SM) .



SMSS Deployment Sequence (2)




190812 JWST Monthly Telecon 9

July 2019: Full 1-G deployment of JWST secondary mirror (SM) ..



SMSS Deployment Sequence (3)



190812 JWST Monthly Telecon 10



July 2019: Full 1-G deployment of JWST secondary mirror (SM) ...



May 2017: JWST in enclosure at Johnson Space Center in Houston.

Program Update: OTIS

NORTHROP GRUMMAN



Program Updates: Spacecraft and Sunshield

NORTHROP GRUMMAN




2017–2018: JWST Flight Sunshield assembled and tested at Northrop.



SCE to Elephant Stand

NORTHROP GRUMMAN



Aug. 2019: Stowed flight sunshield before integration with JWST OTE.




SMSS Deployment

NORTHROP GRUMMAN



190812 JWST Monthly Telecon 39

Aug. 2019: OTE before final integration with Sunshield & spacecraft.



Aug. 2019: JWST OTE+ISIM lowered into Sunshield+Spacecraft



August 2019: JWST OTE+ISIM integrated with Sunshield+Spacecraft!



August 2019: JWST OTE+ISIM integrated with Sunshield and Spacecraft!



Meet the JWST Observatory 1



See NASA Press Release here:

<https://www.nasa.gov/feature/goddard/2019/nasa-s-james-webb-space-telescope-has-been-assembled-for-the-first-time>

190909-JWST Monday Telecon 11

August 2019: JWST OTE+ISIM integrated with Sunshield and Spacecraft!



Solar Array Deployment 1

Five Panel Sunshield

Stowed

Offloading System



200511 JWST Monthly Telecon 12

May 2020: Ready for Solar Array deployment test



Solar Array Deployment 2



200511 JWST Monthly Telecon 13

May 2020: Solar Array deployment with gravity off-loading




Solar Array Deployment 3



200511 JWST Monthly Telecon 14

May 2020: Solar Array fully deployed and motor tested in 1G

7/26/20: Solar Array Installed for Environments



5

Approved for Public Release; NG20-1503
200810 JWST Monthly Telecon Man.

May 2020: Solar Array as installed on JWST Observatory



5/28/20: DTA Deployment



Approved for Public Release; NG20-106
200608 JWST Monthly Telecon 26

June 2020: Deployable Tower Assembly test



5/28/20: DTA Deployment



Approved for Public Release; NG20-106
200608 JWST Monthly Telecon 27

June 2020: Deployable Tower Assembly test with gravity off-loading.



5/29/20: DTA Deployment




Approved for Public Release; NG20-106
200608 JWST Monthly Telecon 28

June 2020: Deployable Tower Assembly motor tested in 1G



Transport to the Large Acoustic Test Facility



En route through the Space Park, Credit: NGSS



Arriving at the LATF Airlock, Credit: NGSS
200914 JWST Monthly Telecon 12

Aug 2020: Transport of JWST into Northrop acoustic chamber



(beautiful)
**The James Webb
Space Telescope**

Stowed for Launch



210913 JWST Monthly Telecon 18

Sept. 2021: JWST ready and stowed for shipping to Kourou



Dec. 9, 2021: JWST transport in Kourou to Ariane Rocket Assembly Building



Webb is finally launched from Kourou on December 25, 2021!



•LIVE




Feb. 2022: Webb seen shortly after launch over Africa using the Ariane V camera.

PRIMARY MIRROR SELFIE




IMAGE ARRAY



Feb. 2022: Webb's first selfie (left) and First Light raw image (right).

COMPLETED SEGMENT ALIGNMENT



COMPLETED IMAGE STACKING




Webb's first segment alignment (left) and first image stack (right).

TELESCOPE ALIGNMENT EVALUATION IMAGE




March 16, 2022: Webb's first fully focused image publicly released !!
Note the plethora of faint galaxies — Webb's looking back in time!

WEBB TELESCOPE IMAGE SHARPNESS CHECK



April 28, 2022: Webb's first fully focused images in all four instruments:
a dense star field in the Large Magellanic Cloud in the South Ecliptic Pole!
(NIRSpec: $1.1 \mu\text{m}$; NIRISS: $1.5 \mu\text{m}$; NIRCam: $2.0 \mu\text{m}$; MIRI $7.7 \mu\text{m}$).




SPITZER IRAC 8.0μ



WEBB MIRI 7.7μ


May 9, 2022: Webb's 7.7μ m MIRI image compared to Spitzer 8.0μ m:
Same dense star field in the Large Magellanic Cloud in the South Ecliptic Pole




July 6, 2022: 32-hr Fine Guidance Sensor deepest wide-band near-IR image

(bright star: 9.2 mag 2MASS 16235798+2826079).



<https://blogs.nasa.gov/webb/2022/07/06/webbs-fine-guidance-sensor-provides-a-preview/>



... Webb reveals the faintest galaxies in the near-infrared!



Webb can see the faintest galaxies to the level where the universe has many
many “billions and billions” !



Webb first NIRSpec near-IR spectra of \sim 100 faint stars near Galactic Center

Webb can take spectra of many 1000's of faint galaxies revealing their distances and chemical composition.



Hubble WFPC2 Eagle Nebula (1995) compared to JWST NIRCam (2022):

- The cradle of cosmic star-formation: NIRCam peers through the dust!
- The 1995 Hubble WFPC2 image (left) was made by Prof. Jeff Hester and Paul Scowen at ASU. It made it onto a US postage stamp!



Webb's MIRI shows the hauntingly beautiful cosmic dust pillars (8–15 μm)

JAMES WEBB SPACE TELESCOPE

TARANTULA NEBULA | NGC 2070



NIRCam Filters


F090W F200W F335M F444W

50 LIGHT-YEARS

Tarantula Nebula “30 Doradus” in Large Magellanic Cloud (163,000 lyrs away)
Cradle of cosmic star-formation: massive stars trigger formation of sun-like stars

JAMES WEBB SPACE TELESCOPE

TARANTULA NEBULA | NGC 2070



MIRI Filters F770W F1000W F1280W F1800W

midIR shows Poly-Aromatic Hydro-carbons: leftover from previous massive stars
Webb's MIRI shows the chemistry of star-formation: cosmic star-dust!



“Cosmic Cliffs” of star-formation in the Carina Nebula (NIR; 7600 light-years).


You will be witnessing the “Cosmic Circle of Life” ...



Cosmic Cliffs of Star-formation in Carina Nebula (NIR+MIR).

Compared to optical+near-IR, mid-IR sees “Cradle of Cosmic Star-formation”

Deep inside the gas and dust, mid-IR reveals birth of young Sun-like stars.



Southern Ring Nebula (Near-IR+Mid-IR; 2500 light-years):

- You **are** witnessing the “Cosmic Circle of Life” here ...
- This is a Sun-like star expelling its outer layers in retirement ...
- It has exhausted its hydrogen and helium as nuclear fuel ...
and expanded to $100\times$ its current size, engulfing the Earth.




This is how our Sun *will* come to an end in 5 Billion years (near-IR).
“... for dust thou art, and unto dust shalt thou return” (Genesis 3:19).





From gas expelled by previous sun-like stars, new stars are born (mid-IR).
And thanks to the dust they expelled, new stars will form with planets ...

Mars

James Webb Space Telescope
NIRCam - September 5, 2022




Simulated Mars image with base maps
from NASA and MOLA data



NASA, ESA, CSA, STScI, MARS JWST/GTO team

Mars' surface with NIRCam: From “hot” to “cold” in the infrared!

ATMOSPHERE COMPOSITION




WEBB
SPACE TELESCOPE

Mars atmosphere NIRSpec spectrum: Plenty of Carbon Dioxide ...
but the search is much harder for Water vapor and Carbon Monoxide



Aug. 2022: JWST NIRcam image of the planet Jupiter: it has beautiful aurorae at its North and South pole — very strong magnetic field!
The Great “Red” Spot: A giant 4-century storm $2 \times$ Earth’s diameter.



NIRCam family portrait of Neptune with 7 of its Moons
Moon Triton is brighter, since methane darkens Neptune's atmosphere



Closeup of planet Neptune with Webb's NIRCam

- Giant planets with (dim) rings more common those than without rings!

Star
HIP 65426

Exoplanet
HIP 65426 b

JWST

NIRCam

F300M

NIRCam

F444W

MIRI


F1140C

MIRI

F1550C

Webb 3–15 micron exoplanet images (10 Jupiter masses; 15 Myr young!)

ATMOSPHERE COMPOSITION



WEBB
SPACE TELESCOPE


Hot exoplanet WASP-96b orbiting a Sun-like star (1150 light-years):

- Near-IR spectrum shows characteristic features of water (steam !).
- It has a temperature of 1000 F and is half Jupiter in mass.
- Webb will scan Earth-like exoplanets for building blocks of life.



Stephan's Quintet: 4 colliding galaxies (40 M-lyr; left spiral is foreground).

- These major “Cosmic Trainwrecks” are much more common in the past.
- Sun-like stars formed in aftermath of minor “Cosmic Fender-benders” .




Stephan's Quintet: 4 colliding galaxies at 40 million light-years (Mid-IR):


- Mid-IR shows molecular gas being pulled out during collision.
- Gravity from collision in top galaxy feeds the Beast: central black hole!




Spiral Galaxy overlapping Elliptical: Tracing cosmic dust (Keel⁺ 22) ...




... and the elliptical also lenses a galaxy seen 2 Byrs after Big Bang!



July 11, 2022: 12-hr Webb Deep Field on galaxy cluster SMACS 0723



Hubble image of SMACS 0723 – Webb sees the dawn of galaxy formation!



Compared to near-IR (right), mid-IR sees some **very red** objects (left):

- These may be gravitationally lensed galaxies seen in the first 1–2 Byrs.
 - Cluster galaxies already are ~ 9 Byrs old, seen at 4.5 Blyr distance!
- ⇒ Sun was just born when these *old* galaxies emitted their light!




JD 1

JD 2

JD 3


Cluster MACS0647 triply lensed a galaxy 0.4 Byrs after BB! (Hsiao, Coe⁺ 22)




Monster cluster El Gordo distorts distant galaxies into “pencils” (Diego⁺²²)




and El Gordo makes a super-lens “El Anzuelo” — Einstein’s fishhook!



Many Webb images are deep fields looking back into the first Billion years.




Irac Dark Field: spot on the Ecliptic Pole Webb can always see (Yan⁺ 22).
Not very many Galactic stars in the foreground (incl some brown dwarfs).
Distant galaxies are between 1 and 13 Byrs looking back in time.




Many deep field galaxies have tidal tails ... galaxies torn by gravity

For details, see: Windhorst et al. (<https://arxiv.org/abs/2209.04119>)



Many deep field galaxies have tidal tails ... galaxies torn by gravity ...
What can I say: Gravity bends!




Many deep field galaxies have tidal tails ... galaxies torn by gravity ...
What can I say: Gravity bends!
and black holes bend harder ... any black hole questions?

SDSS J165202.64+172852.3

MOTIONS OF GAS AROUND AN EXTREMELY RED QUASAR

Hubble ACS + WFC3 Imaging


Webb NIRSpec IFU Spectroscopy



WEBB
SPACE TELESCOPE

NIRSpec spectral cube of a luminous quasar seen 2.2 Byrs after Big Bang.
Colors indicate 3 companion galaxies falling into the quasar host galaxy.
● In the first 2 billion years big galaxies were swallowing little ones!


3) How can Webb measure First Light: What to expect in (Ultra)Deep Fields?



- Can't beat redshift: to see First Light, must observe near-mid IR.
⇒ This is why JWST needs NIRCam at 0.8–5 μm and MIRI at 5–28 μm .


13 filter Hubble UltraDeep Field:

UV-Blue emphasized.




○ z=7-8, ○ z=9, ○ z=10-12.


Hubble UltraDeep Field: Red-IR emphasized.



13 filter Hubble UltraDeep Field:

False-color image: sky is half covered by galaxies.





Conclusion: JWST First Light strategy must consider three aspects:

- (1) The earliest objects (first 0.5 Byr) are very rare and hard to find.
- (2) Cannot-see-the-forest-for-the-trees effect [“Confusion” limit]:
Background objects blend into foreground objects because of their density.
- (3) House-of-mirrors effect [“Gravitational Amplification”]:
 - Lensing is needed to see what Einstein thought was impossible to observe!

(3) Summary and Conclusions

(1) Webb was successfully built, tested and finally launched in Dec. 2021.

(2) Webb was designed to map the epochs of First Light, Galaxy Assembly & Super Massive Black Hole-growth in detail:

- Formation of the first stars and star-clusters after 0.2 Byr.
- How galaxies formed and evolved over 13.5 Billion years.

(3) Webb's first images trace the "Cosmic Circle of Life":

- Formation and evolution of stars and dust over cosmic time.
- How dust helped form exoplanets and building blocks for life.

(4) Webb will have a major impact on astrophysics this decade:

- IR sequel to HST starting 2022: Training next generation researchers.

SPARE CHARTS

- References and other sources of material shown:

<http://ahah.asu.edu/> [Appreciating Hubble at Hyperspeed]

<http://ahah.asu.edu/download.html> [Download Java–tool]

<http://ahah.asu.edu/clickonHUDF/index.html> [Clickable map]

<http://www.jwst.nasa.gov/> & <http://www.stsci.edu/jwst/>

<https://blogs.nasa.gov/webb/>

<http://https://www.nasa.gov/webbfirstrimages>

<http://www.webbcompare.com/index.html>

Gardner, J. P., et al. 2006, *Space Science Reviews*, 123, 485–606


Mather, J., & Stockman, H. 2000, *Proc. SPIE Vol. 4013*, 2

Windhorst, R., et al. 2008, *Advances in Space Research*, 41, 1965

Windhorst, R., et al., 2011, *ApJS*, 193, 27 ([astro-ph/1005.2776](http://arxiv.org/abs/1005.2776)).

What the Scientists See:

What the Project Manager Sees:




The Happy Balance

Found at GP3A.CX

Any (space) mission is a balance between what science demands, what technology can do, and what budget & schedule allows ... (courtesy Prof. R. Ellis).

(4) What Hubble has done: Panchromatic High-Throughput Camera



HST WFC3 and its IR channel: a critical pathfinder for JWST science.

(4) Hubble WFC3: Measuring Galaxy Assembly and SMBH Growth?



10 filters with Hubble WFC3 & ACS reaching AB=26.5–27.0 mag over 40 arcmin² with 0.07–0.15" images from 0.2–1.7μm (UVUBVizYJH). JWST adds 0.05–0.2" FWHM imaging to AB≈31.5 mag (1 FF) at 1–5μm, with 0.2–1.2" images at 5–29μm, tracing young+old stars & dust.

Black Hole growth — Waves that happen in Nature: 1) Sounds Waves:





In solids: Earthquakes



In liquids: Surf!




In gasses: Sound



2) Electromagnetic Waves

3) In space-time: Gravity Waves

Sept. 2015: LIGO added Gravity Waves as a new way to observe Nature!



(1) LIGO first observed Gravitational Waves on Sept. 14, 2015.

(2) These were caused by two merging ($29+36 M_{\odot}$) black holes about 1 Gyr ago!

- $E=Mc^2$: $3 M_{\odot}$ was converted to energy in a fraction of a second!

Visible

Infrared




30 Doradus Nebula and Star Cluster

Hubble Space Telescope • WFC3/UVIS/IR

NASA, ESA, F. Paresce (INAF-IASF, Italy), and the WFC3 Science Oversight Committee

STScI-PRC09-32b

30 Doradus: Giant young star-cluster in Large Magellanic Cloud (150,000 ly), triggering birth of Sun-like stars (and surrounding debris disks).



Ordinary massive stars ($10\text{--}30 M_{\odot}$) leave modest black holes ($\sim 3\text{--}10 M_{\odot}$).



Ordinary massive stars ($10\text{--}30 M_{\odot}$) leave modest black holes ($\sim 3\text{--}10 M_{\odot}$).

Conclusion 1: Most low-mass black holes today are small, slow eaters:



- 29–36 M_{\odot} blackholes may be leftover from First Stars (first 500 Myr).
- Likely too massive to be leftover from ordinary Supernova explosions, ...
- How come only now seen merging by LIGO (12.5 Byr after BB)?
- They were likely not fast & efficient eaters, but slow and messy ...

Elliptical galaxy M87 with Active Galactic Nucleus (AGN) and relativistic jet:



"For God's sake, Edwards. Put the laser pointer away."

The danger of having Quasar-like devices too close to home ...

They are EXTREMELY bright sources if viewed “down-the-pipe”.

~0.5% of the baryonic mass, but produce most of the photons!








Figure 15. Averages of the three fiducial images of M87 for each of the four observed days after restoring each to an equivalent resolution, as in Figure 14. The indicated beam is $20 \mu\text{as}$ (i.e., that of DIFMAP, which is always the largest of the three individual beams).

2019 discovery of Black Hole Shadow in M87 by Event Horizon Telescope:
M87 at 55 Mlyr distance has a black hole mass of $\sim 6.5 \times 10^9 M_\odot$!


Centaurus A
NGC 5128
HST WFC3/UVIS





Blue=X-rays; White=Optical; Orange=Radio

- Quasars: Centers of galaxies with feeding supermassive blackholes:




- Hubble IR-images of the most luminous Quasar known in the universe.
- Seen at redshift 6.42 (universe $7.42 \times$ smaller than today), 900 Myr old!
- Contains 10^{14} solar luminosities within a region as small as Pluto's orbit!
- A feeding monster blackhole ($> 3 \times 10^9$ solar mass) 900 Myr after BB!

Conclusion 2: Supermassive black holes started early & were very rapid eaters:



- Massive galaxies today contain a super-massive blackhole, no exceptions!
- Masses $\sim 3 \times 10^9$ solar, leftover from the First Stars (first 500 Myr)?
- Must have fed enormously rapidly in the first 1 Byr after the Big Bang.
- Were eating *cat*-astrophically (and secretly) until they ran out of food ...
- JWST can image the First Quasars to $z \gtrsim 10$ (*if* we can find them).



Will this ever happen to our own Galaxy?

YES! Hubble showed no lateral motion of Andromeda:
Approaches at -110 km/s.


Hence, Andromeda will merge with Milky Way!

The two blackholes (10^6 – 10^7 suns) will also merge!

Not to worry: only 4–5 Byr from today!



Illustration Sequence of the Milky Way and Andromeda Galaxy Colliding

(5) How can JWST measure Star-Formation and Earth-like exoplanets?




NGC 3603: Young star-cluster triggering star-birth in “Pillars of Creation”

Pillar and Stellar Jet in the Carina Nebula
HST WFC3/UVIS




Fomalhaut
HST ACS/HRC




HST/ACS Coronagraph imaging of planetary debris disk around Fomalhaut:
First direct imaging of a moving planet forming around a nearby star!
JWST can find such planets much closer in for much farther stars.


Exoplanet HR 8799 System



NASA, ESA, and R. Soummer (STScI)



HST NICMOS with additional processing



HR 8799 planet orbits

Relative size of
Neptune's orbit

STScI-PRC11-29


HST/NICMOS imaging of planetary system around the (carefully subtracted) star HR 8799: Direct imaging of planets around a nearby star.

Press release: <http://hubblesite.org/newscenter/archive/releases/2011/29/>


JWST can find such planets much closer in for much farther-away stars.

Schematic of Transit and Eclipse Science

Seager & Deming (2010, ARAA, 48, 631)




Timeline of a Transit Observation



6


13

JWST can do very precise photometry of transiting Earth-like exoplanets.



JWST IR spectra can find water and CO₂ in (super-)Earth-like exoplanets.

Transit Spectrum of Habitable “Ocean Planet”



17

JWST IR spectra can find water and CO_2 in transiting Earth-like exoplanets.