

422 Exam 2

⚠ This is a preview of the published version of the quiz

Started: Jan 20 at 5:25pm

Quiz Instructions

This is the second midterm. It covers material post midterm 1. You have 2 hours from the time that you start the exam to submit. You must show your work on this exam. You are welcome to use a calculator, but you cannot use solvers (think eigenvalue solvers).

Question 1

5 pts

Why did we reformulate the SVM optimization using Lagrange multipliers? In other words, what was the point of moving to the dual form of the SVM?

Edit View Insert Format Tools Table

12pt ▾ Paragraph ▾ | **B** *I* U A ▾  ▾ τ^2 ▾ | ⋮

p



0 words



Question 2**5 pts**

If I am going to apply PCA to the following data, what size will my covariance matrix be? (s_i stands for sample i)

$$s_1 = (0, 1, 2, 1)$$

$$s_2 = (1, 3, 0, 2)$$

$$s_3 = (3, 5, 1, 1)$$

$$s_4 = (2, 2, 1, 4)$$

$$s_5 = (10, 1, 3, 5)$$

Edit View Insert Format Tools Table

12pt ▾ Paragraph ▾ | **B** *I* U A ▾  ▾ T² ▾ | ⋮

p



0 words

**Question 3****5 pts**

In a setting where there is no human oversight, is it more important to have high precision or high recall? Explain.

Edit View Insert Format Tools Table

12pt ▾ Paragraph ▾ | **B** *I* U A ▾  ▾ τ^2 ▾ | ⋮

p



0 words



Question 4

5 pts

I have a large dataset. Should I separate the data into Train, Val & Test? Or should I use 5-fold Cross Validation? Explain why you chose one over the other. Would your answer change if I had a small dataset? Why or why not?

Edit View Insert Format Tools Table

12pt ▾ Paragraph ▾ | **B** *I* U A ▾  ▾ τ^2 ▾ | ⋮

p



0 words

**Question 5****5 pts**

CNNs: If you apply a $J \times K$ filter to an image of size $N \times M$. What size is the resulting feature map? Assume there are p pixels of padding in each direction and a stride $s = 1$. Give an expression for the size of the resulting feature map in terms of these variables.

Edit View Insert Format Tools Table

12pt ▾

Paragraph ▾

B*I*UA ▾ ▾ T^2 ▾

p



0 words



Question 6**10 pts**

True/False: A hard-margin SVM will always give the same results assuming the training data is the same. What about a soft-margin SVM? Briefly explain.

Edit View Insert Format Tools Table

12pt ▾ Paragraph ▾ | **B** *I* U A ▾  ▾ τ^2 ▾ | ⋮

p



0 words


**Question 7****10 pts**

We want to perform Adaboost to train a strong classifier using many weak classifiers. In this case, our weak classifiers are going to be axis-parallel lines (single lines that are either vertical or horizontal) that minimize the weighted training error. Everything on one side of the line will be classified as positive and on the other side will be negative. There are three parts to this question, so make sure to answer all three.

a) Using the dataset below, draw the decision boundary learned by the first classifier in the Adaboost algorithm.

b) Circle the point(s) with the highest weights at the beginning of the second iteration of Adaboost.

c) Draw the decision boundary learned by the the second classifier in the Adaboost Algorithm.



Edit View Insert Format Tools Table

12pt ∨ Paragraph ∨ | **B** *I* U A ∨ ∨ T² ∨ | ∷

p



0 words



Question 8

15 pts





Given the following kernel, find the function $\phi(x)$ such that $K(x, z) = \phi(x) \cdot \phi(z)$. Each sample (x and z) is one dimensional (they have one feature).

$$K(x, z) = (1 + xz + x^2 z^2)^2$$

Edit View Insert Format Tools Table

12pt Paragraph | **B** *I* U A   T² | ⋮


p

  | 0 words |   ⋮

Question 9

20 pts

Given the following data, what w and b would a hard-margin SVM return? Red filled circles are the negative class. Green empty circles are the positive class. Assume all points lie on integer coordinates.



Edit View Insert Format Tools Table

12pt Paragraph | **B** *I* U A τ^2 :

p



0 words



Question 10

20 pts


Given the following neural network, give the expression for $\frac{\partial L}{\partial w_{11}^1}$. Assume $L = (y - \hat{y})^2$, $h_i^1 = e^{w_{i1}^1 \cdot x}$ (that is, the first hidden layer applies e to its input), and there is no nonlinearity at the second hidden layer nor at the output.

Remember the notation for w_{ij}^k means the k^{th} layer of weights going from the i^{th} node to the j^{th} node.

Write out the expression for $\frac{\partial L}{\partial w_{11}^1}$ using partial derivatives (e.g.

$\frac{\partial L}{\partial w_{11}^1} = \frac{\partial L}{\partial \text{something}} \frac{\partial \text{something}}{\partial \text{somethingelse}} \dots$) and then identify each partial derivative





separately (e.g. $\frac{\partial L}{\partial \text{something}} = \text{blahblahblah}$, $\frac{\partial \text{something}}{\partial \text{somethingelse}} = \text{yadayada}$). This is the only way you will receive partial credit!!!



Edit View Insert Format Tools Table

12pt ▾ Paragraph ▾ | **B** *I* U A ▾  ▾ T² ▾ | ⋮

p

  | 0 words |   ⋮

Not saved

Submit Quiz