

$$x \text{ , } I_{ij} \text{ , } e^{x^2}$$

$$L=I_{ij}\cdot\cos(\theta)+R^2\tag{1}$$

$$k=\frac{ab}{cd}\tag{2}$$

$$\begin{aligned} f(x) &= x^2 + 2x + 1 \\ &= (x+1)^2 \end{aligned}\tag{3}$$

$$|y| = \begin{cases} y & \text{if } y \geq 0 \\ -y & \text{if } y < 0 \end{cases}$$

$$f(x_i)=\begin{cases} \frac{x_i^2}{\sigma^2} & \text{if } x_i \geq 0 \\ \frac{|x_i|}{\sigma} & \text{if } x_i < 0 \end{cases}$$

$$T=\sum_{i=1}^n(x_i-\mu)^2=\sum_{i=1}^nx_i^2-2\mu\sum_{i=1}^nx_i+n\mu^2\tag{4}$$

$$N(x_i,\mu,\sigma)=\frac{1}{\sqrt{2\pi}\sigma}e^{-\frac{(x_i-\mu)^2}{2\sigma^2}}$$

$$g(x_i)=\begin{cases} \frac{x_i+1}{x_i-1} & \text{if } x_i \neq 1 \\ 0 & \text{if } x_i = 1 \end{cases}$$

$$g(x_i)=\begin{cases} \frac{x_i+1}{x_i-1} & \text{if } x_i \neq 1 \\ 0 & \text{if } x_i = 1 \end{cases}\tag{5}$$