

(1) European Space Agency. (2017). Land Cover CCI Product User Guide Version 2.0. Retrieved from www.esa-landcover-cci.org

(2) Tadono, T., Nagai, H., Ishida, H., Oda, F., Naito, S., Minakawa, K., & Iwamoto, H. (2016). GENERATION OF THE 30M-MESH GLOBAL DIGITAL SURFACE MODEL BY ALOS PRISM. *ISPRS - International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences, XLI-B4*, 157-162. doi:10.5194/isprs-archives-XLI-B4-157-2016

(3) Marsh, C. J., Sica, Y. V., Burgin, C. J., Dorman, W. A., Anderson, R. C., del Toro Mijares, I., . . . Jetz, W. (2022). Expert range maps of global mammal distributions harmonised to three taxonomic authorities. *Journal of Biogeography, 49*(5), 979-992. doi:https://doi.org/10.1111/jbi.14330

(4) Schug, F., Bar-Massada, A., Carlson, A. R., Cox, H., Hawbaker, T. J., Helmers, D., . . . Radeloff, V. C. (2023). The global wildland–urban interface. *Nature*. doi:10.1038/s41586-023-06320-0

(5) Bauer-Marschallinger, B., Sabel, D., & Wagner, W. (2014). Optimisation of global grids for high-resolution remote sensing data. *Computers & Geosciences, 72*, 84-93. doi:https://doi.org/10.1016/j.cageo.2014.07.005

(6) *Schiavina M., Freire S., Carioli A., MacManus K. (2023):* GHS-POP R2023A - GHS population grid multitemporal (1975-2030). European Commission, Joint Research Centre (JRC)
PID: http://data.europa.eu/89h/2ff68a52-5b5b-4a22-8f40-c41da8332cfe
doi:10.2905/2FF68A52-5B5B-4A22-8F40-C41DA8332CFE