

Residue Classes

Dr. Odelu Vanga

Computer Science and Engineering
Indian Institute of Information Technology Sri City

odelu.vanga@iiits.in

Today's Objectives

- Residue Classes

Today's Objectives

- Residue Classes
- Finding Inverse Modulo m

Today's Objectives

- Residue Classes
- Finding Inverse Modulo m
- General Caesar Cipher

Today's Objectives

- Residue Classes
- Finding Inverse Modulo m
- General Caesar Cipher
- Affine Cipher

Residue Classes

Definition

If m is a positive integer and m divides $(b - a)$, then we say that

- a and b are congruent modulo m
- we write $a \equiv b \pmod{m}$

Residue Classes

Definition

If m is a positive integer and m divides $(b - a)$, then we say that

- a and b are congruent modulo m
- we write $a \equiv b \pmod{m}$

Examples:

- $3 \equiv 9 \pmod{6}$, since 6 divides $9 - 3 = 6$

Residue Classes

Definition

If m is a positive integer and m divides $(b - a)$, then we say that

- a and b are congruent modulo m
- we write $a \equiv b \pmod{m}$

Examples:

- $3 \equiv 9 \pmod{6}$, since 6 divides $9 - 3 = 6$
- $-2 \equiv 28 \pmod{5}$, since 5 divides $28 - (-2) = 30$

Residue Classes

Definition

If m is a positive integer and m divides $(b - a)$, then we say that

- a and b are congruent modulo m
- we write $a \equiv b \pmod{m}$

Examples:

- $3 \equiv 9 \pmod{6}$, since 6 divides $9 - 3 = 6$
- $-2 \equiv 28 \pmod{5}$, since 5 divides $28 - (-2) = 30$
- $0 \equiv -666 \pmod{3}$, since 3 divides $-666 - 0 = -666$

Residue Classes

Definition

If m is a positive integer and m divides $(b - a)$, then we say that

- a and b are congruent modulo m
- we write $a \equiv b \pmod{m}$

Examples:

- $3 \equiv 9 \pmod{6}$, since 6 divides $9 - 3 = 6$
- $-2 \equiv 28 \pmod{5}$, since 5 divides $28 - (-2) = 30$
- $0 \equiv -666 \pmod{3}$, since 3 divides $-666 - 0 = -666$

If m does not divide $b - a$, we say a and b are not congruent mod m , and write $a \not\equiv b \pmod{m}$

Residue Classes

Definition

If m is a positive integer and m divides $(b - a)$, then we say that

- a and b are congruent modulo m
- we write $a \equiv b \pmod{m}$

Examples:

- $3 \equiv 9 \pmod{6}$, since 6 divides $9 - 3 = 6$
- $-2 \equiv 28 \pmod{5}$, since 5 divides $28 - (-2) = 30$
- $0 \equiv -666 \pmod{3}$, since 3 divides $-666 - 0 = -666$

If m does not divide $b - a$, we say a and b are not congruent mod m , and write $a \not\equiv b \pmod{m}$

- $2 \not\equiv 7 \pmod{3}$, because 3 does not divide $7 - 2 = 5$

Residue Classes

- $a = q_1 m + r_1$ and $b = q_2 m + r_2$,
where $0 \leq r_1 \leq m - 1$ and $0 \leq r_2 \leq m - 1$.
- $a \equiv b \pmod{m}$ if and only if $r_1 = r_2$.

Residue Classes

- $a = q_1 m + r_1$ and $b = q_2 m + r_2$,
where $0 \leq r_1 \leq m - 1$ and $0 \leq r_2 \leq m - 1$.
- $a \equiv b \pmod{m}$ if and only if $r_1 = r_2$.
- $r_1 = a \pmod{m}$ denotes the remainder.

Residue Classes

- $a = q_1 m + r_1$ and $b = q_2 m + r_2$,
where $0 \leq r_1 \leq m - 1$ and $0 \leq r_2 \leq m - 1$.
- $a \equiv b \pmod{m}$ if and only if $r_1 = r_2$.
- $r_1 = a \pmod{m}$ denotes the remainder.
- Thus, $a \equiv b \pmod{m}$ if and only if $a \pmod{m} = b \pmod{m}$.

Residue Classes

- $a = q_1m + r_1$ and $b = q_2m + r_2$,
where $0 \leq r_1 \leq m - 1$ and $0 \leq r_2 \leq m - 1$.
- $a \equiv b \pmod{m}$ if and only if $r_1 = r_2$.
- $r_1 = a \pmod{m}$ denotes the remainder.
- Thus, $a \equiv b \pmod{m}$ if and only if $a \pmod{m} = b \pmod{m}$.

Definition (Residue Class)

If a is an integer and $a \equiv b \pmod{m}$, we say that b is a residue of $a \bmod m$.

Residue Classes

- $a = q_1 m + r_1$ and $b = q_2 m + r_2$,
where $0 \leq r_1 \leq m - 1$ and $0 \leq r_2 \leq m - 1$.
- $a \equiv b \pmod{m}$ if and only if $r_1 = r_2$.
- $r_1 = a \pmod{m}$ denotes the remainder.
- Thus, $a \equiv b \pmod{m}$ if and only if $a \pmod{m} = b \pmod{m}$.

Definition (Residue Class)

If a is an integer and $a \equiv b \pmod{m}$, we say that b is a residue of $a \pmod{m}$.

- The residue class of a modulo m , denoted \bar{a} , is the collection of all integers congruent to a modulo m .

Residue Classes

- $a = q_1m + r_1$ and $b = q_2m + r_2$,
where $0 \leq r_1 \leq m - 1$ and $0 \leq r_2 \leq m - 1$.
- $a \equiv b \pmod{m}$ if and only if $r_1 = r_2$.
- $r_1 = a \pmod{m}$ denotes the remainder.
- Thus, $a \equiv b \pmod{m}$ if and only if $a \pmod{m} = b \pmod{m}$.

Definition (Residue Class)

If a is an integer and $a \equiv b \pmod{m}$, we say that b is a residue of $a \pmod{m}$.

- The residue class of a modulo m , denoted \bar{a} , is the collection of all integers congruent to a modulo m .
- Observe that $\bar{a} = \{a + km, k \in \mathbb{Z}\}$.

Set of Residue Class

- The residue class of a modulo m , denoted \bar{a} , is the collection of all integers congruent to a modulo m .
- Observe that $\bar{a} = \{a + km, k \in \mathbb{Z}\}$.

Set of Residue Class

- The residue class of a modulo m , denoted \bar{a} , is the collection of all integers congruent to a modulo m .
- Observe that $\bar{a} = \{a + km, k \in \mathbb{Z}\}$.

Set of residue class $\{0, 1, 2, \dots, m-1\}$ modulo m is denoted by \mathbb{Z}_m , that is,

$$\mathbb{Z}_m = \{0, 1, 2, \dots, m-1\}$$

- Addition and Multiplication works exactly like real addition and multiplication, except reduce modulo m .

Set of Residue Class

- The residue class of a modulo m , denoted \bar{a} , is the collection of all integers congruent to a modulo m .
- Observe that $\bar{a} = \{a + km, k \in \mathbb{Z}\}$.

Set of residue class $\{0, 1, 2, \dots, m-1\}$ modulo m is denoted by \mathbb{Z}_m , that is,

$$\mathbb{Z}_m = \{0, 1, 2, \dots, m-1\}$$

- Addition and Multiplication works exactly like real addition and multiplication, except reduce modulo m .
- $11 \times 13 = 143$ in \mathbb{Z}_{16} , and reduce it to modulo 16:

Set of Residue Class

- The residue class of a modulo m , denoted \bar{a} , is the collection of all integers congruent to a modulo m .
- Observe that $\bar{a} = \{a + km, k \in \mathbb{Z}\}$.

Set of residue class $\{0, 1, 2, \dots, m-1\}$ modulo m is denoted by \mathbb{Z}_m , that is,

$$\mathbb{Z}_m = \{0, 1, 2, \dots, m-1\}$$

- Addition and Multiplication works exactly like real addition and multiplication, except reduce modulo m .
- $11 \times 13 = 143$ in \mathbb{Z}_{16} , and reduce it to modulo 16:
 $143 = 8 \times 16 + 15$, so $143 \pmod{16} = 15 \in \mathbb{Z}_{16}$

Definition (Inverse of an element)

Suppose $a \in \mathbb{Z}_m$. The multiplicative inverse of a is an element $a^{-1} \in \mathbb{Z}_m$ such that $aa^{-1} = a^{-1}a = 1 \pmod{m}$

Finding Inverse Modulo m

Theorem (Multiplicative Inverse Modulo m)

a and m relatively primes if and only if a^{-1} modulo m exists

Finding Inverse Modulo m

Theorem (Multiplicative Inverse Modulo m)

a and m relatively primes if and only if a^{-1} modulo m exists

Proof.

Suppose a and m are relatively prime

Then, $GCD(a, m) = 1$

Finding Inverse Modulo m

Theorem (Multiplicative Inverse Modulo m)

a and m relatively primes if and only if a^{-1} modulo m exists

Proof.

Suppose a and m are relatively prime

Then, $GCD(a, m) = 1$

There exists x and y such that $1 = ax + my$

Now apply modulo m , we get $1 = (ax + 0) \pmod{m}$

That is, $1 = ax \pmod{m}$

Means, there exists x such that $ax = 1 \pmod{m}$

Therefore, x is inverse of a modulo m



Example: Finding Inverse Modulo m

Find multiplicative inverse of $a = 8$ modulo $m = 11$

Example: Finding Inverse Modulo m

Find multiplicative inverse of $a = 8$ modulo $m = 11$

Finding $8^{-1} \pmod{11}$ using Euclidean Algorithm

$$m = qa + r$$

$$a = q_1 r + r_1$$

- $11 = (1) \times 8 + 3$
- $8 = (2) \times 3 + 2$
- $3 = (1) \times 2 + 1$
- $2 = (2) \times 1 + 0$

Example: Finding Inverse Modulo m

Find multiplicative inverse of $a = 8$ modulo $m = 11$

Finding $8^{-1} \pmod{11}$ using Euclidean Algorithm

$$m = qa + r$$

$$a = q_1 r + r_1$$

- $11 = (1) \times 8 + 3$
- $8 = (2) \times 3 + 2$
- $3 = (1) \times 2 + 1$
- $2 = (2) \times 1 + 0$

Rewrite

- $3 = 11 - (1) \times 8$
- $2 = 8 - (2) \times 3$
- $1 = 3 - (1) \times 2$

Example: Finding Inverse Modulo m

Find multiplicative inverse of $a = 8$ modulo $m = 11$

Finding $8^{-1} \pmod{11}$ using Euclidean Algorithm

$$m = qa + r$$

$$a = q_1 r + r_1$$

Reverse the process:

- $11 = (1) \times 8 + 3$
- $8 = (2) \times 3 + 2$
- $3 = (1) \times 2 + 1$
- $2 = (2) \times 1 + 0$

Rewrite

- $3 = 11 - (1) \times 8$
- $2 = 8 - (2) \times 3$
- $1 = 3 - (1) \times 2$

Example: Finding Inverse Modulo m

Find multiplicative inverse of $a = 8$ modulo $m = 11$

Finding $8^{-1} \pmod{11}$ using Euclidean Algorithm

$$m = qa + r$$

$$a = q_1 r + r_1$$

Reverse the process:

find $1 = 8x + 11y$ form

- $11 = (1) \times 8 + 3$
- $8 = (2) \times 3 + 2$
- $3 = (1) \times 2 + 1$
- $2 = (2) \times 1 + 0$

Rewrite

- $3 = 11 - (1) \times 8$
- $2 = 8 - (2) \times 3$
- $1 = 3 - (1) \times 2$

Example: Finding Inverse Modulo m

Find multiplicative inverse of $a = 8$ modulo $m = 11$

Finding $8^{-1} \pmod{11}$ using Euclidean Algorithm

$$m = qa + r$$

$$a = q_1 r + r_1$$

Reverse the process:

find $1 = 8x + 11y$ form

- $11 = (1) \times 8 + 3$

- $8 = (2) \times 3 + 2$

- $3 = (1) \times 2 + 1$

- $2 = (2) \times 1 + 0$

Rewrite

- $3 = 11 - (1) \times 8$

- $2 = 8 - (2) \times 3$

- $1 = 3 - (1) \times 2$

$$1 = 3 - (1) \times 2$$

$$= 3 - (1) \times [8 - (2) \times 3]$$

$$= (-1) \times 8 + (3) \times 3$$

$$= (-1) \times 8 + (3) \times [11 - (1) \times 8]$$

$$= (-4) \times 8 + (3) \times 11$$

$$x = -4 \pmod{11} = 7 = 8^{-1} \pmod{11}$$

Finding Inverse

Find Inverse of 7 modulo 26

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Rewrite

$$5 = 26 - (3) \times 7$$

$$2 = 7 - (1) \times 5$$

$$1 = 5 - (2) \times 2$$

Finding Inverse

Find Inverse of 7 modulo 26

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Rewrite

$$5 = 26 - (3) \times 7$$

$$2 = 7 - (1) \times 5$$

$$1 = 5 - (2) \times 2$$

Reverse Process

$$1 = 5 - (2) \times 2$$

$$1 = 5 - (2) \times [7 - (1) \times 5]$$

$$1 = (-2) \times 7 + (3) \times 5$$

$$1 = (-2) \times 7 + (3) \times [26 - (3) \times 7]$$

$$1 = (-11) \times 7 + (3) \times 26$$

$$x = -11 \pmod{26} = 15 = 7^{-1} \pmod{26}$$

Finding Inverse

Find Inverse of 7 modulo 26

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Rewrite

$$5 = 26 - (3) \times 7$$

$$2 = 7 - (1) \times 5$$

$$1 = 5 - (2) \times 2$$

Reverse Process

$$1 = 5 - (2) \times 2$$

$$1 = 5 - (2) \times [7 - (1) \times 5]$$

$$1 = (-2) \times 7 + (3) \times 5$$

$$1 = (-2) \times 7 + (3) \times [26 - (3) \times 7]$$

$$1 = (-11) \times 7 + (3) \times 26$$

$$x = -11 \pmod{26} = 15 = 7^{-1} \pmod{26}$$

Finding Inverse

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

Finding Inverse

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$
- $5 = 5 \times 1 + 0$

Finding Inverse

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$
- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

Finding Inverse

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

Finding Inverse

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Finding Inverse

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

General Caesar Cipher

- Assign numerical value from 0 - 25 to each letter of plaintext alphabet *a* - *z*, respectively.

General Caesar Cipher

- Assign numerical value from 0 - 25 to each letter of plaintext alphabet $a - z$, respectively.

- $\mathcal{P} = \mathcal{C} = \mathcal{K} = \mathbb{Z}_{26} = \{0, 1, 2, \dots, 25\}$

General Caesar Cipher

- Assign numerical value from 0 - 25 to each letter of plaintext alphabet $a - z$, respectively.
- $\mathcal{P} = \mathcal{C} = \mathcal{K} = \mathbb{Z}_{26} = \{0, 1, 2, \dots, 25\}$
 \mathbb{Z}_{26} - set of remainders when divide by 26
- Encryption function $E_k : \mathcal{P} \rightarrow \mathcal{C}$ and decryption function $D_k : \mathcal{C} \rightarrow \mathcal{P}$, where $k \in \mathcal{K}$, defined as follows:

General Caesar Cipher

- Assign numerical value from 0 - 25 to each letter of plaintext alphabet $a - z$, respectively.
- $\mathcal{P} = \mathcal{C} = \mathcal{K} = \mathbb{Z}_{26} = \{0, 1, 2, \dots, 25\}$
 \mathbb{Z}_{26} - set of remainders when divide by 26
- Encryption function $E_k : \mathcal{P} \rightarrow \mathcal{C}$ and decryption function $D_k : \mathcal{C} \rightarrow \mathcal{P}$, where $k \in \mathcal{K}$, defined as follows:

$$C = E_k(m) = (m + k) \pmod{26}$$

General Caesar Cipher

- Assign numerical value from 0 - 25 to each letter of plaintext alphabet $a - z$, respectively.
- $\mathcal{P} = \mathcal{C} = \mathcal{K} = \mathbb{Z}_{26} = \{0, 1, 2, \dots, 25\}$
 \mathbb{Z}_{26} - set of remainders when divide by 26
- Encryption function $E_k : \mathcal{P} \rightarrow \mathcal{C}$ and decryption function $D_k : \mathcal{C} \rightarrow \mathcal{P}$, where $k \in \mathcal{K}$, defined as follows:

$$C = E_k(m) = (m + k) \pmod{26}$$

$$m = D_k(C) = (C - k) \pmod{26}$$

General Caesar Cipher

- Assign numerical value from 0 - 25 to each letter of plaintext alphabet $a - z$, respectively.
- $\mathcal{P} = \mathcal{C} = \mathcal{K} = Z_{26} = \{0, 1, 2, \dots, 25\}$
 Z_{26} - set of remainders when divide by 26
- Encryption function $E_k : \mathcal{P} \rightarrow \mathcal{C}$ and decryption function $D_k : \mathcal{C} \rightarrow \mathcal{P}$, where $k \in \mathcal{K}$, defined as follows:

$$C = E_k(m) = (m + k) \pmod{26}$$

$$m = D_k(C) = (C - k) \pmod{26}$$

where $m, C \in Z_{26}$

General Caesar Cipher

- Assign numerical value from 0 - 25 to each letter of plaintext alphabet $a - z$, respectively.
- $\mathcal{P} = \mathcal{C} = \mathcal{K} = \mathbb{Z}_{26} = \{0, 1, 2, \dots, 25\}$
 \mathbb{Z}_{26} - set of remainders when divide by 26
- Encryption function $E_k : \mathcal{P} \rightarrow \mathcal{C}$ and decryption function $D_k : \mathcal{C} \rightarrow \mathcal{P}$, where $k \in \mathcal{K}$, defined as follows:

$$C = E_k(m) = (m + k) \pmod{26}$$

$$m = D_k(C) = (C - k) \pmod{26}$$

where $m, C \in \mathbb{Z}_{26}$

Note that, if key $k = 3$, it is simply a Caesar cipher.

General Caesar Cipher

Example

Find the Caesar cipher of a simple message $m = \text{"crypto"}$ with the key $k = 3$

General Caesar Cipher

Example

Find the Caesar cipher of a simple message $m = \text{"crypto"}$ with the key $k = 3$

- Assume that $m = m_1 m_2 \dots m_n$
the plaintext message with n letters m_1 to m_n

General Caesar Cipher

Example

Find the Caesar cipher of a simple message $m = \text{"crypto"}$ with the key $k = 3$

- Assume that $m = m_1 m_2 \dots m_n$
the plaintext message with n letters m_1 to m_n
- Then $m_1 = c, m_2 = r, m_3 = y, m_4 = p, m_5 = t, m_6 = o$

General Caesar Cipher

Example

Find the Caesar cipher of a simple message $m = \text{"crypto"}$ with the key $k = 3$

- Assume that $m = m_1 m_2 \dots m_n$
the plaintext message with n letters m_1 to m_n
- Then $m_1 = c, m_2 = r, m_3 = y, m_4 = p, m_5 = t, m_6 = o$
- Suppose the corresponding ciphertext letters are C_1 to C_n

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13

plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13

plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = \mathbf{5 = F}$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

$$C_5 = E_k(m_5) = (19 + 3) \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

$$C_5 = E_k(m_5) = (19 + 3) \pmod{26} = 22 \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

$$C_5 = E_k(m_5) = (19 + 3) \pmod{26} = 22 \pmod{26} = 22 = W$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

$$C_5 = E_k(m_5) = (19 + 3) \pmod{26} = 22 \pmod{26} = 22 = W$$

$$C_6 = E_k(m_6) = (14 + 3) \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

$$C_5 = E_k(m_5) = (19 + 3) \pmod{26} = 22 \pmod{26} = 22 = W$$

$$C_6 = E_k(m_6) = (14 + 3) \pmod{26} = 17 \pmod{26}$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13
plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z		
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25		

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

$$C_5 = E_k(m_5) = (19 + 3) \pmod{26} = 22 \pmod{26} = 22 = W$$

$$C_6 = E_k(m_6) = (14 + 3) \pmod{26} = 17 \pmod{26} = 17 = R$$

General Caesar Cipher

plaintext (m)	a	b	c	d	e	f	g	h	i	j	k	l	m	n
Assigned No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13

plaintext (m)	o	p	q	r	s	t	u	v	w	x	y	z
Assigned No.	14	15	16	17	18	19	20	21	22	23	24	25

Encryption algorithm works as follows:

$m = \text{"crypto"}$ and $C_i = E_k(m_i) = (m_i + k) \pmod{26}$

$$C_1 = E_k(m_1) = (2 + 3) \pmod{26} = 5 \pmod{26} = 5 = F$$

$$C_2 = E_k(m_2) = (17 + 3) \pmod{26} = 20 \pmod{26} = 20 = U$$

$$C_3 = E_k(m_3) = (24 + 3) \pmod{26} = 27 \pmod{26} = 1 = B$$

$$C_4 = E_k(m_4) = (15 + 3) \pmod{26} = 18 \pmod{26} = 18 = S$$

$$C_5 = E_k(m_5) = (19 + 3) \pmod{26} = 22 \pmod{26} = 22 = W$$

$$C_6 = E_k(m_6) = (14 + 3) \pmod{26} = 17 \pmod{26} = 17 = R$$

The ciphertext C is **"FUBSWR"**, that is, $E_3(\text{crypto}) = \text{FUBSWR}$

Affine Cipher

Let $\mathcal{P} = \mathcal{C} = \mathbb{Z}_{26}$, and

$$\mathcal{K} = \{(a, b) \in \mathbb{Z}_{26} \times \mathbb{Z}_{26} : \text{GCD}(a, 26) = 1\}$$

$$C = E_k(m) = (am + b) \pmod{26}$$

$$m = D_k(C) = a^{-1}(C - b) \pmod{26}$$

where $m, C \in \mathbb{Z}_{26}$

Affine Cipher

Let $\mathcal{P} = \mathcal{C} = \mathbb{Z}_{26}$, and

$$\mathcal{K} = \{(a, b) \in \mathbb{Z}_{26} \times \mathbb{Z}_{26} : \text{GCD}(a, 26) = 1\}$$

$$C = E_k(m) = (am + b) \pmod{26}$$

$$m = D_k(C) = a^{-1}(C - b) \pmod{26}$$

where $m, C \in \mathbb{Z}_{26}$

Correctness proof:

$$D_k(E_k(m)) = D_k(am + b) \pmod{26}$$

Affine Cipher

Let $\mathcal{P} = \mathcal{C} = \mathbb{Z}_{26}$, and

$$\mathcal{K} = \{(a, b) \in \mathbb{Z}_{26} \times \mathbb{Z}_{26} : \text{GCD}(a, 26) = 1\}$$

$$C = E_k(m) = (am + b) \pmod{26}$$

$$m = D_k(C) = a^{-1}(C - b) \pmod{26}$$

where $m, C \in \mathbb{Z}_{26}$

Correctness proof:

$$\begin{aligned} D_k(E_k(m)) &= D_k(am + b) \pmod{26} \\ &= a^{-1}(\textcolor{red}{am} + \textcolor{red}{b} - b) \pmod{26} \end{aligned}$$

Affine Cipher

Let $\mathcal{P} = \mathcal{C} = \mathbb{Z}_{26}$, and

$$\mathcal{K} = \{(a, b) \in \mathbb{Z}_{26} \times \mathbb{Z}_{26} : \text{GCD}(a, 26) = 1\}$$

$$C = E_k(m) = (am + b) \pmod{26}$$

$$m = D_k(C) = a^{-1}(C - b) \pmod{26}$$

where $m, C \in \mathbb{Z}_{26}$

Correctness proof:

$$\begin{aligned} D_k(E_k(m)) &= D_k(am + b) \pmod{26} \\ &= a^{-1}(\textcolor{red}{am} + \textcolor{red}{b}) - b \pmod{26} \\ &= a^{-1}(am) \pmod{26} \\ &= (a^{-1}a)m \pmod{26} \\ &= m \pmod{26} \\ &= m \end{aligned}$$

Affine Cipher: Correctness

Suppose $k = (7, 3)$, then

- $C = E_k(m) = 7m + 3 \pmod{26}$

Affine Cipher: Correctness

Suppose $k = (7, 3)$, then

- $C = E_k(m) = 7m + 3 \pmod{26}$

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Affine Cipher: Correctness

Suppose $k = (7, 3)$, then

- $C = E_k(m) = 7m + 3 \pmod{26}$

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Reverse Process

$$1 = 5 - (2) \times 2$$

$$1 = 5 - (2) \times [7 - (1) \times 5]$$

$$1 = (-2) \times 7 + (3) \times 5$$

$$1 = (-2) \times 7 + (3) \times [26 - (3) \times 7]$$

$$1 = (-11) \times 7 + (3) \times 26$$

$$x = -11 \pmod{26} = 15$$

Affine Cipher: Correctness

Suppose $k = (7, 3)$, then

- $C = E_k(m) = 7m + 3 \pmod{26}$

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Reverse Process

$$1 = 5 - (2) \times 2$$

$$1 = 5 - (2) \times [7 - (1) \times 5]$$

$$1 = (-2) \times 7 + (3) \times 5$$

$$1 = (-2) \times 7 + (3) \times [26 - (3) \times 7]$$

$$1 = (-11) \times 7 + (3) \times 26$$

$$x = -11 \pmod{26} = 15$$

Correctness

$$D_k(C) = 15(C - 3) \pmod{26}$$

Affine Cipher: Correctness

Suppose $k = (7, 3)$, then

- $C = E_k(m) = 7m + 3 \pmod{26}$

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Reverse Process

$$1 = 5 - (2) \times 2$$

$$1 = 5 - (2) \times [7 - (1) \times 5]$$

$$1 = (-2) \times 7 + (3) \times 5$$

$$1 = (-2) \times 7 + (3) \times [26 - (3) \times 7]$$

$$1 = (-11) \times 7 + (3) \times 26$$

$$x = -11 \pmod{26} = 15$$

Correctness

$$\begin{aligned} D_k(C) &= 15(C - 3) \pmod{26} \\ &= 15([7m + 3] - 3) \pmod{26} \end{aligned}$$

Affine Cipher: Correctness

Suppose $k = (7, 3)$, then

- $C = E_k(m) = 7m + 3 \pmod{26}$

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Reverse Process

$$1 = 5 - (2) \times 2$$

$$1 = 5 - (2) \times [7 - (1) \times 5]$$

$$1 = (-2) \times 7 + (3) \times 5$$

$$1 = (-2) \times 7 + (3) \times [26 - (3) \times 7]$$

$$1 = (-11) \times 7 + (3) \times 26$$

$$x = -11 \pmod{26} = 15$$

Correctness

$$\begin{aligned} D_k(C) &= 15(C - 3) \pmod{26} \\ &= 15([7m + 3] - 3) \pmod{26} \\ &= 105m \pmod{26} \end{aligned}$$

Affine Cipher: Correctness

Suppose $k = (7, 3)$, then

- $C = E_k(m) = 7m + 3 \pmod{26}$

Remainder Form

$$26 = (3) \times 7 + 5$$

$$7 = (1) \times 5 + 2$$

$$5 = (2) \times 2 + 1$$

$$2 = (2) \times 1 + 0$$

Reverse Process

$$1 = 5 - (2) \times 2$$

$$1 = 5 - (2) \times [7 - (1) \times 5]$$

$$1 = (-2) \times 7 + (3) \times 5$$

$$1 = (-2) \times 7 + (3) \times [26 - (3) \times 7]$$

$$1 = (-11) \times 7 + (3) \times 26$$

$$x = -11 \pmod{26} = 15$$

Correctness

$$\begin{aligned} D_k(C) &= 15(C - 3) \pmod{26} \\ &= 15([7m + 3] - 3) \pmod{26} \\ &= 105m \pmod{26} \\ &= m \end{aligned}$$

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext		c	r	y	p	t	o
-----------	--	---	---	---	---	---	---

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext
 m

c	r	y	p	t	o
2					

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17				

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24			

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15		

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext
 m

c	r	y	p	t	o
2	17	24	15	19	14

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$						

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12					

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87				

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122			

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77		

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$						

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12					

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9				

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18			

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25		

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M					

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J				

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S			

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S	Z		

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S	Z	T	

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S	Z	T	U

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S	Z	T	U

That is, $E_K(\text{crypto}) = \text{MJSZTU}$.

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S	Z	T	U

That is, $E_K(\text{crypto}) = \text{MJSZTU}$.

- The decryption function is $m = D_k(C) = 5^{-1}(C - 2) \pmod{26}$

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S	Z	T	U

That is, $E_K(\text{crypto}) = \text{MJSZTU}$.

- The decryption function is $m = D_k(C) = 5^{-1}(C - 2) \pmod{26}$
- We have to find the value of $5^{-1} \pmod{26}$

Affine Cipher: Encryption

Find the Affine cipher for given

- the plaintext message m : “*crypto*”
- key $k = (a, b) = (5, 2)$, then $C = E_k(m) = 5m + 2 \pmod{26}$

Encryption

plaintext	c	r	y	p	t	o
m	2	17	24	15	19	14
$5m + 2$	12	87	122	77	97	72
$(5m + 2) \pmod{26}$	12	9	18	25	19	20
ciphertext	M	J	S	Z	T	U

That is, $E_K(\text{crypto}) = \text{MJSZTU}$.

- The decryption function is $m = D_k(C) = 5^{-1}(C - 2) \pmod{26}$
- We have to find the value of $5^{-1} \pmod{26}$

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$
- $5 = 5 \times 1 + 0$

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$
- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$
- $5 = 5 \times 1 + 0$
- $1 = 5x + 26y$
where $x = -5$ and $y = 1$

Rewrite

- $1 = 26 - 5 \times 5$

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext

C

M

J

S

Z

T

U

12

9

18

25

19

20

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext

C

$21(C - 2)$

M

J

S

Z

T

U

12

9

18

25

19

20

210

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext

C

$21(C - 2)$

M

J

S

Z

T

U

12

9

18

25

19

20

210

147

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext

C

$21(C - 2)$

M

J

S

Z

T

U

12

9

18

25

19

20

210

147

336

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext

C

$21(C - 2)$

M

J

S

Z

T

U

12

9

18

25

19

20

210

147

336

483

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext

C

$21(C - 2)$

M

J

S

Z

T

U

12

9

18

25

19

20

210

147

336

483

357

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext

C

$21(C - 2)$

M

J

S

Z

T

U

12

9

18

25

19

20

210

147

336

483

357

378

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2					

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17				

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24			

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15		

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14
plaintext	c					

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14
plaintext	c	r				

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14
plaintext	c	r	y			

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14
plaintext	c	r	y	p		

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14
plaintext	c	r	y	p	t	

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14
plaintext	c	r	y	p	t	o

Affine Cipher: Decryption

Finding $5^{-1} \pmod{26}$ using Euclidean Algorithm

- $26 = 5 \times 5 + 1$

- $5 = 5 \times 1 + 0$

Rewrite

- $1 = 26 - 5 \times 5$

- $1 = 5x + 26y$

where $x = -5$ and $y = 1$

- $1 = 5x \pmod{26}$, that is,
 $x = 5^{-1} = -5 \pmod{26} = 21$

Decryption

ciphertext	M	J	S	Z	T	U
C	12	9	18	25	19	20
$21(C - 2)$	210	147	336	483	357	378
$21(C - 2) \pmod{26}$	2	17	24	15	19	14
plaintext	c	r	y	p	t	o

Remark: If $a = 1$, the Affine cipher becomes simply a Caesar cipher, that is, $C = E_K(m) = x + b \pmod{26}$.

Thank You