

# Sequence Hunt

As you know in Arithmetic Progression (A.P.) difference between every two consecutive terms remains constant; whereas in Geometric Progression (G.P.) ratio of every two consecutive terms remains same. Consider the example:

2 5 8 11 is A.P. having common difference 3. 2 6 18 54 is G.P. having common ratio of 3.

You have some sequences, where each sequence has multiple sub sequences (either A.P. or G.P.) mixed as elaborated by following examples:

1	2 5 6 8 10 11	First subsequence is an A.P. [2 6 10] Second subsequence is also an A.P. [5 8 11]
2	2 5 10 15 50 45 250 135	First subsequence is a G.P. [2 10 50 250] Second subsequence is also a G.P. [5 15 45 135]
3	2 5 5 10 8 20 11 40	First subsequence is an A.P. [2 5 8 11] Second subsequence is a G.P. [5 10 20 40]

Each sequence of n numbers may have 1 upto n/3 sub sequences because to get common difference or ratio minimum 3 elements are required. Each ith sequence starts from ith location as you have noticed in above examples, it can be observed in Input / Output section. Your task is to analyze each sequence and identify no of subsequences that can be 1 or more. Moreover, you have to find type of each subsequence and elements of individual sub sequence.

## Input:

The first line has integer N (the number of mixed sequences), followed by N lines each contains mixed sequences. Each of the next N lines contains C+1 integers starting with C followed by the mixed sequence of C elements.

## Constraint:

$1 \leq N < 1000$

$3 \leq C < 1000$

Every number is in the range of 4 byte integer.

## Output:

For each sequence output shows count of subsequences, type of each subsequence followed by corresponding subsequence.

Sample Input	Sample Output
4 6 2 5 6 8 10 11 8 2 5 10 15 50 45 250 135 8 2 5 5 10 8 20 11 40 12 2 5 7 5 10 14 8 20 21 11 40 28	Count=2 AP 2 6 10 AP 5 8 11 Count=2 GP 2 10 50 250 GP 5 15 45 135 Count=2 AP 2 5 8 11 GP 5 10 20 40 Count=3 AP 2 5 8 11 GP 5 10 20 40 AP 7 14 21 28

