

KZ equation and correspondence in CS

Haolin Fan

Abstract

A sketch from the basics of the WZW model to the KZ equation and its relation to Chern-Simons theory.

Keywords: keyword WZW model;KZ equation;

1 Frame of WZW model

WZW model is a nonlinear sigma model with the target space being a Lie group,in this sketch,we choose the map from Σ to $SU(2)$ as a model.In fact ,this map describes the gauge transformation over the $SU(2)$ principle bundle and relate to Chern-Simons theory.

1.1 Action and Conservation Current

In WZW model,the action is:

$$S^{\text{WZW}} = \frac{k}{16\pi} \int d^2x \text{Tr}' (\partial^\mu g^{-1} \partial_\mu g) + k\Gamma$$

with

$$\Gamma = \frac{-i}{24\pi} \int_B d^3y \epsilon_{\alpha\beta\gamma} \text{Tr}' (\tilde{g}^{-1} \partial^\alpha \tilde{g} \tilde{g}^{-1} \partial^\beta \tilde{g} \tilde{g}^{-1} \partial^\gamma \tilde{g})$$

the kinetic term is nature,since $\partial^\mu g^{-1} = - (g^{-1} \partial_\mu g)(g^{-1} \partial_\mu g)$ and $U = g^{-1} \partial_\mu g$ is the lie algebra,so it's the contraction of killing form with derivation as common kinetic term.But if calculate its emotion function,we get :

$$\delta \mathcal{L}_t = \frac{\partial \text{Tr}' (U^\dagger U)}{\partial U} \cdot \delta U = \text{Tr}' (\delta g \cdot \partial_\mu U) = \text{Tr}' (\delta g \cdot \partial^\mu (g^{-1} \partial_\mu g))$$

which results in the following equation of motion:

$$\partial^\mu (g^{-1} \partial_\mu g) = 0.$$

but if we translate this to complex coordinate,it is:

$$\partial_z \tilde{J}_{\bar{z}} + \partial_{\bar{z}} \tilde{J}_z = 0.$$

with $\tilde{J}_z = g^{-1} \partial_z g$ and $\tilde{J}_{\bar{z}} = g^{-1} \partial_{\bar{z}} g$. yet in classical CFT,we know that the separation of homology and anti-homology(i.e $\partial_{\bar{z}} T = 0$) plays an important role in OPE, and corresponds to conformal invariance under renormalization. So we introduce the term Γ (it can also be considered as the pull-back of the wedge 3-form of

the Maurer-Cartan form) ,here \tilde{g} means extension g to a 3D-manifold with boundary Σ ,since the extension is not unique , k has quantized to some positive integer.

Check its motion equation,we have: $\partial_z (g^{-1} \partial_{\bar{z}} g) = 0$ and let

$$J_z = -k \partial_z g g^{-1}, \quad J_{\bar{z}} = k g^{-1} \partial_{\bar{z}} g$$

,we get the expected property.

Analog with energy-momentum tensor as conservation current as translation,we define the corresponding infinitesimal transform:

$$\delta_\omega g = \omega g \quad \delta_{\bar{\omega}} g = -g \bar{\omega} \quad (\omega \text{ belongs to lie algebra})$$

check $\delta \mathcal{L}$,this is a local symmetry transformation

$$G(z) \times G(\bar{z})$$

1.2 Affine Lie algebra

We extend J and ω :

$$J = \sum_a J^a t^a, \quad \omega = \sum_a \omega^a t^a$$

by directly calculation

$$\delta_\omega J = -k [\partial_z (\delta_\omega g) g^{-1} - \partial_z g g^{-1} \delta_\omega g^{-1}] = -k (\partial_z \omega g + \omega \partial_z g) g^{-1} + k \partial_z g g^{-1} \omega = [\omega, J] - k \partial_z \omega$$

It can be rewritten as

$$\delta_\omega J^a = \sum_{b,c} f_{abc} \omega^b J^c - k \partial_z \omega^a$$

by Ward identity in integration form,

$$\langle \delta X \rangle = \int dx \partial_\mu (j_a^\mu(x) X) \omega_a(x) = \delta_{\omega, \bar{\omega}} \langle X \rangle = -\frac{1}{2\pi i} \oint dz \omega^a \langle J^a X \rangle$$

by conformal Ward identity,

$$[Q_a, \Phi] = -i \delta_a \Phi$$

we get

$$[J_{n'}^a, J_m^b] = \sum_c i f_{abc} J_{n+m}^c + k n \delta_{ab} \delta_{n+m,0} \quad (\text{here the low index } n \text{ means laurent expansion})$$

it's a affine lie algebra structure denoted as

$$\hat{g} = g \otimes \mathbb{C}[m, m^{-1}] \oplus C\hat{k} \oplus CL_0$$

its irreducible representation can be calculate in wight vector space. For the case of $su(2)$. The generator of lie algebra

$$\begin{aligned} t_1 &= \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ t_2 &= \frac{1}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \\ t_3 &= \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \end{aligned}$$

$$[T_a, T_b] = i\epsilon_{abc}T_c \quad (a, b, c = 1, 2, 3)$$

its H_0^1 chosen as $-t^3 \otimes m^0$ then the affine weight of a state is marked by the eigenvalue of $(H_0^1, \hat{k}, -L_0)$.

We can also define highest weight through primary field, where we extend

$$J^a(z) = \sum_i \frac{J_i^a}{z^i + 1}$$

and define primary field as:

$$J_0^a \phi_\lambda = -t_\lambda^a \phi_\lambda \quad J_n^a \phi_\lambda = 0$$

Here, we claim that state defined by representation is equivalent to primary and secondary field, since for highest state,

$$J_0^a |\lambda\rangle = J_0^a \phi_\lambda |0\rangle = -t_\lambda^a \phi_\lambda |0\rangle = -t_\lambda^a |\lambda\rangle = \lambda |\lambda\rangle$$

the condition $J_n^a \phi_\lambda = 0$ can be proved in similar.

To $\widehat{su}(2)$, its representation is limited by Weyl chambers:




Figure 14.2. Affine Weyl chambers for $\widehat{su}(2)$.

Figure 1: Weyl chambers for $\widehat{su}(2)$

when $k > 0$, all its representations are integrable.

2 Relation to Chern-Simons Theory

The action of Chern-Simons Theory is

$$CS(A) = \frac{k}{4\pi} \int_M \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right),$$

here we use the following model:




Figure 2: Boundary model

here, M is a compact oriented 3-D manifold with boundary Σ . We assume

$$M = \Sigma \times I, I \text{ close in } R, \Sigma = U_1 \cup U_2$$

P is the principal $SU(2)$ bundle over M , A_M is the space of connections on P , \mathcal{G} is the gauge group of P so the corresponding path-integral is:

$$Z_{CS} = \int_{\mathcal{A}_M} e^{iS_{CS}(A)}$$

but for manifolds with boundary, it depends on the configuration of A limit on the boundary, we denote A_1, A_2 , so the path integral in fact is a functional of A_1, A_2 , so we denote \mathcal{A}_Σ , the space of connection of P limit on Σ . Now, the path-integral is

$$Z_{CS}(A_1, A_2) = \int_{A|_\Sigma = (A_1, A_2)} \mathcal{D}A e^{iS_{CS}(A)}$$

a function on \mathcal{A}_Σ . It can be seemed as section of a line bundle over \mathcal{A}_Σ with nature hermitian metric bu fusion rule.

In category language, this means "a 3-D QFT assign to 3-D manifold gives a partition function and assign to 2-D manifold gives a quantum Hilbert space". In particular, since Chern-Simons is topology field theory, and observable depends on gauge field, we should process geometric quantization on moduli space of flat connections on boundary modulo gauge transformations, denoted \mathcal{M} .

To realize Kahler polarization on \mathcal{M} , we select a complex structure to Σ (this also means selecting a conformal structure), and we can construct a symplectic form on \mathcal{M} , denoted the space \mathcal{M}_J and get a Quantum Hilbert space $\mathcal{H}_\Sigma^{(J)}$, and we regard it as a modular functor, since we can construct a flat connection to identify complex structure, at last we get functor

$$\mathcal{H} : \Sigma \rightarrow \mathcal{H}_\Sigma$$

In this sense, if we isomorphic \mathcal{H}_Σ with its dual space. The correlation function can be seen as operator from state on U_1 to state on U_2 .

3 KZ equation

3.1 Sugawara Construction

By directly calculation, the energy-momentum tensor of \mathcal{L}_{wzw} is:

$$T(z) = \frac{k}{2} \text{Tr}(\partial_z g g^{-1} \partial_z g g^{-1}) = \frac{k}{2} \text{Tr}(JJ)$$

and by the orthonormal relation of generator we have

$$T(z) = \frac{k}{2} \sum_a (J^a J^a)$$

but there is a abnormal that

$$J^a(w)T(z) \neq \overline{J^a(w)}(J^b(z)J^b(z))$$

here the overline means the contraction in OPE. So we modify it to

$$T(z) = \frac{1}{2(k+g)} \sum_a (J^a J^a)(z) \quad (g \text{ is the dual Coxeter number relate to structure constant}), \text{ for } SU(2) \quad g = 2$$

and under Laurent extension, we have the KZ Equation, which is a necessary and sufficient condition to highest weight state:

$$L_{-1}|\varphi_i\rangle = \frac{1}{k+g} \sum_a (J_{-1}^a J_0^a) |\varphi_i\rangle = -\frac{1}{k+g} \sum_a (J_{-1}^a t_i^a) |\varphi_i\rangle$$

We consider the insertion of the zero vector

$$|x\rangle = \left[L_{-1} + \frac{1}{k+g} \sum_a (J_{-1}^a t_i^a) \right] |\varphi_i\rangle = 0 \quad (15.96)$$

inside the correlation function of a set of primary fields. We note that the insertion of the operator J_{-1}^a in the correlator can be expressed through Laurent extension at each z_i

$$\begin{aligned} \langle \varphi_1(z_1) \cdots (J_{-1}^a \varphi_i)(z_i) \cdots \varphi_n(z_n) \rangle &= \frac{1}{2\pi i} \oint_{z_i} \frac{dz}{z - z_i} \langle J^a(z) \varphi_1(z_1) \cdots \varphi_n(z_n) \rangle \\ &= \frac{1}{2\pi i} \oint_{z_{j \neq i}} \frac{dz}{z - z_i} \sum_{j \neq i} \frac{t_j^a}{z - z_j} (\varphi_1(z_1) \cdots \varphi_n(z_n)) \\ &= \sum_{j \neq i} \frac{t_j^a}{z_i - z_j} \langle \varphi_1(z_1) \cdots \varphi_n(z_n) \rangle \end{aligned}$$

At last we have:

$$\langle \varphi_1(z_1) \cdots \chi(z_i) \cdots \varphi_n(z_n) \rangle = \left[\partial_{z_i} + \frac{1}{k+g} \sum_{j \neq i} \frac{\sum_a t_i^a \otimes t_j^a}{z_i - z_j} \right] \langle \varphi_1(z_1) \cdots \varphi_n(z_n) \rangle = 0$$

this is KZ equation for conformal block and also necessary and sufficient

3.2 KZ connection

To relate with Chern-Simons, we give a geometry description of n-point correlation function.

We first divide n-point correlation function to conformal block by operator algebra:

$$\langle \varphi_1(x_1) \cdots \varphi_i(x_i) \rangle = \sum_{p,q} C_{p,q} F_p(x_i) \bar{F}_q(\bar{x}_i),$$

so the we get the KZ equation for conformal block:

$$\frac{\partial F}{\partial z_i} = \frac{1}{k+h^\vee} \sum_{j \neq i} \frac{\sum_a t_i^a \otimes t_j^a}{z_i - z_j} F,$$

here we can write the correlation function as $\langle \varphi_1(x_1) \cdots \varphi_i(x_i) \rangle = \langle 0 | \lambda_1 \cdots \lambda_i \rangle$, where we use the equivalent between primary field with state, in this case conformal block can be seen as a "channel" from $|\lambda_1 \cdots \lambda_i\rangle$ go through integrable weight $|p\rangle$ to 0. So we can define conformal block as:

$$\Psi : H_{\lambda_1} \otimes \cdots \otimes H_{\lambda_n} \rightarrow \mathbb{C}$$

and the space of all the "channel" as $\mathcal{H}(p_1, \cdots, p_n; \lambda_1, \cdots, \lambda_n)$.

Because of the dependence on position and state, we define

$$\mathcal{E}_{\lambda_1, \dots, \lambda_n} = \bigcup_{(p_1, \dots, p_n) \in \text{Conf}_n(\Sigma)} \mathcal{H}(p_1, \dots, p_n; \lambda_1, \dots, \lambda_n)$$

as a subbundle of E, which defined as:

$$E = \text{Conf}_n(\mathbb{CP}^1) \times \text{Hom}_{\mathbb{C}} \left(\bigotimes_{j=1}^n H_{\lambda_j}, \mathbb{C} \right)$$

is a vector bundle over $\text{Conf}_n(\Sigma)$. To select the subbundle, we define KZ connection as:

$$\nabla_{\frac{\partial}{\partial z_j}} \Psi = \frac{\partial \Psi}{\partial z_j} - L_{(j)-1} \Psi.$$

by KZ equation, we realize conformal block as horizontal section of E. If we regard $\mathcal{H}(p_1, \cdots, p_n; \lambda_1, \cdots, \lambda_n)$ as functor, in fact it's the same with the functor \mathcal{H} (last part of section 2) act on Σ equip with wilson line[2].

3.3 Solution to four point conformal block on SU(2)

Here we choose SU(2), K=1 to show the solution of conformal block. In K=1 case, by Weyl chambers there are only two holomorphic conformal block, denoted F_0 and F_1 , corresponds to 1 and 3 dimension integrable representation. Since conformal block satisfy conformal invariance, it must have this form:

$$F(z_i) = [(z_1 - z_4)(z_2 - z_3)(\bar{z}_1 - \bar{z}_4)(\bar{z}_2 - \bar{z}_3)]^{-2h} F(x)$$

where h is the conformal dimension and

$$x = \frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_4)(z_3 - z_2)}$$

In the following, we set $z_{ij} = z_i - z_j$. so the Knizhnik-Zamolodchikov equation (15.99) becomes

$$\left(\partial_i + \frac{1}{k+N} \sum_{j \neq i} t_i^a \otimes t_j^a \frac{1}{z_i - z_j} \right) [z_{14} z_{23}]^{-2h} F = 0$$

The next step is to transform these partial differential equations into an ordinary differential equation in the variable x . Consider the case $i = 1$; since

$$\partial_1 = \left(\frac{x}{z_{12}} - \frac{x}{z_{14}} \right) \partial_x$$

the equation takes the form

$$\left(-\frac{2h}{z_{14}} + \left(\frac{x}{z_{12}} - \frac{x}{z_{14}} \right) \partial_x + \frac{1}{k+N} \sum_{j=2,3,4} \frac{\sum_a t_j^a \otimes t_j^a}{z_1 - z_j} \right) F = 0$$

Then we choose:

$$z_1 = x, \quad z_2 = 0, \quad z_3 = 1, \quad z_4 = \infty$$

The last equation is:

$$\left\{ \partial_x + \frac{1}{k+N} \sum_a t_1^a \otimes t_2^a \frac{1}{x} + \frac{1}{k+N} \sum_a t_1^a \otimes t_3^a \frac{1}{x-1} \right\} F = 0$$

so we have to determine the representation of tensor of generator on the singlet and triplet state, here we use Casimir Operator:

$$C = \sum_a J^a J^a$$

it's the unique commute with all the generators for SU(2). And for two representations of the tensor product $V_{j_1} \otimes V_{j_2}$, the total Casimir operator is given by:

$$C_{\text{total}} = \sum_a (t_1^a + t_2^a)^2 = C_1 + C_2 + 2 \sum_a t_1^a t_2^a.$$

Thus, the generator factor can be represented as:

$$\sum_a t_1^a \otimes t_2^a = \frac{1}{2} (C_{\text{total}} - C_1 - C_2).$$

for the case the weight of $\phi_1, \phi_2, \phi_3, \phi_4$ is $(1/2, 1/2, 1/2, 1/2)$

$$C = (\lambda, \lambda + 2\rho)$$

λ is weight and ρ is root. So, $C_1 = -\frac{3}{4} = C_2$

For singlet state:

$$\sum_a t_1^a \otimes t_2^a = -\frac{3}{4} \quad \sum_a t_1^a \otimes t_3^a = -\frac{3}{4}$$

here to calculate cross term we have use :

$$\sum_a t_{ij}^a t_{kl}^a = \frac{1}{2} \delta_{il} \delta_{jk} - \frac{1}{2N} \delta_{ij} \delta_{kl}.$$

and notice the first term is permutation which is (-1) for singlet the later is id. now the equation is:

$$\partial_x F_0(x) = -\frac{1}{4} \left(\frac{1}{x} + \frac{1}{x-1} \right) F_0(x),$$

with solution:

$$\mathcal{F}_0(x) = C \cdot x^{-\frac{1}{4}} (1-x)^{-\frac{1}{4}}.$$

Similar, for triplet state,

$$\sum_a t_1^a \otimes t_2^a = \frac{1}{4} \quad \sum_a t_1^a \otimes t_3^a = \frac{1}{4}$$

the solution is:

$$\mathcal{F}_1(x) = C \cdot x^{\frac{1}{4}} (1-x)^{\frac{1}{4}}.$$

References

- [1] Arnaud Beauville, Yves Laszlo, "Conformal blocks and generalized theta functions", *Communications in Mathematical Physics*, Volume 164, Issue 2, pp. 385-419.
- [2] Witten, E. Quantum field theory and the Jones polynomial. *Commun.Math. Phys.* 121, 351–399 (1989).