

SECOND LIVES CORPORATION

DIVISION 9 BIOSECURITY COMMAND

COMPREHENSIVE PARASITOLOGY DOSSIER

CLASSIFICATION: OMEGA BLACK



SecondLives

Entering the second life with your loved ones

Table of Contents

Executive Summary.....	1
Executive Summary.....	2
PARASITE CLASSIFICATION SYSTEM.....	3
PARASITE CLASSIFICATION SYSTEM.....	4
PARASITE CLASSIFICATION SYSTEM.....	5
PARASITE CLASSIFICATION SYSTEM.....	6
PARASITE CLASSIFICATION SYSTEM.....	7
PARASITE CLASSIFICATION SYSTEM.....	8
PARASITE CLASSIFICATION SYSTEM.....	9
PARASITE CLASSIFICATION SYSTEM.....	10
PARASITE CLASSIFICATION SYSTEM.....	11
PARASITE CLASSIFICATION SYSTEM.....	12
Infection Progression Analysis.....	13
Infection Progression Analysis.....	14
Infection Progression Analysis.....	15
Infection Progression Analysis.....	16
Infection Progression Analysis.....	17
Specimen 07-A (Patient Zero).....	18
The Butcher.....	19
The Screamer.....	20
Sporestrider.....	21
Blindmaw.....	22
Halfling.....	23
The Warden.....	24
The Warden.....	25
Conclusion.....	26

Executive Summary

Following the catastrophic Nova Crest Outbreak of 2010, all surviving parasite specimens, infected biological materials, research archives, and containment assets were seized and transferred under the authority of Division 9's Department of Advanced Warfare. What began as a civilian biotechnology initiative under Second Lives Corporation rapidly evolved into one of the most significant biological containment crises in modern history, resulting in the collapse of multiple research sectors, the loss of thousands of civilian and scientific personnel, and the permanent quarantine of Nova Crest City.

Subsequent investigations confirmed the existence of multiple genetically divergent parasite strains originating from a common source organism designated ***Specimen 07-A (Patient Zero)***. Analysis revealed that these organisms possessed unprecedented adaptive capabilities, including accelerated mutation rates, host integration mechanisms, neural infiltration properties, and the capacity to develop specialized evolutionary branches under environmental pressure. As a result, conventional biological classification methods proved insufficient for accurate threat assessment and containment planning.

In response, Division 9 established the **Parasitology Division**, a specialized research and containment program tasked with the identification, cataloging, observation, and control of all known parasite variants. Since its formation, the division has conducted thousands of observation hours, recovered numerous infected specimens, documented multiple mutation events, and developed the first standardized threat classification system for parasite-related entities.

This dossier serves as the primary operational reference for authorized research personnel, containment officers, military response teams, and executive oversight staff. Its purpose is to provide consolidated intelligence regarding parasite morphology, behavioral patterns, infection vectors, environmental adaptations, containment requirements, and emergency response procedures. Information contained herein is derived from direct observation logs, recovered laboratory records, field reports, autopsy examinations, neural scan data, and post-incident investigations conducted between 2010 and the present day.

Current records confirm the existence of multiple high-risk strains, including breeder-class organisms, acoustic predators, spore-dispersal variants, infiltration entities capable of human mimicry, and sovereign-level hybrid organisms exhibiting intelligence and adaptive capabilities beyond initial projections. Several documented specimens have demonstrated behaviors previously believed impossible, including coordinated hunting tactics, environmental manipulation, regenerative adaptation, and signs of emerging collective intelligence.

Personnel are advised that all parasite specimens must be considered capable of continued evolution. Historical data demonstrates that no strain remains biologically static for

extended periods, and mutation events may occur without warning. Threat classifications provided within this document represent the most current assessments available at the time of publication and are subject to revision following future observations.

The information contained within this dossier is classified under **OMEGA BLACK SECURITY CLEARANCE**. Unauthorized access, duplication, distribution, or disclosure of any material contained herein is strictly prohibited and may result in immediate disciplinary action, security review, or detention under Division 9 Biosecurity Directives.

Continued observation, documentation, and containment remain essential to preventing a second Nova Crest-scale event. All personnel are reminded that preservation of scientific data is considered secondary only to successful containment. In the event of a containment failure, emergency response procedures outlined in Appendix D are to be enacted immediately.

PARASITE CLASSIFICATION SYSTEM

Biological Threat Taxonomy

Following the Nova Crest Outbreak, there was established a standardized classification framework to categorize parasite specimens based on origin, behavior, mutation potential, infection methods, intelligence, and containment requirements. This system allows researchers and response teams to rapidly assess threats and determine appropriate containment procedures.

The classifications listed below represent the current understanding of parasite evolution and are subject to revision as new strains emerge.

Level	Designation	Meaning
I	Minimal	Easily contained, low danger
II	Elevated	Dangerous but manageable
III	Severe	Significant risk to personnel
IV	Critical	High mortality and containment risk
V	Catastrophic	Facility-wide threat
VI	Extinction	Regional or civilization-level threat
Ω	Omega Black	Existential threat; military intervention mandatory

PARASITE CLASSIFICATION SYSTEM

The following designations represent the scientific taxonomy assigned to the parasitic organisms discovered and studied by Second Lives Corporation. While individual infected hosts may vary significantly in appearance, each classification refers to a distinct evolutionary branch of parasite worm exhibiting unique biological adaptations, reproductive methods, and host interaction mechanisms.

All currently documented strains are believed to descend from a common ancestor originating from Specimen 07-A.

Classified

PARASITE CLASSIFICATION SYSTEM

PRIME ORIGIN STRAIN

Taxonomic Designation: Prime Origin Strain

Description: The Prime Origin Strain is considered the ancestral form from which all known parasite species evolved. Unlike later specialized variants, the Prime Strain possesses an unusually flexible genome capable of adapting to nearly any host environment. Researchers believe this strain was responsible for the earliest infections observed during pre-Nova Crest testing. Genetic sequencing indicates that every documented parasite contains fragments of Prime-Strain DNA.

Worm Morphology

- Length: 12–40 cm
- Coloration: Pale white to translucent gray
- Body Structure: Smooth-bodied nematode-like organism
- Nervous System: Highly developed neural fibers
- Reproductive Method: Asexual cellular replication

Primary Function: The Prime Strain serves as a biological foundation for future evolutionary branches, allowing adaptation into specialized forms when exposed to environmental pressures.

Research Notes: often refer to the Prime Strain as "the blueprint."



PARASITE CLASSIFICATION SYSTEM

PROGENIVORAX GESTATRIX

Translation: "Life-Devouring Bearer"

Classification: Breeder-Class Parasitic Worm

Description: Progenivorax Gestatrix evolved to maximize reproduction and population growth. Unlike most strains, it invests the majority of its biological resources into generating offspring rather than mobility or stealth.

These worms act as living incubation systems capable of producing hundreds of juvenile organisms throughout their lifespan.

Worm Morphology:

- Length: 40–120 cm
- Diameter: Extremely thick-bodied
- Coloration: Dark red to black
- Surface Texture: Segmented muscular rings
- Internal Structure: Multiple gestation chambers

Biological Function:

- Reproduction
- Colony expansion
- Host conversion
- Nest creation

Research Notes: Researchers frequently compare this species to parasitic queen insects due to its reproductive specialization.



PARASITE CLASSIFICATION SYSTEM

SONOSPORA VOXALIS

Translation: "Sound-Spore Voice"

Classification: Acoustic Predator Worm

Description: Sonospora Voxalis evolved sensory organs capable of detecting microscopic sound vibrations. The species possesses reduced visual organs but compensates with extraordinary auditory perception.

These worms can generate resonant frequencies used for navigation, communication, and host detection.

Worm Morphology

- Length: 20–60 cm
- Coloration: Pale gray
- Sensory Structures: Vibrational receptor nodes
- Eyes: Vestigial or absent
- Vocal Organs: Resonance sacs along the body

Biological Function

- Host tracking
- Environmental mapping
- Colony communication
- Predatory behavior

Research Notes: This species appears to navigate similarly to bats through biological echolocation.



PARASITE CLASSIFICATION SYSTEM

FISSIONIS ECHOFORMIS

Translation: "Dividing Echo Form"

Classification: Spore-Dispersal Worm

Description: Fissionis Echoformis evolved for environmental contamination. Specialized internal organs continuously generate infectious spores that are released through respiratory vents and tissue ruptures.

The species demonstrates rapid movement and exceptional environmental survivability.

Worm Morphology

- Length: 15–45 cm
- Coloration: Green-gray
- Surface Texture: Covered in microscopic spore sacs
- Distinctive Feature: Bioluminescent fungal growths

Biological Function

- Environmental infection
- Spore dispersal
- Territory expansion
- Long-term contamination

Research Notes: A single mature specimen can contaminate several square kilometers under favorable conditions.



PARASITE CLASSIFICATION SYSTEM

SONOSPORA / FISSIONIS HYBRID

Classification: Hybridized Predator-Dispersal Worm

Description: This strain represents an unstable merger between Sonospora Voxalis and Fissionis Echoformis genetic material.

The resulting organism combines the sensory capabilities of Sonospora with the contamination abilities of Fissionis.

Worm Morphology

- Length: 30–90 cm
- Coloration: Dark gray with green bioluminescence
- Specialized Structures:
 - Echolocation nodes
 - Spore glands
 - Reinforced musculature

Biological Function

- Active hunting
- Environmental infection
- Adaptive mutation
- Territorial dominance

Research Notes: Researchers consider this hybrid one of the most dangerous naturally occurring evolutionary branches due to its versatility.



PARASITE CLASSIFICATION SYSTEM

MIMICUS REPLICANTUS

Translation: "Imitating Replica"

Classification: Infiltration Worm

Description: Mimicus Replicantus evolved to interact directly with host neural tissue. Rather than focusing on physical destruction, the species specializes in observation, behavioral replication, and identity mimicry.

Neural scans suggest these worms partially record host memories and behavioral patterns, allowing infected organisms to imitate human mannerisms with alarming accuracy.

Worm Morphology

- Length: 10–25 cm
- Coloration: Nearly translucent
- Neural Appendages: Extensive
- Body Structure: Extremely thin and flexible

Biological Function


- Host infiltration
- Memory acquisition
- Behavioral replication
- Population concealment

Research Notes: Division 9 personnel classify Mimicus Replicantus as one of the most strategically dangerous parasite species due to its ability to compromise security systems through deception rather than force.



PARASITE CLASSIFICATION SYSTEM

Specimen threat assessment



Note: comparative threat analysis of known parasite classifications.

Threat Assessment Summary

Analysis of current Division 9 threat metrics indicates a significant increase in operational danger among later-generation parasite strains. While early variants such as "The Screamer" demonstrate dangerous predatory behavior, their threat profile remains relatively localized due to limited adaptability and predictable hunting patterns.

"The Butcher" presents a substantially greater risk due to its reproductive specialization. Although less intelligent than several advanced strains, its ability to generate large populations of offspring dramatically increases outbreak potential and containment complexity.

"Sporestrider" ranks above most conventional strains due to its environmental contamination capabilities. Division 9 simulations suggest that a single uncontrolled specimen could compromise an entire facility through airborne spore dispersal within a matter of days.

Hybrid organisms represent a major escalation in biological danger. **Blindmaw**, a Sonospora/Fissionis hybrid, combines predatory efficiency with contamination capabilities,

resulting in a threat profile that exceeds either parent species. Extensive regenerative capabilities further complicate containment and termination efforts.

Although not physically dominant, **Mimicus Replicantus** ("Halfling") is classified as one of the most strategically dangerous organisms ever documented. Its ability to replicate human behavior, infiltrate secured facilities, and compromise personnel identity verification systems creates risks extending beyond traditional biological containment scenarios.

Specimen 07-A (Patient Zero) remains one of the highest-priority assets under Division 9 oversight. As the ancestral source of all known parasite lineages, it possesses extraordinary adaptive potential and remains capable of producing previously undocumented mutations.

The highest threat designation is assigned to **The Warden**, a sovereign-class hybrid organism exhibiting advanced intelligence, regenerative adaptation, and traits inherited from multiple evolutionary branches. Current assessments indicate that containment failure involving The Warden would likely exceed the response capabilities of standard facilities and require immediate military intervention.

Conclusion

Current data demonstrates a clear evolutionary trend toward increased adaptability, intelligence, and survivability among successive parasite generations. While individual strains possess unique biological advantages, sovereign-class hybrids represent the greatest long-term threat to containment operations due to their capacity to combine multiple evolutionary traits within a single organism. Therefore highly recommended to prioritize hybrid and sovereign-class specimens for continuous monitoring and enhanced containment protocols.

Infection Progression Analysis


The progression of parasite infection remains one of the most extensively studied phenomena within Division 9. Data collected from controlled laboratory observations and recovered Nova Crest incident records indicate that infection follows a predictable sequence of neurological and physiological changes. While progression speed may vary between strains, the overall pattern remains remarkably consistent.

Researchers believe the parasite's primary objective is the establishment of neural control over the host. Once successful neural integration occurs, recovery rates decline dramatically and host autonomy becomes increasingly compromised.

Infection Progression Timeline

Parasite infection progression

Observed progression from initial exposure to terminal conversion.



Analysis of infection progression demonstrates a gradual increase in parasite influence during the first twenty-four hours, followed by a period of accelerated host deterioration. The most significant increase occurs during the transition between Neural Integration and Behavioral Alteration, where neurological disruption becomes visible through measurable personality changes, memory impairment, and increased aggression.

researchers refer to this period as the **Critical Intervention Window**, as successful treatment outcomes decrease rapidly beyond this stage.

Infection Progression Analysis


Confirmed Infection Vectors

Current evidence suggests that parasite transmission can occur through multiple pathways. While direct implantation remains the most common method observed in containment incidents, environmental contamination events have become increasingly common among advanced strains.

The following chart represents the distribution of confirmed infection events documented.

Observed infection vectors

Relative frequency of documented infection routes.



Results indicate that nearly half of all documented infections originated from direct parasite implantation. However, the growing prevalence of spore-producing strains has resulted in a substantial increase in airborne transmission events. Such infections are particularly dangerous because exposure may occur without the host's immediate awareness.

Environmental contamination incidents involving airborne spores remain a major concern for containment personnel, as entire facility sectors may become compromised before detection systems identify the threat.

Infection Progression Analysis


Host Survival Analysis

One of primary objectives is determining how infection timing influences recovery outcomes. Long-term studies have consistently demonstrated that survival probability decreases in direct proportion to the parasite's level of neurological integration.

The following chart illustrates estimated survival probabilities based on documented treatment attempts and recovered medical records.

Host survival probability

Estimated survival likelihood following confirmed infection.



The decline in survival probability after the first twenty-four hours is particularly concerning. Statistical models suggest that once neural integration has been established, parasite removal procedures frequently result in catastrophic nervous system damage.

After seventy-two hours, successful recovery becomes exceptionally rare. At this stage, most hosts exhibit extensive biological conversion and are no longer considered medically stable.

As a result, Division 9 policy mandates immediate quarantine and medical evaluation following any suspected exposure event.

Infection Progression Analysis

Symptom Development Matrix

While infection severity increases steadily over time, symptom presentation varies considerably between stages. Early symptoms often resemble common illnesses, making initial detection difficult. More advanced stages produce increasingly recognizable neurological and physiological abnormalities.

Stage	Neurological Effects	Physical Effects	Threat Rating
Exposure	None	Minor irritation	Low
Neural Contact	Headaches, dizziness	Elevated temperature	Moderate
Neural Integration	Memory disruption	Fatigue	High
Behavioral Alteration	Aggression, hallucinations	Increased strength	Severe
Biological Conversion	Loss of judgment	Visible mutation	Critical
Terminal Conversion	Host autonomy lost	Full parasite traits	Catastrophic

Researchers are advised to pay particular attention to behavioral alterations, as these symptoms often provide the first reliable indication of a developing infection. Auditory hallucinations, unexplained aggression, memory lapses, and sudden personality changes have been identified as common warning signs across nearly all documented strains.

Infection Progression Analysis

Research Conclusion

Collected data strongly supports the hypothesis that parasite survival depends primarily upon successful neurological integration. Once control over the host nervous system has been established, both mortality rates and mutation frequency increase substantially.

Continued monitoring of infection pathways, progression timelines, and survival outcomes remains essential for improving containment strategies and reducing casualty rates during future outbreak events.

Specimen 07-A (Patient Zero)

SPECIMEN 07-A : "PATIENT ZERO"

Classification : Prime Origin Strain

Origin : The first known human-parasite integration event following the 1967 Catalyst Incident. All documented parasite variants originate from tissue, spores, or genetic descendants of Specimen 07-A.

Physical Description

- Human-like morphology heavily altered by parasitic growth
- Extensive neural infiltration
- Regenerative tissue layers
- Variable body mass due to adaptive biomass production

Behavioral Profile

- Alternates between aggression and periods of lucidity
- Demonstrates memory retention from original host
- Reacts strongly to familiar voices and environmental triggers
- Displays leadership influence over lesser strains

Observation Equipment

- Neural activity monitor
- Thermal imaging system
- Remote bioscanner
- High-resolution optical systems
- Automated containment cameras

Research Warnings

- Direct exposure prohibited
- Capable of rapid tissue regeneration
- Potential psychic/neural influence on personnel
- Unknown mutation potential

Interaction Procedures

- Maintain minimum 20-meter separation.
- Use remote communication systems only.
- Avoid emotional or personal dialogue.
- Record all speech patterns.
- Immediately report abnormal behavioral changes.

Threat Level : V — Catastrophic

The Butcher

THE BUTCHER

Classification : PROGENIVORAX GESTATRIX - Host-Breeder Parasite

Description : Large, heavily muscled parasite designed for reproduction and host conversion.

Characteristics

- Massive physical strength
- Releases offspring worms
- Creates parasite nests
- High durability

Observation Equipment

- Structural stress monitors
- Thermal cameras
- Drone observation units
- Containment pressure sensors

Warnings

- Do not approach offspring clusters.
- Breeding events may occur without warning.
- Physical barriers may fail under sustained assault.

Interaction Procedures

- Observe remotely.
- Terminate offspring immediately if containment is breached.
- Avoid enclosed environments.

Threat Level : IV — Critical

The Screamer

THE SCREAMER

Classification : Acoustic Predator - SONOSPORA VOXALIS

Description : Blind parasite relying entirely on sound-based navigation.

Characteristics

- No functional eyesight
- Extreme hearing sensitivity
- Generates sonic shockwaves
- Tracks movement through vibration
- Can alert all other parasites to the exact

Observation Equipment

- Sound isolation chamber
- Vibration monitors
- Acoustic spectrum analyzers

Warnings

- Speaking is prohibited near containment.
- Sudden noises may trigger aggression.
- Sonic attacks may cause internal injuries.

Interaction Procedures

- Communicate through text systems only.
- Wear hearing protection.
- Move slowly and quietly.
- Avoid metallic impacts.

Threat Level : III — Severe

Sporestrider

SPORESTRIDER

Classification: Spore Dispersal Variant - FISSIONIS ECHOFORMIS

Description : Fast-moving parasite specializing in environmental contamination.

Characteristics

- Rapid movement speed
- Produces toxic spore clouds
- Thrives in humid environments
- Creates secondary infection zones

Observation Equipment

- Air quality monitors
- Spore detectors
- UV contamination scanners
- Sealed observation systems

Warnings

- Respiratory exposure may result in infection.
- Air filtration systems must remain active.
- Protective suits required.

Interaction Procedures

- Never enter contaminated zones without PPE.
- Perform air testing every 15 minutes.
- Isolate contaminated samples immediately.

Threat Level : IV — Critical

Blindmaw

BLINDMAW

Classification: Hybrid Variant - SONOSPORA / FISSIONIS HYBRID

Description : Engineered combination of Screamer and Sporestrider traits.

Characteristics

- Echolocation capabilities
- Rapid tissue regeneration
- Acidic blood
- Enhanced aggression

Observation Equipment

- Acid-resistant containment systems
- Thermal cameras
- Environmental hazard sensors

Warnings

- Blood contact causes severe chemical burns.
- Regeneration may repair extensive damage.
- Hybrid mutations remain unpredictable.

Interaction Procedures

- Avoid damaging containment unnecessarily.
- Use chemical-resistant equipment.
- Monitor regenerative cycles continuously.

Threat Level: V — Catastrophic

Halfling

HALFLING

Classification: Infiltration Parasite - MIMICUS REPLICANTUS

Description : Parasite capable of generating biological replicas of human subjects.

Characteristics

- Mimics human appearance
- Copies behavioral patterns
- Produces deceptive vocalizations
- Decomposes into spores upon death

Observation Equipment

- DNA verification systems
- Biometric scanners
- Neural activity readers

Warnings

- Visual identification is unreliable.
- Potential infiltration of research staff.
- Spores released after termination.

Interaction Procedures

- Verify identity through DNA scans.
- Require multi-factor biometric confirmation.
- Treat all unknown personnel as potential mimics.

Threat Level : V — Catastrophic

The Warden

THE WARDEN

Classification: Apex Hybrid Organism - PARASITE SOVEREIGN

Description : Largest and most dangerous parasite created through multiple strain integrations.

Phase One: Jailor of Flesh

➤ **Characteristics:**

- Massive biomass
- Heavy armor-like tissue
- Slow movement
- Extreme durability

Phase Two: Parasite Sovereign

➤ **Characteristics:**

- Increased speed
- Accelerated regeneration
- Aggressive combat behavior
- Enhanced intelligence

Observation Equipment

- Satellite-linked monitoring systems
- Military-grade sensor arrays
- Structural integrity monitoring
- Full-spectrum surveillance

Warnings

- No direct interaction authorized.
- Containment failure is considered catastrophic.
- Military intervention is required immediately.

Interaction Procedures

- Evacuate personnel.
- Lock down all facility sectors.
- Deploy emergency response teams.
- Notify Division 9 Command.
- Preserve observation data during evacuation.

The Warden

The Warden (Phase One)

Threat Level: V — Catastrophic

Reason:

- Massive biomass
- Extremely durable
- Requires coordinated response

The Warden (Phase Two — Parasite Sovereign)

Threat Level: Ω — Omega Black

Reason:

- Apex hybrid organism
- Rapid regeneration
- High intelligence
- Multi-strain integration
- Capable of overwhelming military containment assets

Conclusion

Final Assessment

The observations, analyses, and specimen examinations contained within this dossier represent the most complete understanding currently available regarding the parasite classifications descended from the Prime Origin Strain. Despite decades of study, numerous aspects of their biology remain poorly understood, and several documented behaviors continue to challenge established scientific models.

Research confirms that these organisms are not static lifeforms. Every classification examined demonstrates some degree of adaptation, specialization, or mutation. While certain strains evolved primarily for reproduction, communication, environmental spread, or infiltration, all share a common characteristic: the ability to change in response to environmental pressures.

Particularly concerning is the continued emergence of hybridized organisms. The successful integration of multiple evolutionary traits into a single specimen suggests that parasite development is ongoing and may eventually produce classifications beyond current predictive models. Existing threat assessments should therefore be regarded as temporary evaluations rather than permanent designations.

Analysis of all available data supports the conclusion that complete eradication remains unlikely with current technology. Containment, observation, and early detection continue to be the most effective methods of limiting spread and preventing large-scale incidents.

Future research should prioritize:

- Monitoring mutation events.
- Tracking evolutionary divergence.
- Identifying reproductive vulnerabilities.
- Improving early infection detection methods.
- Investigating long-term adaptation mechanisms.

The information contained within this dossier reflects only what has been observed to date. Numerous specimens remain unaccounted for, and several recorded anomalies have yet to receive formal classification.

As such, Division 9 recommends that all parasite classifications be considered active biological threats regardless of current containment status.

END OF DOSSIER

**SECOND LIVES CORPORATION
DIVISION 9 PARASITOLOGY DIVISION
OMEGA BLACK CLEARANCE REQUIRED**